

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
11 January 2001 (11.01.2001)

PCT

(10) International Publication Number  
**WO 01/02583 A2**

(51) International Patent Classification<sup>7</sup>: C12N 15/54,  
15/31, 9/00, 9/12, 15/77, C12P 1/04

(DE). HABERHAUER, Gregor; Moselstrasse 42,  
D-67117 Limburgerhof (DE).

(21) International Application Number: PCT/IB00/00973

(22) International Filing Date: 27 June 2000 (27.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/142,691 1 July 1999 (01.07.1999) US  
60/150,310 23 August 1999 (23.08.1999) US  
199 42 095.5 3 September 1999 (03.09.1999) DE  
199 42 097.1 3 September 1999 (03.09.1999) DE

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: BASF AKTIENGESELLSCHAFT  
[DE/DE]; D-67056 Ludwigshafen (DE).

**Published:**

— Without international search report and to be republished upon receipt of that report.

(72) Inventors: POMPEJUS, Markus; Wenjenstrasse 21, D-67251 Freinsheim (DE). KRÖGER, Burkhard; Im Waldhof 1, D-67117 Limburgerhof (DE). SCHRÖDER, Hartwig; Goethestrasse 5, D-69226 Nussloch (DE). ZELDER, Oskar; Rossmarktstrasse 27, D-67346 Speyer

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: *ORYNEBACTERIUM GLUTAMICUM* GENES ENCODING PHOSPHOENOLPYRUVATE: SUGAR PHOSPHOTRANSFERASE SYSTEM PROTEINS

(57) Abstract: Isolated nucleic acid molecules, designated PTS nucleic acid molecules, which encode novel PTS proteins from *Corynebacterium glutamicum* are described. The invention also provides antisense nucleic acid molecules, recombinant expression vectors containing PTS nucleic acid molecules, and host cells into which the expression vectors have been introduced. The invention still further provides isolated PTS proteins, mutated PTS proteins, fusion proteins, antigenic peptides and methods for the improvement of production of a desired compound from *C. glutamicum* based on genetic engineering of PTS genes in this organism.

WO 01/02583 A2

**CORYNEBACTERIUM GLUTAMICUM GENES ENCODING  
PHOSPHOENOLPYRUVATE: SUGAR PHOSPHOTRANSFERASE SYSTEM  
PROTEINS**

**5    Related Applications**

          This application claims priority to U.S. Provisional Patent Application No.: 60/142,691, filed on July 1, 1999, and also to U.S. Provisional Patent Application No.: 60/150,310, filed on August 23, 1999, incorporated herein in their entirety by this reference. This application also claims priority to German Patent Application No.:  
10    19942095.5, filed on September 3, 1999, and also to German Patent Application No.: 19942097.1, filed on September 3, 1999, incorporated herein in their entirety by this reference.

**Background of the Invention**

15           Certain products and by-products of naturally-occurring metabolic processes in cells have utility in a wide array of industries, including the food, feed, cosmetics, and pharmaceutical industries. These molecules, collectively termed 'fine chemicals', include organic acids, both proteinogenic and non-proteinogenic amino acids, nucleotides and nucleosides, lipids and fatty acids, diols, carbohydrates, aromatic  
20    compounds, vitamins and cofactors, and enzymes. Their production is most conveniently performed through large-scale culture of bacteria developed to produce and secrete large quantities of a particular desired molecule. One particularly useful organism for this purpose is *Corynebacterium glutamicum*, a gram positive, nonpathogenic bacterium. Through strain selection, a number of mutant strains have  
25    been developed which produce an array of desirable compounds. However, selection of strains improved for the production of a particular molecule is a time-consuming and difficult process.

**Summary of the Invention**

30           The invention provides novel bacterial nucleic acid molecules which have a variety of uses. These uses include the identification of microorganisms which can be used to produce fine chemicals, the modulation of fine chemical production in *C.*

*glutamicum* or related bacteria, the typing or identification of *C. glutamicum* or related bacteria, as reference points for mapping the *C. glutamicum* genome, and as markers for transformation. These novel nucleic acid molecules encode proteins, referred to herein as phosphoenolpyruvate:sugar phosphotransferase system (PTS) proteins.

5            *C. glutamicum* is a gram positive, aerobic bacterium which is commonly used in industry for the large-scale production of a variety of fine chemicals, and also for the degradation of hydrocarbons (such as in petroleum spills) and for the oxidation of terpenoids. The PTS nucleic acid molecules of the invention, therefore, can be used to identify microorganisms which can be used to produce fine chemicals, *e.g.*, by  
10 fermentation processes. Modulation of the expression of the PTS nucleic acids of the invention, or modification of the sequence of the PTS nucleic acid molecules of the invention, can be used to modulate the production of one or more fine chemicals from a microorganism (*e.g.*, to improve the yield or production of one or more fine chemicals from a *Corynebacterium* or *Brevibacterium* species).

15            The PTS nucleic acids of the invention may also be used to identify an organism as being *Corynebacterium glutamicum* or a close relative thereof, or to identify the presence of *C. glutamicum* or a relative thereof in a mixed population of microorganisms. The invention provides the nucleic acid sequences of a number of *C. glutamicum* genes; by probing the extracted genomic DNA of a culture of a unique or  
20 mixed population of microorganisms under stringent conditions with a probe spanning a region of a *C. glutamicum* gene which is unique to this organism, one can ascertain whether this organism is present. Although *Corynebacterium glutamicum* itself is nonpathogenic, it is related to species pathogenic in humans, such as *Corynebacterium diphtheriae* (the causative agent of diphtheria); the detection of such organisms is of  
25 significant clinical relevance.

            The PTS nucleic acid molecules of the invention may also serve as reference points for mapping of the *C. glutamicum* genome, or of genomes of related organisms. Similarly, these molecules, or variants or portions thereof, may serve as markers for genetically engineered *Corynebacterium* or *Brevibacterium* species.

30            The PTS proteins encoded by the novel nucleic acid molecules of the invention are capable of, for example, transporting high-energy carbon-containing molecules such as glucose into *C. glutamicum*, or of participating in intracellular signal transduction in

this microorganism. Given the availability of cloning vectors for use in *Corynebacterium glutamicum*, such as those disclosed in Sinskey *et al.*, U.S. Patent No. 4,649,119, and techniques for genetic manipulation of *C. glutamicum* and the related *Brevibacterium* species (*e.g.*, *lactofermentum*) (Yoshihama *et al.*, *J. Bacteriol.* 162: 591-597 (1985); Katsumata *et al.*, *J. Bacteriol.* 159: 306-311 (1984); and Santamaria *et al.*, *J. Gen. Microbiol.* 130: 2237-2246 (1984)), the nucleic acid molecules of the invention may be utilized in the genetic engineering of this organism to make it a better or more efficient producer of one or more fine chemicals.

The PTS molecules of the invention may be modified such that the yield, production, and/or efficiency of production of one or more fine chemicals is improved. For example, by modifying a PTS protein involved in the uptake of glucose such that it is optimized in activity, the quantity of glucose uptake or the rate at which glucose is translocated into the cell may be increased. The breakdown of glucose and other sugars within the cell provides energy that may be used to drive energetically unfavorable biochemical reactions, such as those involved in the biosynthesis of fine chemicals. This breakdown also provides intermediate and precursor molecules necessary for the biosynthesis of certain fine chemicals, such as amino acids, vitamins and cofactors. By increasing the amount of intracellular high-energy carbon molecules through modification of the PTS molecules of the invention, one may therefore increase both the energy available to perform metabolic pathways necessary for the production of one or more fine chemicals, and also the intracellular pools of metabolites necessary for such production.

Further, the PTS molecules of the invention may be involved in one or more intracellular signal transduction pathways which may affect the yields and/or rate of production of one or more fine chemical from *C. glutamicum*. For example, proteins necessary for the import of one or more sugars from the extracellular medium (*e.g.*, HPr, Enzyme I, or a member of an Enzyme II complex) are frequently posttranslationally modified upon the presence of a sufficient quantity of the sugar in the cell, such that they are no longer able to import that sugar. While this quantity of sugar at which the transport system is shut off may be sufficient to sustain the normal functioning of the cell, it may be limiting for the overproduction of the desired fine chemical. Thus, it may be desirable to modify the PTS proteins of the invention such that they are no longer



responsive to such negative regulation, thereby permitting greater intracellular concentrations of one or more sugars to be achieved, and, by extension, more efficient production or greater yields of one or more fine chemicals from organisms containing such mutant PTS proteins.

- 5           This invention provides novel nucleic acid molecules which encode proteins, referred to herein as phosphoenolpyruvate:sugar phosphotransferase system (PTS) proteins, which are capable of, for example, participating in the import of high-energy carbon molecules (*e.g.*, glucose, fructose, or sucrose) into *C. glutamicum*, and/or of participating in one or more *C. glutamicum* intracellular signal transduction pathways.
- 10   Nucleic acid molecules encoding a PTS protein are referred to herein as PTS nucleic acid molecules. In a preferred embodiment, the PTS protein participates in the import of high-energy carbon molecules (*e.g.*, glucose, fructose, or sucrose) into *C. glutamicum*, and also may participate in one or more *C. glutamicum* intracellular signal transduction pathways. Examples of such proteins include those encoded by the genes set forth in
- 15   Table 1.

          Accordingly, one aspect of the invention pertains to isolated nucleic acid molecules (*e.g.*, cDNAs, DNAs, or RNAs) comprising a nucleotide sequence encoding a PTS protein or biologically active portions thereof, as well as nucleic acid fragments suitable as primers or hybridization probes for the detection or amplification of PTS-

20   encoding nucleic acid (*e.g.*, DNA or mRNA). In particularly preferred embodiments, the isolated nucleic acid molecule comprises one of the nucleotide sequences set forth in as the odd-numbered SEQ ID NOs in the Sequence Listing (*e.g.*, SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7....) or the coding region or a complement thereof of one of these nucleotide sequences. In other particularly preferred embodiments, the

25   isolated nucleic acid molecule of the invention comprises a nucleotide sequence which hybridizes to or is at least about 50%, preferably at least about 60%, more preferably at least about 70%, 80% or 90%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more homologous to a nucleotide sequence set forth, as an odd-numbered SEQ ID NO in the Sequence Listing (*e.g.*, SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5,

30   SEQ ID NO:7....), or a portion thereof. In other preferred embodiments, the isolated nucleic acid molecule encodes one of the amino acid sequences set forth in as an even-numbered SEQ ID NO in the Sequence Listing (*e.g.*, SEQ ID NO:2, SEQ ID NO:4, SEQ

ID NO:6, SEQ ID NO:8....). The preferred PTS proteins of the present invention also preferably possess at least one of the PTS activities described herein.

In another embodiment, the isolated nucleic acid molecule encodes a protein or portion thereof wherein the protein or portion thereof includes an amino acid sequence  
5 which is sufficiently homologous to an amino acid sequence of the invention (*e.g.*, a sequence having an even-numbered SEQ ID NO: in the Sequence Listing), *e.g.*, sufficiently homologous to an amino acid sequence of the invention such that the protein or portion thereof maintains a PTS activity. Preferably, the protein or portion thereof encoded by the nucleic acid molecule maintains the ability to participate in the import of  
10 high-energy carbon molecules (*e.g.*, glucose, fructose, or sucrose) into *C. glutamicum*, and/or to participate in one or more *C. glutamicum* intracellular signal transduction pathways. In one embodiment, the protein encoded by the nucleic acid molecule is at least about 50%, preferably at least about 60%, and more preferably at least about 70%, 80%, or 90% and most preferably at least about 95%, 96%, 97%, 98%, or 99% or more  
15 homologous to an amino acid sequence of the invention (*e.g.*, an entire amino acid sequence selected from those having an even-numbered SEQ ID NO in the Sequence Listing). In another preferred embodiment, the protein is a full length *C. glutamicum* protein which is substantially homologous to an entire amino acid sequence of the invention (encoded by an open reading frame shown in the corresponding odd-numbered  
20 SEQ ID NOs in the Sequence Listing (*e.g.*, SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7....)).

In another preferred embodiment, the isolated nucleic acid molecule is derived from *C. glutamicum* and encodes a protein (*e.g.*, a PTS fusion protein) which includes a biologically active domain which is at least about 50% or more homologous to one of  
25 the amino acid sequences of the invention (*e.g.*, a sequence of one of the even-numbered SEQ ID NOs in the Sequence Listing) and is able to participate in the import of high-energy carbon molecules (*e.g.*, glucose, fructose, or sucrose) into *C. glutamicum*, and/or to participate in one or more *C. glutamicum* intracellular signal transduction pathways, or possesses one or more of the activities set forth in Table 1, and which also includes  
30 heterologous nucleic acid sequences encoding a heterologous polypeptide or regulatory regions.

- 6 -

In another embodiment, the isolated nucleic acid molecule is at least 15 nucleotides in length and hybridizes under stringent conditions to a nucleic acid molecule comprising a nucleotide sequence of the invention (*e.g.*, a sequence of an odd-numbered SEQ ID NO in the Sequence Listing). Preferably, the isolated nucleic acid molecule corresponds to a naturally-occurring nucleic acid molecule. More preferably, the isolated nucleic acid encodes a naturally-occurring *C. glutamicum* PTS protein, or a biologically active portion thereof.

Another aspect of the invention pertains to vectors, *e.g.*, recombinant expression vectors, containing the nucleic acid molecules of the invention, and host cells into which such vectors have been introduced. In one embodiment, such a host cell is used to produce a PTS protein by culturing the host cell in a suitable medium. The PTS protein can be then isolated from the medium or the host cell.

Yet another aspect of the invention pertains to a genetically altered microorganism in which a PTS gene has been introduced or altered. In one embodiment, the genome of the microorganism has been altered by the introduction of a nucleic acid molecule of the invention encoding wild-type or mutated PTS sequence as a transgene. In another embodiment, an endogenous PTS gene within the genome of the microorganism has been altered, *e.g.*, functionally disrupted, by homologous recombination with an altered PTS gene. In another embodiment, an endogenous or introduced PTS gene in a microorganism has been altered by one or more point mutations, deletions, or inversions, but still encodes a functional PTS protein. In still another embodiment, one or more of the regulatory regions (*e.g.*, a promoter, repressor, or inducer) of a PTS gene in a microorganism has been altered (*e.g.*, by deletion, truncation, inversion, or point mutation) such that the expression of the PTS gene is modulated. In a preferred embodiment, the microorganism belongs to the genus *Corynebacterium* or *Brevibacterium*, with *Corynebacterium glutamicum* being particularly preferred. In a preferred embodiment, the microorganism is also utilized for the production of a desired compound, such as an amino acid, with lysine being particularly preferred.

In another aspect, the invention provides a method of identifying the presence or activity of *Cornyebacterium diphtheriae* in a subject. This method includes detection of one or more of the nucleic acid or amino acid sequences of the invention (*e.g.*, the

sequences set forth in the Sequence Listing as SEQ ID NOs 1 through 34)) in a subject, thereby detecting the presence or activity of *Corynebacterium diphtheriae* in the subject.

Still another aspect of the invention pertains to an isolated PTS protein or a portion, *e.g.*, a biologically active portion, thereof. In a preferred embodiment, the isolated PTS protein or portion thereof can participate in the import of high-energy carbon molecules (*e.g.*, glucose, fructose, or sucrose) into *C. glutamicum*, and also may participate in one or more *C. glutamicum* intracellular signal transduction pathways. In another preferred embodiment, the isolated PTS protein or portion thereof is sufficiently homologous to an amino acid sequence of the invention (*e.g.*, a sequence of an even-numbered SEQ ID NO: in the Sequence Listing) such that the protein or portion thereof maintains the ability to participate in the import of high-energy carbon molecules (*e.g.*, glucose, fructose, or sucrose) into *C. glutamicum*, and /or to participate in one or more *C. glutamicum* intracellular signal transduction pathways.

The invention also provides an isolated preparation of a PTS protein. In preferred embodiments, the PTS protein comprises an amino acid sequence of the invention (*e.g.*, a sequence of an even-numbered SEQ ID NO: of the Sequence Listing). In another preferred embodiment, the invention pertains to an isolated full length protein which is substantially homologous to an entire amino acid sequence of the invention (*e.g.*, a sequence of an even-numbered SEQ ID NO: of the Sequence Listing) (encoded by an open reading frame set in a corresponding odd-numbered SEQ ID NO: of the Sequence Listing). In yet another embodiment, the protein is at least about 50%, preferably at least about 60%, and more preferably at least about 70%, 80%, or 90%, and most preferably at least about 95%, 96%, 97%, 98%, or 99% or more homologous to an entire amino acid sequence of the invention (*e.g.*, a sequence of an even-numbered SEQ ID NO: of the Sequence Listing). In other embodiments, the isolated PTS protein comprises an amino acid sequence which is at least about 50% or more homologous to one of the amino acid sequences of the invention (*e.g.*, a sequence of an even-numbered SEQ ID NO: of the Sequence Listing) and is able to participate in the import of high-energy carbon molecules (*e.g.*, glucose, fructose, or sucrose) into *C. glutamicum*, and/or to participate in one or more *C. glutamicum* intracellular signal transduction pathways, or has one or more of the activities set forth in Table 1.

Alternatively, the isolated PTS protein can comprise an amino acid sequence which is encoded by a nucleotide sequence which hybridizes, *e.g.*, hybridizes under stringent conditions, or is at least about 50%, preferably at least about 60%, more preferably at least about 70%, 80%, or 90%, and even more preferably at least about 5 95%, 96%, 97%, 98,%, or 99% or more homologous, to a nucleotide sequence of one of the even-numbered SEQ ID NOs set forth in the Sequence Listing. It is also preferred that the preferred forms of PTS proteins also have one or more of the PTS bioactivities described herein.

The PTS polypeptide, or a biologically active portion thereof, can be operatively 10 linked to a non-PTS polypeptide to form a fusion protein. In preferred embodiments, this fusion protein has an activity which differs from that of the PTS protein alone. In other preferred embodiments, this fusion protein results in increased yields, production, and/or efficiency of production of a desired fine chemical from *C. glutamicum*. In particularly preferred embodiments, integration of this fusion protein into a host cell 15 modulates the production of a desired compound from the cell.

In another aspect, the invention provides methods for screening molecules which modulate the activity of a PTS protein, either by interacting with the protein itself or a substrate or binding partner of the PTS protein, or by modulating the transcription or translation of a PTS nucleic acid molecule of the invention.

20 Another aspect of the invention pertains to a method for producing a fine chemical. This method involves the culturing of a cell containing a vector directing the expression of a PTS nucleic acid molecule of the invention, such that a fine chemical is produced. In a preferred embodiment, this method further includes the step of obtaining a cell containing such a vector, in which a cell is transfected with a vector directing the 25 expression of a PTS nucleic acid. In another preferred embodiment, this method further includes the step of recovering the fine chemical from the culture. In a particularly preferred embodiment, the cell is from the genus *Corynebacterium* or *Brevibacterium*, or is selected from those strains set forth in Table 3.

Another aspect of the invention pertains to methods for modulating production of 30 a molecule from a microorganism. Such methods include contacting the cell with an agent which modulates PTS protein activity or PTS nucleic acid expression such that a cell associated activity is altered relative to this same activity in the absence of the

agent. In a preferred embodiment, the cell is modulated for the uptake of one or more sugars, such that the yields or rate of production of a desired fine chemical by this microorganism is improved. The agent which modulates PTS protein activity can be an agent which stimulates PTS protein activity or PTS nucleic acid expression. Examples  
5 of agents which stimulate PTS protein activity or PTS nucleic acid expression include small molecules, active PTS proteins, and nucleic acids encoding PTS proteins that have been introduced into the cell. Examples of agents which inhibit PTS activity or expression include small molecules, and antisense PTS nucleic acid molecules.

Another aspect of the invention pertains to methods for modulating yields of a  
10 desired compound from a cell, involving the introduction of a wild-type or mutant PTS gene into a cell, either maintained on a separate plasmid or integrated into the genome of the host cell. If integrated into the genome, such integration can random, or it can take place by homologous recombination such that the native gene is replaced by the  
15 introduced copy, causing the production of the desired compound from the cell to be modulated. In a preferred embodiment, said yields are increased. In another preferred embodiment, said chemical is a fine chemical. In a particularly preferred embodiment, said fine chemical is an amino acid. In especially preferred embodiments, said amino acid is L-lysine.

## 20 Detailed Description of the Invention

The present invention provides PTS nucleic acid and protein molecules which are involved in the uptake of high-energy carbon molecules (*e.g.*, sucrose, fructose, or glucose) into *C. glutamicum*, and may also participate in intracellular signal transduction pathways in this microorganism. The molecules of the invention may be utilized in the  
25 modulation of production of fine chemicals from microorganisms. Such modulation may be due to increased intracellular levels of high-energy molecules needed to produce, *e.g.*, ATP, GTP and other molecules utilized to drive energetically unfavorable biochemical reactions in the cell, such as the biosynthesis of a fine chemical. This modulation of fine chemical production may also be due to the fact that the breakdown  
30 products of many sugars serve as intermediates or precursors for other biosynthetic pathways, including those of certain fine chemicals. Further, PTS proteins are known to participate in certain intracellular signal transduction pathways which may have

regulatory activity for one or more fine chemical metabolic pathways; by manipulating these PTS proteins, one may thereby activate a fine chemical biosynthetic pathways or repress a fine chemical degradation pathway. Aspects of the invention are further explicated below.

5

#### I. Fine Chemicals

The term 'fine chemical' is art-recognized and includes molecules produced by an organism which have applications in various industries, such as, but not limited to, the pharmaceutical, agriculture, and cosmetics industries. Such compounds include  
10 organic acids, such as tartaric acid, itaconic acid, and diaminopimelic acid, both proteinogenic and non-proteinogenic amino acids, purine and pyrimidine bases, nucleosides, and nucleotides (as described *e.g.* in Kuninaka, A. (1996) Nucleotides and related compounds, p. 561-612, in Biotechnology vol. 6, Rehm *et al.*, eds. VCH: Weinheim, and references contained therein), lipids, both saturated and unsaturated fatty  
15 acids (*e.g.*, arachidonic acid), diols (*e.g.*, propane diol, and butane diol), carbohydrates (*e.g.*, hyaluronic acid and trehalose), aromatic compounds (*e.g.*, aromatic amines, vanillin, and indigo), vitamins and cofactors (as described in Ullmann's Encyclopedia of Industrial Chemistry, vol. A27, "Vitamins", p. 443-613 (1996) VCH: Weinheim and references therein; and Ong, A.S., Niki, E. & Packer, L. (1995) "Nutrition, Lipids,  
20 Health, and Disease" Proceedings of the UNESCO/Confederation of Scientific and Technological Associations in Malaysia, and the Society for Free Radical Research – Asia, held Sept. 1-3, 1994 at Penang, Malaysia, AOCS Press, (1995)), enzymes, polyketides (Cane *et al.* (1998) *Science* 282: 63-68), and all other chemicals described in Gutcho (1983) Chemicals by Fermentation, Noyes Data Corporation, ISBN:  
25 0818805086 and references therein. The metabolism and uses of certain of these fine chemicals are further explicated below.

#### *A. Amino Acid Metabolism and Uses*

Amino acids comprise the basic structural units of all proteins, and as such are  
30 essential for normal cellular functioning in all organisms. The term "amino acid" is art-recognized. The proteinogenic amino acids, of which there are 20 species, serve as structural units for proteins, in which they are linked by peptide bonds, while the

nonproteinogenic amino acids (hundreds of which are known) are not normally found in proteins (see Ulmann's Encyclopedia of Industrial Chemistry, vol. A2, p. 57-97 VCH: Weinheim (1985)). Amino acids may be in the D- or L- optical configuration, though L-amino acids are generally the only type found in naturally-occurring proteins.

- 5 Biosynthetic and degradative pathways of each of the 20 proteinogenic amino acids have been well characterized in both prokaryotic and eukaryotic cells (see, for example, Stryer, L. Biochemistry, 3<sup>rd</sup> edition, pages 578-590 (1988)). The 'essential' amino acids (histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, and valine), so named because they are generally a nutritional requirement due to the
- 10 complexity of their biosyntheses, are readily converted by simple biosynthetic pathways to the remaining 11 'nonessential' amino acids (alanine, arginine, asparagine, aspartate, cysteine, glutamate, glutamine, glycine, proline, serine, and tyrosine). Higher animals do retain the ability to synthesize some of these amino acids, but the essential amino acids must be supplied from the diet in order for normal protein synthesis to occur.
- 15 Aside from their function in protein biosynthesis, these amino acids are interesting chemicals in their own right, and many have been found to have various applications in the food, feed, chemical, cosmetics, agriculture, and pharmaceutical industries. Lysine is an important amino acid in the nutrition not only of humans, but also of monogastric animals such as poultry and swine. Glutamate is most commonly
- 20 used as a flavor additive (mono-sodium glutamate, MSG) and is widely used throughout the food industry, as are aspartate, phenylalanine, glycine, and cysteine. Glycine, L-methionine and tryptophan are all utilized in the pharmaceutical industry. Glutamine, valine, leucine, isoleucine, histidine, arginine, proline, serine and alanine are of use in both the pharmaceutical and cosmetics industries. Threonine, tryptophan, and D/ L-
- 25 methionine are common feed additives. (Leuchtenberger, W. (1996) Amino acids – technical production and use, p. 466-502 in Rehm *et al.* (eds.) Biotechnology vol. 6, chapter 14a, VCH: Weinheim). Additionally, these amino acids have been found to be useful as precursors for the synthesis of synthetic amino acids and proteins, such as N-acetylcysteine, S-carboxymethyl-L-cysteine, (S)-5-hydroxytryptophan, and others
- 30 described in Ulmann's Encyclopedia of Industrial Chemistry, vol. A2, p. 57-97, VCH: Weinheim, 1985.



The biosynthesis of these natural amino acids in organisms capable of producing them, such as bacteria, has been well characterized (for review of bacterial amino acid biosynthesis and regulation thereof, see Umbarger, H.E.(1978) *Ann. Rev. Biochem.* 47: 533-606). Glutamate is synthesized by the reductive amination of  $\alpha$ -ketoglutarate, an intermediate in the citric acid cycle. Glutamine, proline, and arginine are each subsequently produced from glutamate. The biosynthesis of serine is a three-step process beginning with 3-phosphoglycerate (an intermediate in glycolysis), and resulting in this amino acid after oxidation, transamination, and hydrolysis steps. Both cysteine and glycine are produced from serine; the former by the condensation of homocysteine with serine, and the latter by the transferal of the side-chain  $\beta$ -carbon atom to tetrahydrofolate, in a reaction catalyzed by serine transhydroxymethylase. Phenylalanine, and tyrosine are synthesized from the glycolytic and pentose phosphate pathway precursors erythrose 4-phosphate and phosphoenolpyruvate in a 9-step biosynthetic pathway that differ only at the final two steps after synthesis of prephenate. Tryptophan is also produced from these two initial molecules, but its synthesis is an 11-step pathway. Tyrosine may also be synthesized from phenylalanine, in a reaction catalyzed by phenylalanine hydroxylase. Alanine, valine, and leucine are all biosynthetic products of pyruvate, the final product of glycolysis. Aspartate is formed from oxaloacetate, an intermediate of the citric acid cycle. Asparagine, methionine, threonine, and lysine are each produced by the conversion of aspartate. Isoleucine is formed from threonine. A complex 9-step pathway results in the production of histidine from 5-phosphoribosyl-1-pyrophosphate, an activated sugar.

Amino acids in excess of the protein synthesis needs of the cell cannot be stored, and are instead degraded to provide intermediates for the major metabolic pathways of the cell (for review see Stryer, L. *Biochemistry* 3<sup>rd</sup> ed. Ch. 21 "Amino Acid Degradation and the Urea Cycle" p. 495-516 (1988)). Although the cell is able to convert unwanted amino acids into useful metabolic intermediates, amino acid production is costly in terms of energy, precursor molecules, and the enzymes necessary to synthesize them. Thus it is not surprising that amino acid biosynthesis is regulated by feedback inhibition, in which the presence of a particular amino acid serves to slow or entirely stop its own production (for overview of feedback mechanisms in amino acid biosynthetic pathways, see Stryer, L. *Biochemistry*, 3<sup>rd</sup> ed. Ch. 24: "Biosynthesis of Amino Acids and Heme" p.

575-600 (1988)). Thus, the output of any particular amino acid is limited by the amount of that amino acid present in the cell.

*B. Vitamin, Cofactor, and Nutraceutical Metabolism and Uses*

5 Vitamins, cofactors, and nutraceuticals comprise another group of molecules which the higher animals have lost the ability to synthesize and so must ingest, although they are readily synthesized by other organisms, such as bacteria. These molecules are either bioactive substances themselves, or are precursors of biologically active substances which may serve as electron carriers or intermediates in a variety of  
10 metabolic pathways. Aside from their nutritive value, these compounds also have significant industrial value as coloring agents, antioxidants, and catalysts or other processing aids. (For an overview of the structure, activity, and industrial applications of these compounds, see, for example, Ullman's Encyclopedia of Industrial Chemistry, "Vitamins" vol. A27, p. 443-613, VCH: Weinheim, 1996.) The term "vitamin" is art-  
15 recognized, and includes nutrients which are required by an organism for normal functioning, but which that organism cannot synthesize by itself. The group of vitamins may encompass cofactors and nutraceutical compounds. The language "cofactor" includes nonproteinaceous compounds required for a normal enzymatic activity to occur. Such compounds may be organic or inorganic; the cofactor molecules of the  
20 invention are preferably organic. The term "nutraceutical" includes dietary supplements having health benefits in plants and animals, particularly humans. Examples of such molecules are vitamins, antioxidants, and also certain lipids (*e.g.*, polyunsaturated fatty acids).

The biosynthesis of these molecules in organisms capable of producing them,  
25 such as bacteria, has been largely characterized (Ullman's Encyclopedia of Industrial Chemistry, "Vitamins" vol. A27, p. 443-613, VCH: Weinheim, 1996; Michal, G. (1999) Biochemical Pathways: An Atlas of Biochemistry and Molecular Biology, John Wiley & Sons; Ong, A.S., Niki, E. & Packer, L. (1995) "Nutrition, Lipids, Health, and Disease" Proceedings of the UNESCO/Confederation of Scientific and Technological  
30 Associations in Malaysia, and the Society for Free Radical Research – Asia, held Sept. 1-3, 1994 at Penang, Malaysia, AOCS Press: Champaign, IL X, 374 S).

Thiamin (vitamin B<sub>1</sub>) is produced by the chemical coupling of pyrimidine and thiazole moieties. Riboflavin (vitamin B<sub>2</sub>) is synthesized from guanosine-5'-triphosphate (GTP) and ribose-5'-phosphate. Riboflavin, in turn, is utilized for the synthesis of flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD). The family of  
5 compounds collectively termed 'vitamin B<sub>6</sub>' (e.g., pyridoxine, pyridoxamine, pyridoxal-5'-phosphate, and the commercially used pyridoxin hydrochloride) are all derivatives of the common structural unit, 5-hydroxy-6-methylpyridine. Pantothenate (pantothenic acid, (R)-(+)-N-(2,4-dihydroxy-3,3-dimethyl-1-oxobutyl)- $\beta$ -alanine) can be produced either by chemical synthesis or by fermentation. The final steps in pantothenate  
10 biosynthesis consist of the ATP-driven condensation of  $\beta$ -alanine and pantoic acid. The enzymes responsible for the biosynthesis steps for the conversion to pantoic acid, to  $\beta$ -alanine and for the condensation to pantothenic acid are known. The metabolically active form of pantothenate is Coenzyme A, for which the biosynthesis proceeds in 5 enzymatic steps. Pantothenate, pyridoxal-5'-phosphate, cysteine and ATP are the  
15 precursors of Coenzyme A. These enzymes not only catalyze the formation of pantothenate, but also the production of (R)-pantoic acid, (R)-pantolacton, (R)-panthenol (provitamin B<sub>5</sub>), pantetheine (and its derivatives) and coenzyme A.

Biotin biosynthesis from the precursor molecule pimeloyl-CoA in microorganisms has been studied in detail and several of the genes involved have been  
20 identified. Many of the corresponding proteins have been found to also be involved in Fe-cluster synthesis and are members of the nifS class of proteins. Lipic acid is derived from octanoic acid, and serves as a coenzyme in energy metabolism, where it becomes part of the pyruvate dehydrogenase complex and the  $\alpha$ -ketoglutarate dehydrogenase complex. The folates are a group of substances which are all derivatives  
25 of folic acid, which in turn is derived from L-glutamic acid, p-amino-benzoic acid and 6-methylpterin. The biosynthesis of folic acid and its derivatives, starting from the metabolism intermediates guanosine-5'-triphosphate (GTP), L-glutamic acid and p-amino-benzoic acid has been studied in detail in certain microorganisms.

Corrinoids (such as the cobalamines and particularly vitamin B<sub>12</sub>) and  
30 porphyrines belong to a group of chemicals characterized by a tetrapyrrole ring system. The biosynthesis of vitamin B<sub>12</sub> is sufficiently complex that it has not yet been completely characterized, but many of the enzymes and substrates involved are now

known. Nicotinic acid (nicotinate), and nicotinamide are pyridine derivatives which are also termed 'niacin'. Niacin is the precursor of the important coenzymes NAD (nicotinamide adenine dinucleotide) and NADP (nicotinamide adenine dinucleotide phosphate) and their reduced forms.

5       The large-scale production of these compounds has largely relied on cell-free chemical syntheses, though some of these chemicals have also been produced by large-scale culture of microorganisms, such as riboflavin, Vitamin B<sub>6</sub>, pantothenate, and biotin. Only Vitamin B<sub>12</sub> is produced solely by fermentation, due to the complexity of its synthesis. *In vitro* methodologies require significant inputs of materials and time,  
10       often at great cost.

### *C. Purine, Pyrimidine, Nucleoside and Nucleotide Metabolism and Uses*

Purine and pyrimidine metabolism genes and their corresponding proteins are important targets for the therapy of tumor diseases and viral infections. The language  
15       "purine" or "pyrimidine" includes the nitrogenous bases which are constituents of nucleic acids, co-enzymes, and nucleotides. The term "nucleotide" includes the basic structural units of nucleic acid molecules, which are comprised of a nitrogenous base, a pentose sugar (in the case of RNA, the sugar is ribose; in the case of DNA, the sugar is D-deoxyribose), and phosphoric acid. The language "nucleoside" includes molecules  
20       which serve as precursors to nucleotides, but which are lacking the phosphoric acid moiety that nucleotides possess. By inhibiting the biosynthesis of these molecules, or their mobilization to form nucleic acid molecules, it is possible to inhibit RNA and DNA synthesis; by inhibiting this activity in a fashion targeted to cancerous cells, the ability of tumor cells to divide and replicate may be inhibited. Additionally, there are  
25       nucleotides which do not form nucleic acid molecules, but rather serve as energy stores (i.e., AMP) or as coenzymes (i.e., FAD and NAD).

Several publications have described the use of these chemicals for these medical indications, by influencing purine and/or pyrimidine metabolism (*e.g.* Christopherson, R.I. and Lyons, S.D. (1990) "Potent inhibitors of *de novo* pyrimidine and purine  
30       biosynthesis as chemotherapeutic agents." *Med. Res. Reviews* 10: 505-548). Studies of enzymes involved in purine and pyrimidine metabolism have been focused on the development of new drugs which can be used, for example, as immunosuppressants or

anti-proliferants (Smith, J.L., (1995) "Enzymes in nucleotide synthesis." *Curr. Opin. Struct. Biol.* 5: 752-757; (1995) *Biochem Soc. Transact.* 23: 877-902). However, purine and pyrimidine bases, nucleosides and nucleotides have other utilities: as intermediates in the biosynthesis of several fine chemicals (e.g., thiamine, S-adenosyl-methionine, folates, or riboflavin), as energy carriers for the cell (e.g., ATP or GTP), and for chemicals themselves, commonly used as flavor enhancers (e.g., IMP or GMP) or for several medicinal applications (see, for example, Kuninaka, A. (1996) *Nucleotides and Related Compounds in Biotechnology* vol. 6, Rehm *et al.*, eds. VCH: Weinheim, p. 561-612). Also, enzymes involved in purine, pyrimidine, nucleoside, or nucleotide metabolism are increasingly serving as targets against which chemicals for crop protection, including fungicides, herbicides and insecticides, are developed.

The metabolism of these compounds in bacteria has been characterized (for reviews see, for example, Zalkin, H. and Dixon, J.E. (1992) "*de novo* purine nucleotide biosynthesis", in: *Progress in Nucleic Acid Research and Molecular Biology*, vol. 42, Academic Press:, p. 259-287; and Michal, G. (1999) "Nucleotides and Nucleosides", Chapter 8 in: *Biochemical Pathways: An Atlas of Biochemistry and Molecular Biology*, Wiley: New York). Purine metabolism has been the subject of intensive research, and is essential to the normal functioning of the cell. Impaired purine metabolism in higher animals can cause severe disease, such as gout. Purine nucleotides are synthesized from ribose-5-phosphate, in a series of steps through the intermediate compound inosine-5'-phosphate (IMP), resulting in the production of guanosine-5'-monophosphate (GMP) or adenosine-5'-monophosphate (AMP), from which the triphosphate forms utilized as nucleotides are readily formed. These compounds are also utilized as energy stores, so their degradation provides energy for many different biochemical processes in the cell. Pyrimidine biosynthesis proceeds by the formation of uridine-5'-monophosphate (UMP) from ribose-5-phosphate. UMP, in turn, is converted to cytidine-5'-triphosphate (CTP). The deoxy- forms of all of these nucleotides are produced in a one step reduction reaction from the diphosphate ribose form of the nucleotide to the diphosphate deoxyribose form of the nucleotide. Upon phosphorylation, these molecules are able to participate in DNA synthesis.

#### D. Trehalose Metabolism and Uses

Trehalose consists of two glucose molecules, bound in  $\alpha$ ,  $\alpha$ -1,1 linkage. It is commonly used in the food industry as a sweetener, an additive for dried or frozen foods, and in beverages. However, it also has applications in the pharmaceutical, cosmetics and biotechnology industries (see, for example, Nishimoto *et al.*, (1998) U.S. Patent No. 5,759,610; Singer, M.A. and Lindquist, S. (1998) *Trends Biotech.* 16: 460-467; Paiva, C.L.A. and Panek, A.D. (1996) *Biotech. Ann. Rev.* 2: 293-314; and Shiosaka, M. (1997) *J. Japan* 172: 97-102). Trehalose is produced by enzymes from many microorganisms and is naturally released into the surrounding medium, from which it can be collected using methods known in the art.

#### II. The Phosphoenolpyruvate: Sugar Phosphotransferase System

The ability of cells to grow and divide rapidly in culture is to a great degree dependent on the extent to which the cells are able to take up and utilize high energy molecules, such as glucose and other sugars. Different transporter proteins exist to transport different carbon sources into the cell. There are transport proteins for sugars, such as glucose, fructose, mannose, galactose, ribose, sorbose, ribulose, lactose, maltose, sucrose, or raffinose, and also transport proteins for starch or cellulose degradation products. Other transport systems serve to import alcohols (*e.g.*, methanol or ethanol), alkanes, fatty acids and organic acids like acetic acid or lactic acid. In bacteria, sugars may be transported into the cell across the cellular membrane by a variety of mechanisms. Aside from the symport of sugars with protons, one of the most commonly utilized processes for sugar uptake is the bacterial phosphoenolpyruvate: sugar phosphotransferase system (PTS). This system not only catalyzes the translocation (with concomitant phosphorylation) of sugars and hexitols, but it also regulates cellular metabolism in response to the availability of carbohydrates. Such PTS systems are ubiquitous in bacteria but do not occur in archaeobacteria or eukaryotes.

Functionally, the PTS system consists of two cytoplasmic proteins, Enzyme I and HPr, and a variable number of sugar-specific integral and peripheral membrane transport complexes (each termed 'Enzyme II' with a sugar-specific subscript, *e.g.*, 'Enzyme II<sup>Glu</sup>' for the Enzyme II complex which binds glucose). Enzymes II specific for mono-, di-, or oligosaccharides, like glucose, fructose, mannose, galactose, ribose,

- 18 -

sorbose, ribulose, lactose, maltose, sucrose, raffinose, and others are known. Enzyme I transfers phosphoryl groups from phosphoenolpyruvate (PEP) to the phosphoryl carrier protein, HPr. HPr then transfers the phosphoryl groups to the different Enzyme II transport complexes. While the amino acid sequences of Enzyme I and HPr are quite similar in all bacteria, the sequences for PTS transporters can be grouped into structurally unrelated families. Further, the number and homology between these genes vary from bacteria to bacteria. The *E. coli* genome encodes 38 different PTS proteins, 33 of which are subunits belonging to 22 different transporters. The *M. genitalium* genome contains one gene each for Enzyme I and HPr, and only two genes for PTS transporters. The genomes of *T. palladium* and *C. trachomatis* contain genes for Enzyme I- and HPr-like proteins but no PTS transporters.

All PTS transporters consist of three functional units, IIA, IIB, and IIC, which occur either as protein subunits in a complex (*e.g.*, IIA<sup>Glc</sup>IICB<sup>Glc</sup>) or as domains of a single polypeptide chain (*e.g.*, IICBA<sup>GlcNAc</sup>). IIA and IIB sequentially transfer phosphoryl groups from HPr to the transported sugars. IIC contains the sugar binding site, and spans the inner membrane six or eight times. Sugar translocation is coupled to the transient phosphorylation of the IIB domain. Enzyme I, HPr, and IIA are phosphorylated at histidine residues, while IIB subunits are phosphorylated at either cysteine or histidine residues, depending on the particular transporter involved. Phosphorylation of the sugar being imported has the advantage of blocking the diffusion of the sugar back through the cellular membrane to the extracellular medium, since the charged phosphate group cannot readily traverse the hydrophobic core of the membrane.

Some PTS proteins play a role in intracellular signal transduction in addition to their function in the active transport of sugars. These subunits regulate their targets either allosterically, or by phosphorylation. Their regulatory activity varies with the degree of their phosphorylation (*i.e.*, the ratio of the non-phosphorylated to the phosphorylated form), which in turn varies with the ratio of sugar-dependent dephosphorylation and phosphoenolpyruvate-dependent rephosphorylation. Examples of such intracellular regulation by PTS proteins in *E. coli* include the inhibition of glycerol kinase by dephosphorylated IIA<sup>Glc</sup>, and the activation of adenylate cyclase by the phosphorylated version of this protein. Also, the HPr and the IIB domains of some transporters in these microorganisms regulate gene expression by reversible

phosphorylation of transcription antiterminators. In gram-positive bacteria, the activity of HPr is modulated by HPr-specific serine kinases and phosphatases. For example, HPr phosphorylated at serine-46 functions as a co-repressor of the transcriptional repressor CcpA. Lastly, it has been found that unphosphorylated Enzyme I inhibits the sensor kinase CheA of the bacterial chemotaxis machinery, providing a direct link between the sugar binding and transport systems of the bacterium and those systems governing movement of the bacterium (Sonenshein, A. L., *et al.*, eds. *Bacillus subtilis* and other gram-positive bacteria. ASM: Washington, D.C.; Neidhardt, F.C., *et al.*, eds. (1996) *Escherichia coli* and *Salmonella*. ASM Press: Washington, D.C.; Lengeler *et al.*, (1999). Biology of Prokaryotes. Section II, pp. 68-87, Thieme Verlag: Stuttgart).

### III. Elements and Methods of the Invention

The present invention is based, at least in part, on the discovery of novel molecules, referred to herein as PTS nucleic acid and protein molecules, which participate in the uptake of high-energy carbon molecules (*e.g.*, glucose, sucrose, and fructose) into *C. glutamicum*, and may also participate in one or more intracellular signal transduction pathways in these microorganisms. In one embodiment, the PTS molecules function to import high-energy carbon molecules into the cell, where the energy produced by their degradation may be utilized to power less energetically favorable biochemical reactions, and their degradation products may serve as intermediates and precursors for a number of other metabolic pathways. In another embodiment, the PTS molecules may participate in one or more intracellular signal transduction pathways, wherein the presence of a modified form of a PTS molecule (*e.g.*, a phosphorylated PTS protein) may participate in a signal transduction cascade which regulates one or more cellular processes. In a preferred embodiment, the activity of the PTS molecules of the present invention has an impact on the production of a desired fine chemical by this organism. In a particularly preferred embodiment, the PTS molecules of the invention are modulated in activity, such that the yield, production or efficiency of production of one or more fine chemicals from *C. glutamicum* is also modulated.

The language, "PTS protein" or "PTS polypeptide" includes proteins which participate in the uptake of one or more high-energy carbon compounds (*e.g.*, mono-, di, or oligosaccharides, such as fructose, mannose, sucrose, glucose, raffinose, galactose,



ribose, lactose, maltose, and ribulose) from the extracellular medium to the interior of the cell. Such PTS proteins may also participate in one or more intracellular signal transduction pathways, such as, but not limited to, those governing the uptake of different sugars into the cell. Examples of PTS proteins include those encoded by the

5 PTS genes set forth in Table 1 and by the odd-numbered SEQ ID NOs. For general references pertaining to the PTS system, see: Stryer, L. (1988) *Biochemistry*. Chapter 37: "Membrane Transport", W.H. Freeman: New York, p. 959-961; Darnell, J. *et al.* (1990) *Molecular Cell Biology* Scientific American Books: New York, p. 552-553, and Michal, G., ed. (1999) *Biochemical Pathways: An Atlas of Biochemistry and Molecular*

10 *Biology*, Chapter 15 "Special Bacterial Metabolism". The terms "PTS gene" or "PTS nucleic acid sequence" include nucleic acid sequences encoding a PTS protein, which consist of a coding region and also corresponding untranslated 5' and 3' sequence regions. Examples of PTS genes include those set forth in Table 1. The terms "production" or "productivity" are art-recognized and include the concentration of the

15 fermentation product (for example, the desired fine chemical) formed within a given time and a given fermentation volume (*e.g.*, kg product per hour per liter). The term "efficiency of production" includes the time required for a particular level of production to be achieved (for example, how long it takes for the cell to attain a particular rate of output of a fine chemical). The term "yield" or "product/carbon yield" is art-recognized

20 and includes the efficiency of the conversion of the carbon source into the product (*i.e.*, fine chemical). This is generally written as, for example, kg product per kg carbon source. By increasing the yield or production of the compound, the quantity of recovered molecules, or of useful recovered molecules of that compound in a given amount of culture over a given amount of time is increased. The terms "biosynthesis" or

25 a "biosynthetic pathway" are art-recognized and include the synthesis of a compound, preferably an organic compound, by a cell from intermediate compounds in what may be a multistep and highly regulated process. The terms "degradation" or a "degradation pathway" are art-recognized and include the breakdown of a compound, preferably an organic compound, by a cell to degradation products (generally speaking, smaller or less

30 complex molecules) in what may be a multistep and highly regulated process. The language "metabolism" is art-recognized and includes the totality of the biochemical reactions that take place in an organism. The metabolism of a particular compound,

then, (e.g., the metabolism of an amino acid such as glycine) comprises the overall biosynthetic, modification, and degradation pathways in the cell related to this compound. The language "transport" or "import" is art-recognized and includes the facilitated movement of one or more molecules across a cellular membrane through  
5 which the molecule would otherwise be unable to pass.

In another embodiment, the PTS molecules of the invention are capable of modulating the production of a desired molecule, such as a fine chemical, in a microorganism such as *C. glutamicum*. Using recombinant genetic techniques, one or more of the PTS proteins of the invention may be manipulated such that its function is  
10 modulated. For example, a protein involved in the PTS-mediated import of glucose may be altered such that it is optimized in activity, and the PTS system for the importation of glucose may thus be able to translocate increased amounts of glucose into the cell. Since glucose molecules are utilized not only for energy to drive energetically unfavorable biochemical reactions, such as fine chemical biosyntheses, but also as  
15 precursors and intermediates in a number of fine chemical biosynthetic pathways (e.g., serine is synthesized from 3-phosphoglycerate). In each case, the overall yield or rate of production of one of these desired fine chemicals may be increased, either by increasing the energy available for such production to occur, or by increasing the availability of compounds necessary for such production to take place.

20 Further, many PTS proteins are known to play key roles in intracellular signal transduction pathways which regulate cellular metabolism and sugar uptake in keeping with the availability of carbon sources. For example, it is known that an increased intracellular level of fructose 1,6-bisphosphate (a compound produced during glycolysis) results in the phosphorylation of a serine residue on HPr which prevents this  
25 protein from serving as a phosphoryl donor in any PTS sugar transport process, thereby blocking further sugar uptake. By mutagenizing HPr such that this serine residue cannot be phosphorylated, one may constitutively activate HPr and thereby increase sugar transport into the cell, which in turn will ensure greater intracellular energy stores and intermediate/precursor molecules for the biosynthesis of one or more desired fine  
30 chemicals.

The isolated nucleic acid sequences of the invention are contained within the genome of a *Corynebacterium glutamicum* strain available through the American Type

Culture Collection, given designation ATCC 13032. The nucleotide sequence of the isolated *C. glutamicum* PTS DNAs and the predicted amino acid sequences of the *C. glutamicum* PTS proteins are shown in the Sequence Listing as odd-numbered SEQ ID NOs and even-numbered SEQ ID NOs, respectively.

- 5           Computational analyses were performed which classified and/or identified these nucleotide sequences as sequences which encode metabolic pathway proteins.

          The present invention also pertains to proteins which have an amino acid sequence which is substantially homologous to an amino acid sequence of the invention (e.g., the sequence of an even-numbered SEQ ID NO of the Sequence Listing). As used  
10   herein, a protein which has an amino acid sequence which is substantially homologous to a selected amino acid sequence is least about 50% homologous to the selected amino acid sequence, e.g., the entire selected amino acid sequence. A protein which has an amino acid sequence which is substantially homologous to a selected amino acid sequence can also be least about 50-60%, preferably at least about 60-70%, and more  
15   preferably at least about 70-80%, 80-90%, or 90-95%, and most preferably at least about 96%, 97%, 98%, 99% or more homologous to the selected amino acid sequence.

          The PTS protein or a biologically active portion or fragment thereof of the invention can participate in the transport of high-energy carbon-containing molecules such as glucose into *C. glutamicum*, or can participate in intracellular signal transduction  
20   in this microorganism, or may have one or more of the activities set forth in Table 1.

          Various aspects of the invention are described in further detail in the following subsections:

#### *A. Isolated Nucleic Acid Molecules*

- 25           One aspect of the invention pertains to isolated nucleic acid molecules that encode PTS polypeptides or biologically active portions thereof, as well as nucleic acid fragments sufficient for use as hybridization probes or primers for the identification or amplification of PTS-encoding nucleic acid (e.g., PTS DNA). As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (e.g., cDNA or genomic  
30   DNA) and RNA molecules (e.g., mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. This term also encompasses untranslated sequence located at both the 3' and 5' ends of the coding region of the gene: at least about 100 nucleotides

of sequence upstream from the 5' end of the coding region and at least about 20 nucleotides of sequence downstream from the 3' end of the coding region of the gene. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA. An "isolated" nucleic acid molecule is one which is separated  
5 from other nucleic acid molecules which are present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated PTS nucleic acid molecule can contain less than  
10 about 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived (e.g., a *C. glutamicum* cell). Moreover, an "isolated" nucleic acid molecule, such as a DNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or  
15 other chemicals when chemically synthesized.

A nucleic acid molecule of the present invention, e.g., a nucleic acid molecule having a nucleotide sequence of an odd-numbered SEQ ID NO of the Sequence Listing, or a portion thereof, can be isolated using standard molecular biology techniques and the sequence information provided herein. For example, a *C. glutamicum* PTS DNA can be  
20 isolated from a *C. glutamicum* library using all or portion of one of the odd-numbered SEQ ID NO sequences of the Sequence Listing as a hybridization probe and standard hybridization techniques (e.g., as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).  
25 Moreover, a nucleic acid molecule encompassing all or a portion of one of the nucleic acid sequences of the invention (e.g., an odd-numbered SEQ ID NO of the Sequence Listing) can be isolated by the polymerase chain reaction using oligonucleotide primers designed based upon this sequence (e.g., a nucleic acid molecule encompassing all or a portion of one of the nucleic acid sequences of the invention (e.g., an odd-numbered  
30 SEQ ID NO of the Sequence Listing) can be isolated by the polymerase chain reaction using oligonucleotide primers designed based upon this same sequence). For example, mRNA can be isolated from normal endothelial cells (e.g., by the guanidinium-

- 24 -

thiocyanate extraction procedure of Chirgwin *et al.* (1979) *Biochemistry* 18: 5294-5299) and DNA can be prepared using reverse transcriptase (*e.g.*, Moloney MLV reverse transcriptase, available from Gibco/BRL, Bethesda, MD; or AMV reverse transcriptase, available from Seikagaku America, Inc., St. Petersburg, FL). Synthetic oligonucleotide  
5 primers for polymerase chain reaction amplification can be designed based upon one of the nucleotide sequences shown in the Sequence Listing. A nucleic acid of the invention can be amplified using cDNA or, alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and  
10 characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to a PTS nucleotide sequence can be prepared by standard synthetic techniques, *e.g.*, using an automated DNA synthesizer.

In a preferred embodiment, an isolated nucleic acid molecule of the invention comprises one of the nucleotide sequences shown in the Sequence Listing. The nucleic  
15 acid sequences of the invention, as set forth in the Sequence Listing, correspond to the *Corynebacterium glutamicum* PTS DNAs of the invention. This DNA comprises sequences encoding PTS proteins (*i.e.*, the "coding region", indicated in each odd-numbered SEQ ID NO: sequence in the Sequence Listing), as well as 5' untranslated sequences and 3' untranslated sequences, also indicated in each odd-numbered SEQ ID  
20 NO: in the Sequence Listing. Alternatively, the nucleic acid molecule can comprise only the coding region of any of the nucleic acid sequences of the Sequence Listing.

For the purposes of this application, it will be understood that each of the nucleic acid and amino acid sequences set forth in the Sequence Listing has an identifying RXA, RXN, RXS, or RXC number having the designation "RXA", "RXN", "RXS", or "RXC"  
25 followed by 5 digits (*i.e.*, RXA01503, RXN01299, RXS00315, or RXC00953). Each of the nucleic acid sequences comprises up to three parts: a 5' upstream region, a coding region, and a downstream region. Each of these three regions is identified by the same RXA, RXN, RXS, or RXC designation to eliminate confusion. The recitation "one of the odd-numbered sequences of the Sequence Listing" then, refers to any of the nucleic  
30 acid sequences in the Sequence Listing, which may be also be , distinguished by their differing RXA, RXN, RXS, or RXC designations. The coding region of each of these sequences is translated into a corresponding amino acid sequence, which is also set forth

in the Sequence Listing, as an even-numbered SEQ ID NO: immediately following the corresponding nucleic acid sequence. For example, the coding region for RXA02229 is set forth in SEQ ID NO:1, while the amino acid sequence which it encodes is set forth as SEQ ID NO:2. The sequences of the nucleic acid molecules of the invention are

5 identified by the same RXA, RXN, RXS, or RXC designations as the amino acid molecules which they encode, such that they can be readily correlated. For example, the amino acid sequences designated RXA01503, RXN01299, RXS00315, and RXC00953 are translations of the coding regions of the nucleotide sequence of nucleic acid molecules RXA01503, RXN01299, RXS00315, and RXC00953, respectively. The

10 correspondence between the RXA, RXN, RXS, and RXC nucleotide and amino acid sequences of the invention and their assigned SEQ ID NOs, is set forth in Table 1. For example, as set forth in Table 1, the nucleotide sequence of RXN01299 is SEQ ID NO: 7, and the corresponding amino acid sequence is SEQ ID NO:8.

Several of the genes of the invention are "F-designated genes". An F-designated

15 gene includes those genes set forth in Table 1 which have an 'F' in front of the RXA, RXN, RXS, or RXC designation. For example, SEQ ID NO:3, designated, as indicated on Table 1, as "F RXA00315", is an F-designated gene, as are SEQ ID NOs: 9, 11, and 13 (designated on Table 1 as "F RXA01299", "F RXA01883", and "F RXA01889", respectively).

20 In one embodiment, the nucleic acid molecules of the present invention are not intended to include *C. glutamicum* those compiled in Table 2. In the case of the dapD gene, a sequence for this gene was published in Wehrmann, A., *et al.* (1998) *J. Bacteriol.* 180(12): 3159-3165. However, the sequence obtained by the inventors of the present application is significantly longer than the published version. It is believed that

25 the published version relied on an incorrect start codon, and thus represents only a fragment of the actual coding region.

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which is a complement of one of the nucleotide sequences of the invention (*e.g.*, a sequence of an odd-numbered SEQ ID

30 NO: of the Sequence Listing), or a portion thereof. A nucleic acid molecule which is complementary to one of the nucleotide sequences of the invention is one which is sufficiently complementary to one of the nucleotide sequences shown in the Sequence

- 26 -

Listing (*e.g.*, the sequence of an odd-numbered SEQ ID NO:) such that it can hybridize to one of the nucleotide sequences of the invention, thereby forming a stable duplex.

In still another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleotide sequence which is at least about 50%, 51%, 52%, 53%,  
5 54%, 55%, 56%, 57%, 58%, 59%, or 60%, preferably at least about 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, or 70%, more preferably at least about 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, or 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, or 90%, or 91%, 92%, 93%, 94%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more homologous to a nucleotide sequence of the  
10 invention (*e.g.*, a sequence of an odd-numbered SEQ ID NO: of the Sequence Listing), or a portion thereof. Ranges and identity values intermediate to the above-recited ranges, (*e.g.*, 70-90% identical or 80-95% identical) are also intended to be encompassed by the present invention. For example, ranges of identity values using a combination of any of the above values recited as upper and/or lower limits are intended  
15 to be included. In an additional preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleotide sequence which hybridizes, *e.g.*, hybridizes under stringent conditions, to one of the nucleotide sequences of the invention (*e.g.*, a sequence of an odd-numbered SEQ ID NO: of the Sequence Listing), or a portion thereof.

20 Moreover, the nucleic acid molecule of the invention can comprise only a portion of the coding region of the sequence of one of the odd-numbered SEQ ID NOs of the Sequence Listing, for example a fragment which can be used as a probe or primer or a fragment encoding a biologically active portion of a PTS protein. The nucleotide sequences determined from the cloning of the PTS genes from *C. glutamicum* allows for  
25 the generation of probes and primers designed for use in identifying and/or cloning PTS homologues in other cell types and organisms, as well as PTS homologues from other *Corynebacteria* or related species. The probe/primer typically comprises substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, preferably about  
30 25, more preferably about 40, 50 or 75 consecutive nucleotides of a sense strand of one of the nucleotide sequences of the invention (*e.g.*, a sequence of one of the odd-numbered SEQ ID NOs of the Sequence Listing, an anti-sense sequence of one of these

- 27 -

sequences, or naturally occurring mutants thereof. Primers based on a nucleotide sequence of the invention can be used in PCR reactions to clone PTS homologues. Probes based on the PTS nucleotide sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In preferred  
5   embodiments, the probe further comprises a label group attached thereto, *e.g.* the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells which misexpress a PTS protein, such as by measuring a level of a PTS-encoding nucleic acid in a sample of cells *e.g.*, detecting PTS mRNA levels or determining  
10   whether a genomic PTS gene has been mutated or deleted.

In one embodiment, the nucleic acid molecule of the invention encodes a protein or portion thereof which includes an amino acid sequence which is sufficiently homologous to an amino acid sequence of the invention (*e.g.*, a sequence of an even-numbered SEQ ID NO of the Sequence Listing), such that the protein or portion thereof  
15   maintains the ability to participate in the transport of high-energy carbon molecules (such as glucose) into *C. glutamicum*, and may also participate in one or more intracellular signal transduction pathways. As used herein, the language "sufficiently homologous" refers to proteins or portions thereof which have amino acid sequences which include a minimum number of identical or equivalent (*e.g.*, an amino acid residue  
20   which has a similar side chain as an amino acid residue in a sequence of one of the even-numbered SEQ ID NOs of the Sequence Listing) amino acid residues to an amino acid sequence of the invention such that the protein or portion thereof is capable of transporting high-energy carbon-containing molecules such as glucose into *C. glutamicum*, and may also participate in intracellular signal transduction in this  
25   microorganism. Protein members of such metabolic pathways, as described herein, function to transport high-energy carbon-containing molecules such as glucose into *C. glutamicum*, and may also participate in intracellular signal transduction in this microorganism. Examples of such activities are also described herein. Thus, "the function of a PTS protein" contributes to the overall functioning and/or regulation of one  
30   or more phosphoenolpyruvate-based sugar transport pathway, and /or contributes, either directly or indirectly, to the yield, production, and/or efficiency of production of one or more fine chemicals. Examples of PTS protein activities are set forth in Table 1.



In another embodiment, the protein is at least about 50-60%, preferably at least about 60-70%, and more preferably at least about 70-80%, 80-90%, 90-95%, and most preferably at least about 96%, 97%, 98%, 99% or more homologous to an entire amino acid sequence of the invention (*e.g.*, a sequence of an even-numbered SEQ ID NO: of the Sequence Listing).

Portions of proteins encoded by the PTS nucleic acid molecules of the invention are preferably biologically active portions of one of the PTS proteins. As used herein, the term "biologically active portion of a PTS protein" is intended to include a portion, *e.g.*, a domain/motif, of a PTS protein that is capable of transporting high-energy carbon-containing molecules such as glucose into *C. glutamicum*, or of participating in intracellular signal transduction in this microorganism, or has an activity as set forth in Table 1. To determine whether a PTS protein or a biologically active portion thereof can participate in the transportation of high-energy carbon-containing molecules such as glucose into *C. glutamicum*, or can participate in intracellular signal transduction in this microorganism, an assay of enzymatic activity may be performed. Such assay methods are well known to those of ordinary skill in the art, as detailed in Example 8 of the Exemplification.

Additional nucleic acid fragments encoding biologically active portions of a PTS protein can be prepared by isolating a portion of one of the amino acid sequences of the invention (*e.g.*, a sequence of an even-numbered SEQ ID NO: of the Sequence Listing), expressing the encoded portion of the PTS protein or peptide (*e.g.*, by recombinant expression *in vitro*) and assessing the activity of the encoded portion of the PTS protein or peptide.

The invention further encompasses nucleic acid molecules that differ from one of the nucleotide sequences of the invention (*e.g.*, a sequence of an odd-numbered SEQ ID NO: of the Sequence Listing) (and portions thereof) due to degeneracy of the genetic code and thus encode the same PTS protein as that encoded by the nucleotide sequences of the invention. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in the Sequence Listing (*e.g.*, an even-numbered SEQ ID NO:). In a still further embodiment, the nucleic acid molecule of the invention encodes a full length *C.*

- 29 -

*glutamicum* protein which is substantially homologous to an amino acid sequence of the invention (encoded by an open reading frame shown in an odd-numbered SEQ ID NO: of the Sequence Listing).

It will be understood by one of ordinary skill in the art that in one embodiment the sequences of the invention are not meant to include the sequences of the prior art, such as those Genbank sequences set forth in Tables 2 or 4 which were available prior to the present invention. In one embodiment, the invention includes nucleotide and amino acid sequences having a percent identity to a nucleotide or amino acid sequence of the invention which is greater than that of a sequence of the prior art (*e.g.*, a Genbank sequence (or the protein encoded by such a sequence) set forth in Tables 2 or 4). For example, the invention includes a nucleotide sequence which is greater than and/or at least 44% identical to the nucleotide sequence designated RXA01503 (SEQ ID NO:5), a nucleotide sequence which is greater than and/or at least 41% identical to the nucleotide sequence designated RXA00951 (SEQ ID NO:15), and a nucleotide sequence which is greater than and/or at least 38% identical to the nucleotide sequence designated RXA01300 (SEQ ID NO:21). One of ordinary skill in the art would be able to calculate the lower threshold of percent identity for any given sequence of the invention by examining the GAP-calculated percent identity scores set forth in Table 4 for each of the three top hits for the given sequence, and by subtracting the highest GAP-calculated percent identity from 100 percent. One of ordinary skill in the art will also appreciate that nucleic acid and amino acid sequences having percent identities greater than the lower threshold so calculated (*e.g.*, at least 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, or 60%, preferably at least about 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, or 70%, more preferably at least about 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, or 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, or 90%, or 91%, 92%, 93%, 94%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more identical) are also encompassed by the invention.

In addition to the *C. glutamicum* PTS nucleotide sequences set forth in the Sequence Listing as odd-numbered SEQ ID NOs, it will be appreciated by those of ordinary skill in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of PTS proteins may exist within a population (*e.g.*, the *C. glutamicum* population). Such genetic polymorphism in the PTS gene may exist among individuals

- 30 -

within a population due to natural variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding a PTS protein, preferably a *C. glutamicum* PTS protein. Such natural variations can typically result in 1-5% variance in the nucleotide sequence of the PTS gene. Any and all such nucleotide variations and resulting amino acid polymorphisms in PTS that are the result of natural variation and that do not alter the functional activity of PTS proteins are intended to be within the scope of the invention.

Nucleic acid molecules corresponding to natural variants and non-*C. glutamicum* homologues of the *C. glutamicum* PTS DNA of the invention can be isolated based on their homology to the *C. glutamicum* PTS nucleic acid disclosed herein using the *C. glutamicum* DNA, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 15 nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising a nucleotide sequence of an odd-numbered SEQ ID NO: of the Sequence Listing. In other embodiments, the nucleic acid is at least 30, 50, 100, 250 or more nucleotides in length. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other. Preferably, the conditions are such that sequences at least about 65%, more preferably at least about 70%, and even more preferably at least about 75% or more homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those of ordinary skill in the art and can be found in Ausubel *et al.*, *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to a nucleotide sequence of the invention corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (*e.g.*,

encodes a natural protein). In one embodiment, the nucleic acid encodes a natural *C. glutamicum* PTS protein.

In addition to naturally-occurring variants of the PTS sequence that may exist in the population, one of ordinary skill in the art will further appreciate that changes can be introduced by mutation into a nucleotide sequence of the invention, thereby leading to changes in the amino acid sequence of the encoded PTS protein, without altering the functional ability of the PTS protein. For example, nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues can be made in a nucleotide sequence of the invention. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of one of the PTS proteins (*e.g.*, an even-numbered SEQ ID NO: of the Sequence Listing ) without altering the activity of said PTS protein, whereas an "essential" amino acid residue is required for PTS protein activity. Other amino acid residues, however, (*e.g.*, those that are not conserved or only semi-conserved in the domain having PTS activity) may not be essential for activity and thus are likely to be amenable to alteration without altering PTS activity.

Accordingly, another aspect of the invention pertains to nucleic acid molecules encoding PTS proteins that contain changes in amino acid residues that are not essential for PTS activity. Such PTS proteins differ in amino acid sequence from a sequence of an even-numbered SEQ ID NO: of the Sequence Listing yet retain at least one of the PTS activities described herein. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 50% homologous to an amino acid sequence of the invention and is capable of transporting high-energy carbon-containing molecules such as glucose into *C. glutamicum*, or of participating in intracellular signal transduction in this microorganism, or has one or more activities set forth in Table 1. Preferably, the protein encoded by the nucleic acid molecule is at least about 50-60% homologous to the amino acid sequence of one of the odd-numbered SEQ ID NOs of the Sequence Listing, more preferably at least about 60-70% homologous to one of these sequences, even more preferably at least about 70-80%, 80-90%, 90-95% homologous to one of these sequences, and most preferably at least about 96%, 97%, 98%, or 99% homologous to one of the amino acid sequences of the invention.

To determine the percent homology of two amino acid sequences (*e.g.*, one of the amino acid sequences of the invention and a mutant form thereof) or of two nucleic acids, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the sequence of one protein or nucleic acid for optimal alignment with the other protein or nucleic acid). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in one sequence (*e.g.*, one of the amino acid sequences of the invention) is occupied by the same amino acid residue or nucleotide as the corresponding position in the other sequence (*e.g.*, a mutant form of the amino acid sequence), then the molecules are homologous at that position (*i.e.*, as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity"). The percent homology between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % homology = # of identical positions/total # of positions x 100).

An isolated nucleic acid molecule encoding a PTS protein homologous to a protein sequence of the invention (*e.g.*, a sequence of an even-numbered SEQ ID NO: of the Sequence Listing) can be created by introducing one or more nucleotide substitutions, additions or deletions into a nucleotide sequence of the invention such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced into one of the nucleotide sequences of the invention by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in a PTS protein is preferably replaced with another amino acid residue from the same

side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a PTS coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for a PTS activity described herein to identify mutants that retain PTS activity. Following mutagenesis of one of the nucleotide  
5 sequence of one of the odd-numbered SEQ ID NOs of the Sequence Listing, the encoded protein can be expressed recombinantly and the activity of the protein can be determined using, for example, assays described herein (see Example 8 of the Exemplification).

In addition to the nucleic acid molecules encoding PTS proteins described above,  
10 another aspect of the invention pertains to isolated nucleic acid molecules which are antisense thereto. An "antisense" nucleic acid comprises a nucleotide sequence which is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a double-stranded DNA molecule or complementary to an mRNA  
sequence. Accordingly, an antisense nucleic acid can hydrogen bond to a sense nucleic  
15 acid. The antisense nucleic acid can be complementary to an entire PTS coding strand, or to only a portion thereof. In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding a  
PTS protein. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues (*e.g.*, the entire coding  
20 region of SEQ ID NO. 5 (RXA01503) comprises nucleotides 1 to 249). In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding PTS. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into  
amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

25 Given the coding strand sequences encoding PTS disclosed herein (*e.g.*, the sequences set forth as odd-numbered SEQ ID NOs in the Sequence Listing), antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick base pairing. The antisense nucleic acid molecule can be complementary to the  
entire coding region of PTS mRNA, but more preferably is an oligonucleotide which is  
30 antisense to only a portion of the coding or noncoding region of PTS mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of PTS mRNA. An antisense oligonucleotide can be, for

- 34 -

example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives and acridine substituted nucleotides can be used. Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (*v*), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (*v*), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)*w*, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a cell or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a PTS protein to thereby inhibit expression of the protein, *e.g.*, by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. The antisense molecule can

be modified such that it specifically binds to a receptor or an antigen expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecule to a peptide or an antibody which binds to a cell surface receptor or antigen. The antisense nucleic acid molecule can also be delivered to cells using the vectors described herein. To achieve  
5 sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong prokaryotic, viral, or eukaryotic promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an  $\alpha$ -anomeric nucleic acid molecule. An  $\alpha$ -anomeric nucleic acid molecule forms  
10 specific double-stranded hybrids with complementary RNA in which, contrary to the usual  $\beta$ -units, the strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-*o*-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) *FEBS Lett.* 215:327-330).

15 In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity which are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (*e.g.*, hammerhead ribozymes (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to  
20 catalytically cleave PTS mRNA transcripts to thereby inhibit translation of PTS mRNA. A ribozyme having specificity for a PTS-encoding nucleic acid can be designed based upon the nucleotide sequence of a PTS DNA disclosed herein (*i.e.*, SEQ ID NO:5 (RXA01503)). For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the  
25 nucleotide sequence to be cleaved in a PTS-encoding mRNA. See, *e.g.*, Cech *et al.* U.S. Patent No. 4,987,071 and Cech *et al.* U.S. Patent No. 5,116,742. Alternatively, PTS mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, *e.g.*, Bartel, D. and Szostak, J.W. (1993) *Science* 261:1411-1418.

30 Alternatively, PTS gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of a PTS nucleotide sequence (*e.g.*, a PTS promoter and/or enhancers) to form triple helical structures that prevent



- 36 -

transcription of a PTS gene in target cells. See generally, Helene, C. (1991) *Anticancer Drug Des.* 6(6):569-84; Helene, C. *et al.* (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher, L.J. (1992) *Bioassays* 14(12):807-15.

## 5 B. Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding a PTS protein (or a portion thereof). As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid",  
10 which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other  
15 vectors (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are  
20 often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

25 The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid sequence to be expressed. Within a recombinant  
30 expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a

- host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell and those which direct expression of the nucleotide sequence only in certain host cells. Preferred regulatory sequences are, for example, promoters such as *cos*-, *tac*-, *trp*-, *tet*-, *trp-tet*-, *lpp*-, *lac*-, *lpp-lac*-, *lacI<sup>q</sup>*-, *T7*-, *T5*-, *T3*-, *gal*-, *trc*-, *ara*-, *SP6*-, *amy*, *SPO2*,  $\lambda$ -*P<sub>R</sub>*- or  $\lambda$  *P<sub>L</sub>*, which are used preferably in bacteria. Additional regulatory sequences are, for example, promoters from yeasts and fungi, such as *ADC1*, *MFA*, *AC*, *P-60*, *CYC1*, *GAPDH*, *TEF*, *rp28*, *ADH*, promoters from plants such as *CaMV/35S*, *SSU*, *OCS*, *lib4*, *usp*, *STLS1*, *B33*, *nos* or *ubiquitin*- or *phaseolin*-promoters. It is also possible to use artificial promoters. It will be appreciated by one of ordinary skill in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (*e.g.*, PTS proteins, mutant forms of PTS proteins, fusion proteins, etc.).
- The recombinant expression vectors of the invention can be designed for expression of PTS proteins in prokaryotic or eukaryotic cells. For example, PTS genes can be expressed in bacterial cells such as *C. glutamicum*, insect cells (using baculovirus expression vectors), yeast and other fungal cells (see Romanos, M.A. *et al.* (1992) "Foreign gene expression in yeast: a review", *Yeast* 8: 423-488; van den Hondel, C.A.M.J.J. *et al.* (1991) "Heterologous gene expression in filamentous fungi" in: More Gene Manipulations in Fungi, J.W. Bennet & L.L. Lasure, eds., p. 396-428: Academic Press: San Diego; and van den Hondel, C.A.M.J.J. & Punt, P.J. (1991) "Gene transfer systems and vector development for filamentous fungi, in: Applied Molecular Genetics of Fungi, Peberdy, J.F. *et al.*, eds., p. 1-28, Cambridge University Press: Cambridge), algae and multicellular plant cells (see Schmidt, R. and Willmitzer, L. (1988) High efficiency *Agrobacterium tumefaciens* -mediated transformation of *Arabidopsis thaliana* leaf and cotyledon explants" *Plant Cell Rep.*: 583-586), or mammalian cells.

Suitable host cells are discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

5           Expression of proteins in prokaryotes is most often carried out with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: 1) to increase expression of recombinant protein;  
10       2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes,  
15       and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase.

          Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith, D.B. and Johnson, K.S. (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant  
20       protein. In one embodiment, the coding sequence of the PTS protein is cloned into a pGEX expression vector to create a vector encoding a fusion protein comprising, from the N-terminus to the C-terminus, GST-thrombin cleavage site-X protein. The fusion protein can be purified by affinity chromatography using glutathione-agarose resin. Recombinant PTS protein unfused to GST can be recovered by cleavage of the fusion  
25       protein with thrombin.

          Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, (1988) *Gene* 69:301-315) pLG338, pACYC184, pBR322, pUC18, pUC19, pKC30, pRep4, pHS1, pHS2, pPLc236, pMBL24, pLG200, pUR290, pIN-III113-B1,  $\lambda$ gt11, pBdCl, and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89 ; and  
30       Pouwels *et al.*, eds. (1985) *Cloning Vectors*. Elsevier: New York ISBN 0 444 904018). Target gene expression from the pTrc vector relies on host RNA polymerase

transcription from a hybrid trp-lac fusion promoter. Target gene expression from the pET 11d vector relies on transcription from a T7 gn10-lac fusion promoter mediated by a coexpressed viral RNA polymerase (T7 gn1). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident  $\lambda$  prophage harboring a T7 gn1 gene under the transcriptional control of the lacUV 5 promoter. For transformation of other varieties of bacteria, appropriate vectors may be selected. For example, the plasmids pIJ101, pIJ364, pIJ702 and pIJ361 are known to be useful in transforming *Streptomyces*, while plasmids pUB110, pC194, or pBD214 are suited for transformation of *Bacillus* species. Several plasmids of use in the transfer of genetic information into *Corynebacterium* include pHM1519, pBL1, pSA77, or pAJ667 (Pouwels *et al.*, eds. (1985) *Cloning Vectors*. Elsevier: New York ISBN 0 444 904018).

One strategy to maximize recombinant protein expression is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in the bacterium chosen for expression, such as *C. glutamicum* (Wada *et al.* (1992) *Nucleic Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the PTS protein expression vector is a yeast expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, (1987) *Embo J.* 6:229-234), 2  $\mu$ , pAG-1, Yep6, Yep13, pEMBLYe23, pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz *et al.*, (1987) *Gene* 54:113-123), and pYES2 (Invitrogen Corporation, San Diego, CA). Vectors and methods for the construction of vectors appropriate for use in other fungi, such as the filamentous fungi, include those detailed in: van den Hondel, C.A.M.J.J. & Punt, P.J. (1991) "Gene transfer systems and vector development for filamentous fungi, in: *Applied Molecular Genetics of Fungi*, J.F. Peberdy, *et al.*, eds., p. 1-28, Cambridge University Press: Cambridge, and Pouwels *et al.*, eds. (1985) *Cloning Vectors*. Elsevier: New York (ISBN 0 444 904018).

Alternatively, the PTS proteins of the invention can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith *et al.* (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39).

In another embodiment, the PTS proteins of the invention may be expressed in unicellular plant cells (such as algae) or in plant cells from higher plants (e.g., the spermatophytes, such as crop plants). Examples of plant expression vectors include those detailed in: Becker, D., Kemper, E., Schell, J. and Masterson, R. (1992) "New plant binary vectors with selectable markers located proximal to the left border", *Plant Mol. Biol.* 20: 1195-1197; and Bevan, M.W. (1984) "Binary *Agrobacterium* vectors for plant transformation", *Nucl. Acid. Res.* 12: 8711-8721, and include pLGV23, pGHIac+, pBIN19, pAK2004, and pDH51 (Pouwels *et al.*, eds. (1985) *Cloning Vectors*. Elsevier: New York ISBN 0 444 904018).

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, B. (1987) *Nature* 329:840) and pMT2PC (Kaufman *et al.* (1987) *EMBO J.* 6:187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. For other suitable expression systems for both prokaryotic and eukaryotic cells see chapters 16 and 17 of Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual. 2nd, ed.*, Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert *et al.* (1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988) *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and

Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji *et al.* (1983) *Cell* 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (*e.g.*, the neurofilament promoter; Byrne and Ruddle (1989) *PNAS* 86:5473-5477), pancreas-specific promoters (Edlund *et al.* (1985) *Science* 230:912-916), and mammary gland-specific promoters (*e.g.*, milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, for example the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the  $\alpha$ -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev.* 3:537-546).

10       The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in a manner which allows for expression (by transcription of the DNA molecule) of an RNA molecule which is antisense to PTS mRNA. Regulatory sequences operatively  
15 linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen which direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid  
20 or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes see Weintraub, H. *et al.*, Antisense RNA as a molecular tool for genetic analysis, *Reviews - Trends in Genetics*, Vol. 1(1) 1986.

25       Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due  
30 to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, a PTS protein can be expressed in bacterial cells such as *C. glutamicum*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to one of ordinary skill in the art. Microorganisms related to *Corynebacterium glutamicum* which may be conveniently used as host cells for the nucleic acid and protein molecules of the invention are set forth in Table 3.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (e.g., linear DNA or RNA (e.g., a linearized vector or a gene construct alone without a vector) or nucleic acid in the form of a vector (e.g., a plasmid, phage, phasmid, phagemid, transposon or other DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual*, 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (e.g., resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker can be introduced into a host cell on the same vector as that encoding a PTS protein or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (e.g., cells that have incorporated the selectable marker gene will survive, while the other cells die).

To create a homologous recombinant microorganism, a vector is prepared which contains at least a portion of a PTS gene into which a deletion, addition or substitution has been introduced to thereby alter, e.g., functionally disrupt, the PTS gene. Preferably, this PTS gene is a *Corynebacterium glutamicum* PTS gene, but it can be a

homologue from a related bacterium or even from a mammalian, yeast, or insect source. In a preferred embodiment, the vector is designed such that, upon homologous recombination, the endogenous PTS gene is functionally disrupted (i.e., no longer encodes a functional protein; also referred to as a "knock out" vector). Alternatively, the vector can be designed such that, upon homologous recombination, the endogenous PTS gene is mutated or otherwise altered but still encodes functional protein (*e.g.*, the upstream regulatory region can be altered to thereby alter the expression of the endogenous PTS protein). In the homologous recombination vector, the altered portion of the PTS gene is flanked at its 5' and 3' ends by additional nucleic acid of the PTS gene to allow for homologous recombination to occur between the exogenous PTS gene carried by the vector and an endogenous PTS gene in a microorganism. The additional flanking PTS nucleic acid is of sufficient length for successful homologous recombination with the endogenous gene. Typically, several kilobases of flanking DNA (both at the 5' and 3' ends) are included in the vector (see *e.g.*, Thomas, K.R., and Capecchi, M.R. (1987) *Cell* 51: 503 for a description of homologous recombination vectors). The vector is introduced into a microorganism (*e.g.*, by electroporation) and cells in which the introduced PTS gene has homologously recombined with the endogenous PTS gene are selected, using art-known techniques.

In another embodiment, recombinant microorganisms can be produced which contain selected systems which allow for regulated expression of the introduced gene. For example, inclusion of a PTS gene on a vector placing it under control of the *lac* operon permits expression of the PTS gene only in the presence of IPTG. Such regulatory systems are well known in the art.

In another embodiment, an endogenous PTS gene in a host cell is disrupted (*e.g.*, by homologous recombination or other genetic means known in the art) such that expression of its protein product does not occur. In another embodiment, an endogenous or introduced PTS gene in a host cell has been altered by one or more point mutations, deletions, or inversions, but still encodes a functional PTS protein. In still another embodiment, one or more of the regulatory regions (*e.g.*, a promoter, repressor, or inducer) of a PTS gene in a microorganism has been altered (*e.g.*, by deletion, truncation, inversion, or point mutation) such that the expression of the PTS gene is modulated. One of ordinary skill in the art will appreciate that host cells containing



more than one of the described PTS gene and protein modifications may be readily produced using the methods of the invention, and are meant to be included in the present invention.

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in  
5 culture, can be used to produce (i.e., express) a PTS protein. Accordingly, the invention further provides methods for producing PTS proteins using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding a PTS protein has been introduced, or into which genome has been introduced a gene encoding a wild-type or  
10 altered PTS protein) in a suitable medium until PTS protein is produced. In another embodiment, the method further comprises isolating PTS proteins from the medium or the host cell.

### *C. Isolated PTS Proteins*

15 Another aspect of the invention pertains to isolated PTS proteins, and biologically active portions thereof. An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material when produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes  
20 preparations of PTS protein in which the protein is separated from cellular components of the cells in which it is naturally or recombinantly produced. In one embodiment, the language "substantially free of cellular material" includes preparations of PTS protein having less than about 30% (by dry weight) of non-PTS protein (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-PTS protein,  
25 still more preferably less than about 10% of non-PTS protein, and most preferably less than about 5% non-PTS protein. When the PTS protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, i.e., culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein  
30 preparation. The language "substantially free of chemical precursors or other chemicals" includes preparations of PTS protein in which the protein is separated from chemical precursors or other chemicals which are involved in the synthesis of the

protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of PTS protein having less than about 30% (by dry weight) of chemical precursors or non-PTS chemicals, more preferably less than about 20% chemical precursors or non-PTS chemicals, still more preferably less than about 10% chemical precursors or non-PTS chemicals, and most preferably less than about 5% chemical precursors or non-PTS chemicals. In preferred embodiments, isolated proteins or biologically active portions thereof lack contaminating proteins from the same organism from which the PTS protein is derived. Typically, such proteins are produced by recombinant expression of, for example, a *C. glutamicum* PTS protein in a microorganism such as *C. glutamicum*.

An isolated PTS protein or a portion thereof of the invention can participate in the transport of high-energy carbon-containing molecules such as glucose into *C. glutamicum*, and may also participate in intracellular signal transduction in this microorganism, or has one or more of the activities set forth in Table 1. In preferred embodiments, the protein or portion thereof comprises an amino acid sequence which is sufficiently homologous to an amino acid sequence of the invention (*e.g.*, a sequence of an even-numbered SEQ ID NO: of the Sequence Listing) such that the protein or portion thereof maintains the ability to transport high-energy carbon-containing molecules such as glucose into *C. glutamicum*, or to participate in intracellular signal transduction in this microorganism. The portion of the protein is preferably a biologically active portion as described herein. In another preferred embodiment, a PTS protein of the invention has an amino acid sequence set forth as an even-numbered SEQ ID NO: of the Sequence Listing. In yet another preferred embodiment, the PTS protein has an amino acid sequence which is encoded by a nucleotide sequence which hybridizes, *e.g.*, hybridizes under stringent conditions, to a nucleotide sequence of the invention (*e.g.*, a sequence of an odd-numbered SEQ ID NO: of the Sequence Listing). In still another preferred embodiment, the PTS protein has an amino acid sequence which is encoded by a nucleotide sequence that is at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, or 60%, preferably at least about 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, or 70%, more preferably at least about 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, or 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, or 90%, or 91%, 92%, 93%, 94%, and even more preferably at least about 95%, 96%, 97%, 98%,

- 46 -

99% or more homologous to one of the nucleic acid sequences of the invention, or a portion thereof. Ranges and identity values intermediate to the above-recited values, (e.g., 70-90% identical or 80-95% identical) are also intended to be encompassed by the present invention. For example, ranges of identity values using a combination of any of the above values recited as upper and/or lower limits are intended to be included. The preferred PTS proteins of the present invention also preferably possess at least one of the PTS activities described herein. For example, a preferred PTS protein of the present invention includes an amino acid sequence encoded by a nucleotide sequence which hybridizes, e.g., hybridizes under stringent conditions, to a nucleotide sequence of the invention, and which can participate in the transport of high-energy carbon-containing molecules such as glucose into *C. glutamicum*, and may also participate in intracellular signal transduction in this microorganism, or which has one or more of the activities set forth in Table 1.

In other embodiments, the PTS protein is substantially homologous to an amino acid sequence of the invention (e.g., a sequence of an even-numbered SEQ ID NO: of the Sequence Listing) and retains the functional activity of the protein of one of the amino acid sequences of the invention yet differs in amino acid sequence due to natural variation or mutagenesis, as described in detail in subsection I above. Accordingly, in another embodiment, the PTS protein is a protein which comprises an amino acid sequence which is at least about 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, or 60%, preferably at least about 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, or 70%, more preferably at least about 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, or 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, or 90%, or 91%, 92%, 93%, 94%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more homologous to an entire amino acid sequence of the invention and which has at least one of the PTS activities described herein. Ranges and identity values intermediate to the above-recited values, (e.g., 70-90% identical or 80-95% identical) are also intended to be encompassed by the present invention. For example, ranges of identity values using a combination of any of the above values recited as upper and/or lower limits are intended to be included. In another embodiment, the invention pertains to a full length *C. glutamicum* protein which is substantially homologous to an entire amino acid sequence of the invention.

Biologically active portions of a PTS protein include peptides comprising amino acid sequences derived from the amino acid sequence of a PTS protein, *e.g.*, an amino acid sequence of an even-numbered SEQ ID NO: of the Sequence Listing or the amino acid sequence of a protein homologous to a PTS protein, which include fewer amino acids than a full length PTS protein or the full length protein which is homologous to a PTS protein, and exhibit at least one activity of a PTS protein. Typically, biologically active portions (peptides, *e.g.*, peptides which are, for example, 5, 10, 15, 20, 30, 35, 36, 37, 38, 39, 40, 50, 100 or more amino acids in length) comprise a domain or motif with at least one activity of a PTS protein. Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the activities described herein. Preferably, the biologically active portions of a PTS protein include one or more selected domains/motifs or portions thereof having biological activity.

PTS proteins are preferably produced by recombinant DNA techniques. For example, a nucleic acid molecule encoding the protein is cloned into an expression vector (as described above), the expression vector is introduced into a host cell (as described above) and the PTS protein is expressed in the host cell. The PTS protein can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques. Alternative to recombinant expression, a PTS protein, polypeptide, or peptide can be synthesized chemically using standard peptide synthesis techniques. Moreover, native PTS protein can be isolated from cells (*e.g.*, endothelial cells), for example using an anti-PTS antibody, which can be produced by standard techniques utilizing a PTS protein or fragment thereof of this invention.

The invention also provides PTS chimeric or fusion proteins. As used herein, a PTS "chimeric protein" or "fusion protein" comprises a PTS polypeptide operatively linked to a non-PTS polypeptide. An "PTS polypeptide" refers to a polypeptide having an amino acid sequence corresponding to PTS, whereas a "non-PTS polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the PTS protein, *e.g.*, a protein which is different from the PTS protein and which is derived from the same or a different organism. Within the fusion protein, the term "operatively linked" is intended to indicate that the PTS polypeptide and the non-PTS polypeptide are fused in-frame to each other. The non-

PTS polypeptide can be fused to the N-terminus or C-terminus of the PTS polypeptide. For example, in one embodiment the fusion protein is a GST-PTS fusion protein in which the PTS sequences are fused to the C-terminus of the GST sequences. Such fusion proteins can facilitate the purification of recombinant PTS proteins. In another  
5 embodiment, the fusion protein is a PTS protein containing a heterologous signal sequence at its N-terminus. In certain host cells (*e.g.*, mammalian host cells), expression and/or secretion of a PTS protein can be increased through use of a heterologous signal sequence.

Preferably, a PTS chimeric or fusion protein of the invention is produced by  
10 standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, for example by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid  
15 undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and reamplified to generate a chimeric  
20 gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel *et al.* John Wiley & Sons: 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide). A PTS-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the PTS protein.

25 Homologues of the PTS protein can be generated by mutagenesis, *e.g.*, discrete point mutation or truncation of the PTS protein. As used herein, the term "homologue" refers to a variant form of the PTS protein which acts as an agonist or antagonist of the activity of the PTS protein. An agonist of the PTS protein can retain substantially the same, or a subset, of the biological activities of the PTS protein. An antagonist of the  
30 PTS protein can inhibit one or more of the activities of the naturally occurring form of the PTS protein, by, for example, competitively binding to a downstream or upstream member of the PTS system which includes the PTS protein. Thus, the *C. glutamicum*

PTS protein and homologues thereof of the present invention may modulate the activity of one or more sugar transport pathways or intracellular signal transduction pathways in which PTS proteins play a role in this microorganism.

In an alternative embodiment, homologues of the PTS protein can be identified  
5 by screening combinatorial libraries of mutants, *e.g.*, truncation mutants, of the PTS protein for PTS protein agonist or antagonist activity. In one embodiment, a variegated library of PTS variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of PTS variants can be produced by, for example, enzymatically ligating a mixture of synthetic  
10 oligonucleotides into gene sequences such that a degenerate set of potential PTS sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (*e.g.*, for phage display) containing the set of PTS sequences therein. There are a variety of methods which can be used to produce libraries of potential PTS homologues from a degenerate oligonucleotide sequence. Chemical synthesis of a  
15 degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential PTS sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, *e.g.*, Narang, S.A. (1983) *Tetrahedron* 39:3; Itakura *et al.* (1984) *Annu. Rev. Biochem.* 53:323; Itakura *et al.* (1984) *Science* 20 198:1056; Ike *et al.* (1983) *Nucleic Acid Res.* 11:477.

In addition, libraries of fragments of the PTS protein coding can be used to generate a variegated population of PTS fragments for screening and subsequent selection of homologues of a PTS protein. In one embodiment, a library of coding  
25 sequence fragments can be generated by treating a double stranded PCR fragment of a PTS coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA which can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by  
30 treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal, C-terminal and internal fragments of various sizes of the PTS protein.

Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of PTS homologues. The most widely used techniques, which are amenable to high through-put analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify PTS homologues (Arkin and Yourvan (1992) *PNAS* 89:7811-7815; Delgrave *et al.* (1993) *Protein Engineering* 6(3):327-331).

In another embodiment, cell based assays can be exploited to analyze a variegated PTS library, using methods well known in the art.

#### *D. Uses and Methods of the Invention*

The nucleic acid molecules, proteins, protein homologues, fusion proteins, primers, vectors, and host cells described herein can be used in one or more of the following methods: identification of *C. glutamicum* and related organisms; mapping of genomes of organisms related to *C. glutamicum*; identification and localization of *C. glutamicum* sequences of interest; evolutionary studies; determination of PTS protein regions required for function; modulation of a PTS protein activity; modulation of the activity of a PTS pathway; and modulation of cellular production of a desired compound, such as a fine chemical.

The PTS nucleic acid molecules of the invention have a variety of uses. First, they may be used to identify an organism as being *Corynebacterium glutamicum* or a close relative thereof. Also, they may be used to identify the presence of *C. glutamicum* or a relative thereof in a mixed population of microorganisms. The invention provides the nucleic acid sequences of a number of *C. glutamicum* genes; by probing the extracted genomic DNA of a culture of a unique or mixed population of microorganisms

under stringent conditions with a probe spanning a region of a *C. glutamicum* gene which is unique to this organism, one can ascertain whether this organism is present.

Although *Corynebacterium glutamicum* itself is nonpathogenic, it is related to pathogenic species, such as *Corynebacterium diphtheriae*. *Corynebacterium*

5 *diphtheriae* is the causative agent of diphtheria, a rapidly developing, acute, febrile infection which involves both local and systemic pathology. In this disease, a local lesion develops in the upper respiratory tract and involves necrotic injury to epithelial cells; the bacilli secrete toxin which is disseminated through this lesion to distal susceptible tissues of the body. Degenerative changes brought about by the inhibition of  
10 protein synthesis in these tissues, which include heart, muscle, peripheral nerves, adrenals, kidneys, liver and spleen, result in the systemic pathology of the disease. Diphtheria continues to have high incidence in many parts of the world, including Africa, Asia, Eastern Europe and the independent states of the former Soviet Union. An ongoing epidemic of diphtheria in the latter two regions has resulted in at least 5,000  
15 deaths since 1990.

In one embodiment, the invention provides a method of identifying the presence or activity of *Corynebacterium diphtheriae* in a subject. This method includes detection of one or more of the nucleic acid or amino acid sequences of the invention (*e.g.*, the sequences set forth as odd-numbered or even-numbered SEQ ID NOs, respectively, in  
20 the Sequence Listing) in a subject, thereby detecting the presence or activity of *Corynebacterium diphtheriae* in the subject. *C. glutamicum* and *C. diphtheriae* are related bacteria, and many of the nucleic acid and protein molecules in *C. glutamicum* are homologous to *C. diphtheriae* nucleic acid and protein molecules, and can therefore be used to detect *C. diphtheriae* in a subject.

25 The nucleic acid and protein molecules of the invention may also serve as markers for specific regions of the genome. This has utility not only in the mapping of the genome, but also for functional studies of *C. glutamicum* proteins. For example, to identify the region of the genome to which a particular *C. glutamicum* DNA-binding protein binds, the *C. glutamicum* genome could be digested, and the fragments incubated  
30 with the DNA-binding protein. Those which bind the protein may be additionally probed with the nucleic acid molecules of the invention, preferably with readily detectable labels; binding of such a nucleic acid molecule to the genome fragment enables the



localization of the fragment to the genome map of *C. glutamicum*, and, when performed multiple times with different enzymes, facilitates a rapid determination of the nucleic acid sequence to which the protein binds. Further, the nucleic acid molecules of the invention may be sufficiently homologous to the sequences of related species such that  
5 these nucleic acid molecules may serve as markers for the construction of a genomic map in related bacteria, such as *Brevibacterium lactofermentum*.

The PTS nucleic acid molecules of the invention are also useful for evolutionary and protein structural studies. The sugar uptake system in which the molecules of the invention participate are utilized by a wide variety of bacteria; by comparing the  
10 sequences of the nucleic acid molecules of the present invention to those encoding similar enzymes from other organisms, the evolutionary relatedness of the organisms can be assessed. Similarly, such a comparison permits an assessment of which regions of the sequence are conserved and which are not, which may aid in determining those regions of the protein which are essential for the functioning of the enzyme. This type  
15 of determination is of value for protein engineering studies and may give an indication of what the protein can tolerate in terms of mutagenesis without losing function.

Manipulation of the PTS nucleic acid molecules of the invention may result in the production of PTS proteins having functional differences from the wild-type PTS proteins. These proteins may be improved in efficiency or activity, may be present in  
20 greater numbers in the cell than is usual, or may be decreased in efficiency or activity.

The invention provides methods for screening molecules which modulate the activity of a PTS protein, either by interacting with the protein itself or a substrate or binding partner of the PTS protein, or by modulating the transcription or translation of a PTS nucleic acid molecule of the invention. In such methods, a microorganism  
25 expressing one or more PTS proteins of the invention is contacted with one or more test compounds, and the effect of each test compound on the activity or level of expression of the PTS protein is assessed.

The PTS molecules of the invention may be modified such that the yield, production, and/or efficiency of production of one or more fine chemicals is improved.  
30 For example, by modifying a PTS protein involved in the uptake of glucose such that it is optimized in activity, the quantity of glucose uptake or the rate at which glucose is translocated into the cell may be increased. The breakdown of glucose and other sugars

within the cell provides energy that may be used to drive energetically unfavorable biochemical reactions, such as those involved in the biosynthesis of fine chemicals. This breakdown also provides intermediate and precursor molecules necessary for the biosynthesis of certain fine chemicals, such as amino acids, vitamins and cofactors. By  
5 increasing the amount of intracellular high-energy carbon molecules through modification of the PTS molecules of the invention, one may therefore increase both the energy available to perform metabolic pathways necessary for the production of one or more fine chemicals, and also the intracellular pools of metabolites necessary for such production. Conversely, by decreasing the importation of a sugar whose breakdown  
10 products include a compound which is used solely in metabolic pathways which compete with pathways utilized for the production of a desired fine chemical for enzymes, cofactors, or intermediates, one may downregulate this pathway and thus perhaps increase production through the desired biosynthetic pathway.

Further, the PTS molecules of the invention may be involved in one or more  
15 intracellular signal transduction pathways which may affect the yields and/or rate of production of one or more fine chemical from *C. glutamicum*. For example, proteins necessary for the import of one or more sugars from the extracellular medium (e.g., HPr, Enzyme I, or a member of an Enzyme II complex) are frequently posttranslationally modified upon the presence of a sufficient quantity of the sugar in the cell, such that  
20 they are no longer able to import that sugar. An example of this occurs in *E. coli*, where high intracellular levels of fructose 1,6-bisphosphate result in the phosphorylation of HPr at serine-46, upon which this molecule is no longer able to participate in the transport of any sugar. While this intracellular level of sugar at which the transport system is shut off may be sufficient to sustain the normal functioning of the cell, it may  
25 be limiting for the overproduction of the desired fine chemical. Thus, it may be desirable to modify the PTS proteins of the invention such that they are no longer responsive to such negative regulation, thereby permitting greater intracellular concentrations of one or more sugars to be achieved, and, by extension, more efficient production or greater yields of one or more fine chemicals from organisms containing  
30 such mutant PTS proteins.

This aforementioned list of mutagenesis strategies for PTS proteins to result in increased yields of a desired compound is not meant to be limiting; variations on these

mutagenesis strategies will be readily apparent to one of ordinary skill in the art. By these mechanisms, the nucleic acid and protein molecules of the invention may be utilized to generate *C. glutamicum* or related strains of bacteria expressing mutated PTS nucleic acid and protein molecules such that the yield, production, and/or efficiency of  
5 production of a desired compound is improved. This desired compound may be any natural product of *C. glutamicum*, which includes the final products of biosynthesis pathways and intermediates of naturally-occurring metabolic pathways, as well as molecules which do not naturally occur in the metabolism of *C. glutamicum*, but which are produced by a *C. glutamicum* strain of the invention.

10 This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patent applications, patents, published patent applications, Tables, and the Sequence Listing cited throughout this application are hereby incorporated by reference.

TABLE 1: Genes Included in the Invention

## PHOSPHOENOLPYRUVATE: SUGAR PHOSPHOTRANSFERASE SYSTEM

<u>Nucleotide SEQ ID NO</u>	<u>Amino Acid SEQ ID NO</u>	<u>Identification Code</u>	<u>Contig.</u>	<u>NT Start</u>	<u>NT Stop</u>	<u>Function</u>
1	2	RXS00315				PTS SYSTEM, SUCROSE-SPECIFIC IABC COMPONENT (EIIABC-SCR) (SUCROSE- PERMEASE IABC COMPONENT)(PHOSPHOTRANSFERASE ENZYME II, ABC COMPONENT) (EC 2.7.1.69)
3	4	F RXA00315	GR00053	6537	5452	PTS SYSTEM, BETA-GLUCOSIDES-SPECIFIC IABC COMPONENT (EIIABC-BGL) (BETA- GLUCOSIDES- PERMEASE IABC COMPONENT) (PHOSPHOTRANSFERASE ENZYME II, ABC COMPONENT) (EC 2.7.1.69)
5	6	RXA01503	GR00424	10392	10640	PTS SYSTEM, BETA-GLUCOSIDES-SPECIFIC IABC COMPONENT (EIIABC-BGL) (BETA- GLUCOSIDES- PERMEASE IABC COMPONENT) (PHOSPHOTRANSFERASE ENZYME II, ABC COMPONENT) (EC 2.7.1.69)
7	8	RXN01299	VW0068	11954	9891	PTS SYSTEM, FRUCTOSE-SPECIFIC IIBC COMPONENT (EC 2.7.1.69)
9	10	F RXA01299	GR00375	6	446	PTS SYSTEM, FRUCTOSE-SPECIFIC IIBC COMPONENT (EC 2.7.1.69)
11	12	F RXA01883	GR00538	2154	2633	PTS SYSTEM, FRUCTOSE-SPECIFIC IIBC COMPONENT (EC 2.7.1.69)
13	14	F RXA01889	GR00540	77	631	PTS SYSTEM, FRUCTOSE-SPECIFIC IIBC COMPONENT (EC 2.7.1.69)
15	16	RXA00951	GR00261	564	172	PTS SYSTEM, FRUCTOSE-SPECIFIC IIBC COMPONENT (EC 2.7.1.69)
17	18	RXN01244	VW0068	14141	15844	PTS SYSTEM, MANNITOL (CRYPTIC) -SPECIFIC IIA COMPONENT (EIIA-(C)MTL) (MANNITOL (CRYPTIC)- PERMEASE IIA COMPONENT) (PHOSPHOTRANSFERASE ENZYME II, A COMPONENT) (EC 2.7.1.69)
19	20	F RXA01244	GR00359	4837	3329	PHOSPHOENOLPYRUVATE-PROTEIN PHOSPHOTRANSFERASE (EC 2.7.3.9)
21	22	RXA01300	GR00375	637	903	PHOSPHOENOLPYRUVATE-PROTEIN PHOSPHOTRANSFERASE (EC 2.7.3.9)
23	24	RXN03002	VW0236	1437	1844	PHOSPHOCARRIER PROTEIN HPR
25	26	RXC00953	VW0260	1834	1082	PTS SYSTEM, MANNITOL (CRYPTIC) -SPECIFIC IIA COMPONENT (EIIA-(C)MTL) (MANNITOL (CRYPTIC)-PERMEASE IIA COMPONENT) (PHOSPHOTRANSFERASE ENZYME II, A COMPONENT) (EC 2.7.1.69)
27	28	RXC03001				Membrane Spanning Protein involved in PTS system
29	30	RXN01943	VW0120	4326	6374	Membrane Spanning Protein involved in PTS system
31	32	F RXA02191	GR00642	3395	4633	PTS SYSTEM, GLUCOSE-SPECIFIC IABC COMPONENT (EC 2.7.1.69)
33	34	F RXA01943	GR00557	3944	3540	PHOSPHOENOLPYRUVATE SUGAR PHOSPHOTRANSFERASE crr gene; phosphotransferase system glucose-specific enzyme III

TABLE 2 - Excluded Genes

GenBank™ Accession No.	Gene Name	Gene Function	Reference
A09073	ppg	Phosphoenol pyruvate carboxylase	Bachmann, B. et al. "DNA fragment coding for phosphoenolpyruvate carboxylase, recombinant DNA carrying said fragment, strains carrying the recombinant DNA and method for producing L-aminino acids using said strains," Patent: EP 0358940-A 3 03/21/90
A45579, A45581, A45583, A45585 A45587		Threonine dehydratase	Moeckel, B. et al. "Production of L-isoleucine by means of recombinant micro-organisms with deregulated threonine dehydratase," Patent: WO 9519442-A 5 07/20/95
AB003132	murC; ftsQ; ftsZ		Kobayashi, M. et al. "Cloning, sequencing, and characterization of the ftsZ gene from coryneform bacteria," <i>Biochem. Biophys. Res. Commun.</i> , 236(2):383-388 (1997)
AB015023	murC; ftsQ		Wachi, M. et al. "A murC gene from Coryneform bacteria," <i>Appl. Microbiol. Biotechnol.</i> , 51(2):223-228 (1999)
AB018530	ftsR		Kimura, E. et al. "Molecular cloning of a novel gene, ftsR, which rescues the detergent sensitivity of a mutant derived from <i>Brevibacterium lacticfermentum</i> ," <i>Biosci. Biotechnol. Biochem.</i> , 60(10):1565-1570 (1996)
AB018531	ftsR1; ftsR2		
AB020624	murI	D-glutamate racemase	
AB023377	tkl	transketolase	
AB024708	glbB; gltD	Glutamine 2-oxoglutarate aminotransferase large and small subunits	
AB025424	acn	aconitase	
AB027714	rep	Replication protein	
AB027715	rep; aad	Replication protein; aminoglycoside adenyltransferase	
AF005242	argC	N-acetylglutamate-5-semialdehyde dehydrogenase	
AF005635	glnA	Glutamine synthetase	
AF030405	hisF	cyclase	
AF030520	argG	Argininosuccinate synthetase	
AF031518	argF	Omithine carbamoyltransferase	
AF036932	aroD	3-dehydroquininate dehydratase	
AF038548	pyc	Pyruvate carboxylase	

Table 2 (continued)

	dcIAE; apt; rel	Dipeptide-binding protein; adenine phosphoribosyltransferase; GTP pyrophosphokinase	Wehmeier, L. et al. "The role of the Corynebacterium glutamicum rel gene in (p)ppGpp metabolism," <i>Microbiology</i> , 144:1853-1862 (1998)
AF038651			
AF041436	argR	Arginine repressor	
AF045998	impA	Inositol monophosphate phosphatase	
AF048764	argH	Argininosuccinate lyase	
AF049897	argC; argI; argB; argD; argF; argR; argG; argH	N-acetylglutamylphosphate reductase; ornithine acetyltransferase; N-acetylglutamate kinase; acetylornithine transaminase; ornithine carbamoyltransferase; arginine repressor; argininosuccinate synthase; argininosuccinate lyase	
AF050109	inhA	Enoyl-acyl carrier protein reductase	
AF050166	hisG	ATP phosphoribosyltransferase	
AF051846	hisA	Phosphoribosylformimino-5-amino-1-phosphoribosyl-4-imidazolecarboxamide isomerase	
AF052652	metA	Homoserine O-acetyltransferase	Park, S. et al. "Isolation and analysis of metA, a methionine biosynthetic gene encoding homoserine acetyltransferase in <i>Corynebacterium glutamicum</i> ," <i>Mol. Cells</i> , 8(3):286-294 (1998)
AF053071	aroB	Dehydroquinase synthetase	
AF060558	hisH	Glutamine amidotransferase	
AF086704	hisE	Phosphoribosyl-ATP-pyrophosphohydrolase	
AF114233	aroA	5-enolpyruvylshikimate 3-phosphate synthase	
AF116184	panD	L-aspartate-alpha-decarboxylase precursor	Dusch, N. et al. "Expression of the <i>Corynebacterium glutamicum</i> panD gene encoding L-aspartate-alpha-decarboxylase leads to pantothenate overproduction in <i>Escherichia coli</i> ," <i>Appl. Environ. Microbiol.</i> , 65(4):1530-1539 (1999)
AF124518	aroD; aroE	3-dehydroquinase; shikimate dehydrogenase	
AF124600	aroC; aroK; aroB; pepQ	Chorismate synthase; shikimate kinase; 3-dehydroquinase synthase; putative cytoplasmic peptidase	
AF145897	inhA		
AF145898	inhA		

Table 2 (continued)

AJ001436	ectP	Transport of ectoine, glycine betaine, proline	Peter, H. et al. "Corynebacterium glutamicum is equipped with four secondary carriers for compatible solutes: Identification, sequencing, and characterization of the proline/ectoine uptake system, ProP, and the ectoine/proline/glycine betaine carrier, EctP," <i>J. Bacteriol.</i> , 180(22):6005-6012 (1998)
AJ004934	dapD	Tetrahydrodipicolinate succinylase (incomplete)	Wehrmann, A. et al. "Different modes of diamino pimelate synthesis and their role in cell wall integrity: A study with <i>Corynebacterium glutamicum</i> ," <i>J. Bacteriol.</i> , 180(12):3159-3165 (1998)
AJ007732	ppc; secG; amt; ocd; soxA	Phosphoenolpyruvate-carboxylase; ?; high affinity ammonium uptake protein; putative ornithine-cyclodecarboxylase; sarcosine oxidase	
AJ010319	ftsY, glnB, glnD; srp; amtP	Involved in cell division; PII protein; uridylyltransferase (uridylyl-removing enzyme); signal recognition particle; low affinity ammonium uptake protein	Jakoby, M. et al. "Nitrogen regulation in <i>Corynebacterium glutamicum</i> ; Isolation of genes involved in biochemical characterization of corresponding proteins," <i>FEMS Microbiol.</i> , 173(2):303-310 (1999)
AJ132968	cat	Chloramphenicol acetyl transferase	
AJ224946	mgo	L-malate: quinone oxidoreductase	Molenaar, D. et al. "Biochemical and genetic characterization of the membrane-associated malate dehydrogenase (acceptor) from <i>Corynebacterium glutamicum</i> ," <i>Eur. J. Biochem.</i> , 254(2):395-403 (1998)
AJ238250	ndh	NADH dehydrogenase	
AJ238703	porA	Porin	Lichtinger, T. et al. "Biochemical and biophysical characterization of the cell wall porin of <i>Corynebacterium glutamicum</i> : The channel is formed by a low molecular mass polypeptide," <i>Biochemistry</i> , 37(43):15024-15032 (1998)
D17429		Transposable element IS31831	Vertes et al. "Isolation and characterization of IS31831, a transposable element from <i>Corynebacterium glutamicum</i> ," <i>Mol. Microbiol.</i> , 11(4):739-746 (1994)
D84102	odhA	2-oxoglutarate dehydrogenase	Usuda, Y. et al. "Molecular cloning of the <i>Corynebacterium glutamicum</i> (Brevibacterium lactofermentum AJ12036) odhA gene encoding a novel type of 2-oxoglutarate dehydrogenase," <i>Microbiology</i> , 142:3347-3354 (1996)
E01358	hdh; hk	Homoserine dehydrogenase; homoserine kinase	Katsumata, R. et al. "Production of L-threonine and L-isoleucine," Patent: JP 1987232392-A 1 10/12/87
E01359		Upstream of the start codon of homoserine kinase gene	Katsumata, R. et al. "Production of L-threonine and L-isoleucine," Patent: JP 1987232392-A 2 10/12/87
E01375		Tryptophan operon	
E01376	trpL; trpE	Leader peptide; anthranilate synthase	Matsui, K. et al. "Tryptophan operon, peptide and protein coded thereby, utilization of tryptophan operon gene expression and production of tryptophan," Patent: JP 1987244382-A 1 10/24/87

Table 2 (continued)

E01377	Promoter and operator regions of tryptophan operon	Matsui, K. et al. "Tryptophan operon, peptide and protein coded thereby, utilization of tryptophan operon gene expression and production of tryptophan," Patent: JP 1987244382-A 1 10/24/87
E03937	Biotin-synthase	Hatakeyama, K. et al. "DNA fragment containing gene capable of coding biotin synthetase and its utilization," Patent: JP 1992278088-A 1 10/02/92
E04040	Diamino pelargonic acid aminotransferase	Kohama, K. et al. "Gene coding diaminopelargonic acid aminotransferase and deshiobiotin synthetase and its utilization," Patent: JP 1992330284-A 1 11/18/92
E04041	Deshiobiotinsynthetase	Kohama, K. et al. "Gene coding diaminopelargonic acid aminotransferase and deshiobiotin synthetase and its utilization," Patent: JP 1992330284-A 1 11/18/92
E04307	Flavum aspartase	Kurusu, Y. et al. "Gene DNA coding aspartase and utilization thereof," Patent: JP 1993030977-A 1 02/09/93
E04376	Isocitric acid lyase	Katsumata, R. et al. "Gene manifestation controlling DNA," Patent: JP 1993056782-A 3 03/09/93
E04377	Isocitric acid lyase N-terminal fragment	Katsumata, R. et al. "Gene manifestation controlling DNA," Patent: JP 1993056782-A 3 03/09/93
E04484	Prephenate dehydratase	Sotouchi, N. et al. "Production of L-phenylalanine by fermentation," Patent: JP 1993076352-A 2 03/30/93
E05108	Aspartokinase	Fugono, N. et al. "Gene DNA coding Aspartokinase and its use," Patent: JP 1993184366-A 1 07/27/93
E05112	Dihydro-dipichorinate synthetase	Hatakeyama, K. et al. "Gene DNA coding dihydrodipicolinic acid synthetase and its use," Patent: JP 1993184371-A 1 07/27/93
E05776	Diaminopimelic acid dehydrogenase	Kobayashi, M. et al. "Gene DNA coding Diaminopimelic acid dehydrogenase and its use," Patent: JP 1993284970-A 1 11/02/93
E05779	Threonine synthase	Kohama, K. et al. "Gene DNA coding threonine synthase and its use," Patent: JP 1993284972-A 1 11/02/93
E06110	Prephenate dehydratase	Kikuchi, T. et al. "Production of L-phenylalanine by fermentation method," Patent: JP 1993344881-A 1 12/27/93
E06111	Mutated Prephenate dehydratase	Kikuchi, T. et al. "Production of L-phenylalanine by fermentation method," Patent: JP 1993344881-A 1 12/27/93
E06146	Acetohydroxy acid synthetase	Inui, M. et al. "Gene capable of coding Acetohydroxy acid synthetase and its use," Patent: JP 1993344893-A 1 12/27/93
E06825	Aspartokinase	Sugimoto, M. et al. "Mutant aspartokinase gene," patent: JP 1994062866-A 1 03/08/94
E06826	Mutated aspartokinase alpha subunit	Sugimoto, M. et al. "Mutant aspartokinase gene," patent: JP 1994062866-A 1 03/08/94



Table 2 (continued)

E06827		Mutated aspartokinase alpha subunit	Sugimoto, M. et al. "Mutant aspartokinase gene," patent: JP 1994062866-A 1 03/08/94
E07701	secY		Honno, N. et al. "Gene DNA participating in integration of membrane protein to membrane," Patent: JP 1994169780-A 1 06/21/94
E08177		Aspartokinase	Sato, Y. et al. "Genetic DNA capable of coding Aspartokinase released from feedback inhibition and its utilization," Patent: JP 1994261766-A 1 09/20/94
E08178, E08179, E08180, E08181, E08182		Feedback inhibition-released Aspartokinase	Sato, Y. et al. "Genetic DNA capable of coding Aspartokinase released from feedback inhibition and its utilization," Patent: JP 1994261766-A 1 09/20/94
E08232		Acetohydroxy-acid isomeroreductase	Inui, M. et al. "Gene DNA coding acetohydroxy acid isomeroreductase," Patent: JP 1994277067-A 1 10/04/94
E08234	secE		Asai, Y. et al. "Gene DNA coding for translocation machinery of protein," Patent: JP 1994277073-A 1 10/04/94
E08643		FT aminotransferase and desthiobiotin synthetase promoter region	Hatakeyama, K. et al. "DNA fragment having promoter function in coryneform bacterium," Patent: JP 1995031476-A 1 02/03/95
E08646		Biotin synthetase	Hatakeyama, K. et al. "DNA fragment having promoter function in coryneform bacterium," Patent: JP 1995031476-A 1 02/03/95
E08649		Aspartase	Kohama, K. et al. "DNA fragment having promoter function in coryneform bacterium," Patent: JP 1995031478-A 1 02/03/95
E08900		Dihydrodipicolinate reductase	Madori, M. et al. "DNA fragment containing gene coding Dihydrodipicolinate acid reductase and utilization thereof," Patent: JP 1995055578-A 1 03/20/95
E08901		Diaminopimelic acid decarboxylase	Madori, M. et al. "DNA fragment containing gene coding Diaminopimelic acid decarboxylase and utilization thereof," Patent: JP 1995055579-A 1 03/20/95
E12594		Serine hydroxymethyltransferase	Hatakeyama, K. et al. "Production of L-tryptophan," Patent: JP 1997028391-A 1 02/04/97
E12760, E12759, E12758		transposase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E12764		Arginyl-tRNA synthetase; diaminopimelic acid decarboxylase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E12767		Dihydrodipicolinic acid synthetase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E12770		aspartokinase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97
E12773		Dihydrodipicolinic acid reductase	Moriya, M. et al. "Amplification of gene using artificial transposon," Patent: JP 1997070291-A 03/18/97

Table 2 (continued)

E13655		Glucose-6-phosphate dehydrogenase	Hatakeyama, K. et al. "Glucose-6-phosphate dehydrogenase and DNA capable of coding the same," Patent: JP 1997224661-A 1 09/02/97
L01508	IlvA	Threonine dehydratase	Moockel, B. et al. "Functional and structural analysis of the threonine dehydratase of Corynebacterium glutamicum," <i>J. Bacteriol.</i> , 174:8065-8072 (1992)
L07603	EC 4.2.1.15	3-deoxy-D-arabinoheptulosonate-7-phosphate synthase	Chen, C. et al. "The cloning and nucleotide sequence of Corynebacterium glutamicum 3-deoxy-D-arabinoheptulosonate-7-phosphate synthase gene," <i>FEMS Microbiol. Lett.</i> , 107:223-230 (1993)
L09232	IlvB; ilvN; ilvC	Acetohydroxy acid synthase large subunit; Acetohydroxy acid synthase small subunit; Acetohydroxy acid isomeroreductase	Keilhauer, C. et al. "Isolation and synthesis in Corynebacterium glutamicum: molecular analysis of the ilvB-ilvN-ilvC operon," <i>J. Bacteriol.</i> , 175(17):5595-5603 (1993)
L18874	PtsM	Phosphoenolpyruvate sugar phosphotransferase	Fouet, A. et al. "Bacillus subtilis sucrose-specific enzyme II of the phosphotransferase system: expression in Escherichia coli and homology to enzymes II from enteric bacteria," <i>PNAS USA</i> , 84(24):8773-8777 (1987); Lee, J.K. et al. "Nucleotide sequence of the gene encoding the Corynebacterium glutamicum mannose enzyme II and analyses of the deduced protein sequence," <i>FEMS Microbiol. Lett.</i> , 119(1-2):137-145 (1994)
L27123	aceB	Malate synthase	Lee, H.-S. et al. "Molecular characterization of aceB, a gene encoding malate synthase in Corynebacterium glutamicum," <i>J. Microbiol. Biotechnol.</i> , 4(4):256-263 (1994)
L27126		Pyruvate kinase	Jetten, M. S. et al. "Structural and functional analysis of pyruvate kinase from Corynebacterium glutamicum," <i>Appl. Environ. Microbiol.</i> , 60(7):2501-2507 (1994)
L28760	aceA	Isocitrate lyase	
L35906	dxr	Diphtheria toxin repressor	Oguiza, J.A. et al. "Molecular cloning, DNA sequence analysis, and characterization of the Corynebacterium diphtheriae dxr from Brevibacterium lactofermentum," <i>J. Bacteriol.</i> , 177(2):465-467 (1995)
M13774		Prephenate dehydratase	Follettie, M.T. et al. "Molecular cloning and nucleotide sequence of the Corynebacterium glutamicum pheA gene," <i>J. Bacteriol.</i> , 167:695-702 (1986)
M16175	5S rRNA		Park, Y.-H. et al. "Phylogenetic analysis of the coryneform bacteria by 5S rRNA sequences," <i>J. Bacteriol.</i> , 169:1801-1806 (1987)
M16663	trpE	Anthranyl synthase, 5' end	Sano, K. et al. "Structure and function of the trp operon control regions of Brevibacterium lactofermentum, a glutamic-acid-producing bacterium," <i>Gene</i> , 52:191-200 (1987)
M16664	trpA	Tryptophan synthase, 3' end	Sano, K. et al. "Structure and function of the trp operon control regions of Brevibacterium lactofermentum, a glutamic-acid-producing bacterium," <i>Gene</i> , 52:191-200 (1987)

Table 2 (continued)

		Phosphoenolpyruvate carboxylase	
M25819			O'Regan, M. et al. "Cloning and nucleotide sequence of the Phosphoenolpyruvate carboxylase-coding gene of <i>Corynebacterium glutamicum</i> ATCC13032," <i>Gene</i> , 77(2):237-251 (1989)
M85106		23S rRNA gene insertion sequence	Roller, C. et al. "Gram-positive bacteria with a high DNA G+C content are characterized by a common insertion within their 23S rRNA genes," <i>J. Gen. Microbiol.</i> , 138:1167-1175 (1992)
M85107, M85108		23S rRNA gene insertion sequence	Roller, C. et al. "Gram-positive bacteria with a high DNA G+C content are characterized by a common insertion within their 23S rRNA genes," <i>J. Gen. Microbiol.</i> , 138:1167-1175 (1992)
M89931	aecD; brnQ; yhbW	Beta C-S lyase; branched-chain amino acid uptake carrier; hypothetical protein yhbW	Rosol, I. et al. "The <i>Corynebacterium glutamicum</i> aecD gene encodes a C-S lyase with alpha, beta-elimination activity that degrades aminoethylcysteine," <i>J. Bacteriol.</i> , 174(9):2968-2977 (1992); Tauch, A. et al. "Isoleucine uptake in <i>Corynebacterium glutamicum</i> ATCC 13032 is directed by the brnQ gene product," <i>Arch. Microbiol.</i> , 169(4):303-312 (1998)
S59299	trp	Leader gene (promoter)	Herry, D.M. et al. "Cloning of the trp gene cluster from a tryptophan-hyperproducing strain of <i>Corynebacterium glutamicum</i> : identification of a mutation in the trp leader sequence," <i>Appl. Environ. Microbiol.</i> , 59(3):791-799 (1993)
U11545	trpD	Anthraniolate phosphoribosyltransferase	O'Gara, J.P. and Dunican, L.K. (1994) Complete nucleotide sequence of the <i>Corynebacterium glutamicum</i> ATCC 21850 trpD gene." Thesis, Microbiology Department, University College Galway, Ireland.
U13922	cgIIIM; cgIIR; cglIIR	Putative type II 5-cytosine methyltransferase; putative type II restriction endonuclease; type I or type III restriction endonuclease	Schafer, A. et al. "Cloning and characterization of a DNA region encoding a stress-sensitive restriction system from <i>Corynebacterium glutamicum</i> ATCC 13032 and analysis of its role in intergeneric conjugation with <i>Escherichia coli</i> ," <i>J. Bacteriol.</i> , 176(23):7309-7319 (1994); Schafer, A. et al. "The <i>Corynebacterium glutamicum</i> cglIM gene encoding a 5-cytosine in an McrBC-deficient <i>Escherichia coli</i> strain," <i>Gene</i> , 203(2):95-101 (1997)
U14965	recA		
U31224	ppx		Ankri, S. et al. "Mutations in the <i>Corynebacterium glutamicum</i> proline biosynthetic pathway: A natural bypass of the proA step," <i>J. Bacteriol.</i> , 178(15):4412-4419 (1996)
U31225	proC	L-proline: NADP+ 5-oxidoreductase	Ankri, S. et al. "Mutations in the <i>Corynebacterium glutamicum</i> proline biosynthetic pathway: A natural bypass of the proA step," <i>J. Bacteriol.</i> , 178(15):4412-4419 (1996)
U31230	obg; proB; unkdh	?; gamma glutamyl kinase; similar to D-isomer specific 2-hydroxyacid dehydrogenases	Ankri, S. et al. "Mutations in the <i>Corynebacterium glutamicum</i> proline biosynthetic pathway: A natural bypass of the proA step," <i>J. Bacteriol.</i> , 178(15):4412-4419 (1996)

Table 2 (continued)

U31281	bioB	Biotin synthase	Serebriskii, I.G., "Two new members of the bio B superfamily: Cloning, sequencing and expression of bio B genes of <i>Methylobacillus flagellatum</i> and <i>Corynebacterium glutamicum</i> ," <i>Gene</i> , 175:15-22 (1996)
U35023	thtR; accBC	Thiosulfate sulfurtransferase; acyl CoA carboxylase	Jager, W. et al. "A <i>Corynebacterium glutamicum</i> gene encoding a two-domain protein similar to biotin carboxylases and biotin-carboxyl-carrier proteins," <i>Arch. Microbiol.</i> , 166(2):76-82 (1996)
U43535	cmr	Multidrug resistance protein	Jager, W. et al. "A <i>Corynebacterium glutamicum</i> gene conferring multidrug resistance in the heterologous host <i>Escherichia coli</i> ," <i>J. Bacteriol.</i> , 179(7):2449-2451 (1997)
U43536	clpB	Heat shock ATP-binding protein	
U3587	aphA-3	3'-5'-aminoglycoside phosphotransferase	
U89648		<i>Corynebacterium glutamicum</i> unidentified sequence involved in histidine biosynthesis, partial sequence	
X04960	trpA; trpB; trpC; trpD; trpE; trpG; trpL	Tryptophan operon	Matsui, K. et al. "Complete nucleotide and deduced amino acid sequences of the <i>Brevibacterium lactofermentum</i> tryptophan operon," <i>Nucleic Acids Res.</i> , 14(24):10113-10114 (1986)
X07563	lys A	DAP decarboxylase (meso-diaminopimelate decarboxylase, EC 4.1.1.20)	Yeh, P. et al. "Nucleic sequence of the lysA gene of <i>Corynebacterium glutamicum</i> and possible mechanisms for modulation of its expression," <i>Mol. Gen. Genet.</i> , 212(1):112-119 (1988)
X14234	EC 4.1.1.31	Phosphoenolpyruvate carboxylase	Eikmanns, B.J. et al. "The Phosphoenolpyruvate carboxylase gene of <i>Corynebacterium glutamicum</i> : Molecular cloning, nucleotide sequence, and expression," <i>Mol. Gen. Genet.</i> , 218(2):330-339 (1989); Lepiniec, L. et al. "Sorghum Phosphoenolpyruvate carboxylase gene family: structure, function and molecular evolution," <i>Plant. Mol. Biol.</i> , 21 (3):487-502 (1993)
X17313	fda	Fructose-bisphosphate aldolase	Von der Osten, C.H. et al. "Molecular cloning, nucleotide sequence and fine-structural analysis of the <i>Corynebacterium glutamicum</i> fda gene: structural comparison of <i>C. glutamicum</i> fructose-1, 6-bisphosphate aldolase to class I and class II aldolases," <i>Mol. Microbiol.</i> ,
X53993	dapA	L-2, 3-dihydrodipicolinate synthetase (EC 4.2.1.52)	Bonnassie, S. et al. "Nucleic sequence of the dapA gene from <i>Corynebacterium glutamicum</i> ," <i>Nucleic Acids Res.</i> , 18(21):6421 (1990)
X54223		AttB-related site	Cianciotto, N. et al. "DNA sequence homology between att B-related sites of <i>Corynebacterium diphtheriae</i> , <i>Corynebacterium ulcerans</i> , <i>Corynebacterium glutamicum</i> , and the attP site of lambda dacycophage," <i>FEMS. Microbiol. Lett.</i> , 66:299-302 (1990)
X54740	argS; lysA	Arginyl-tRNA synthetase; Diaminopimelate decarboxylase	Marcel, T. et al. "Nucleotide sequence and organization of the upstream region of the <i>Corynebacterium glutamicum</i> lysA gene," <i>Mol. Microbiol.</i> , 4(11):1819-1830 (1990)

Table 2 (continued)

	trpL; trpE	Purative leader peptide; anthranilate synthase component 1	
X55994			Heery, D.M. et al. "Nucleotide sequence of the <i>Corynebacterium glutamicum</i> trpE gene," <i>Nucleic Acids Res.</i> , 18(23):7138 (1990)
X56037	thrC	Threonine synthase	Han, K.S. et al. "The molecular structure of the <i>Corynebacterium glutamicum</i> threonine synthase gene," <i>Mol. Microbiol.</i> , 4(10):1693-1702 (1990)
X56075	attB-related site	Attachment site	Cianciotto, N. et al. "DNA sequence homology between att B-related sites of <i>Corynebacterium diphtheriae</i> , <i>Corynebacterium ulcerans</i> , <i>Corynebacterium glutamicum</i> , and the attP site of <i>lambdacorynephage</i> ," <i>FEMS. Microbiol. Lett.</i> , 66:299-302 (1990)
X57226	lysC-alpha; lysC-beta; asd	Aspartokinase-alpha subunit; Aspartokinase-beta subunit; aspartate beta semialdehyde dehydrogenase	Kalinowski, J. et al. "Genetic and biochemical analysis of the Aspartokinase from <i>Corynebacterium glutamicum</i> ," <i>Mol. Microbiol.</i> , 5(5):1197-1204 (1991); Kalinowski, J. et al. "Aspartokinase genes lysC alpha and lysC beta overlap and are adjacent to the aspartate beta-semialdehyde dehydrogenase gene asd in <i>Corynebacterium glutamicum</i> ," <i>Mol. Gen. Genet.</i> , 224(3):317-324 (1990)
X59403	gap;pgk; tpi	Glyceraldehyde-3-phosphate; phosphoglycerate kinase; triosephosphate isomerase	Eikmanns, B.J. "Identification, sequence analysis, and expression of a <i>Corynebacterium glutamicum</i> gene cluster encoding the three glycolytic enzymes glyceraldehyde-3-phosphate dehydrogenase, 3-phosphoglycerate kinase, and triosephosphate isomerase," <i>J. Bacteriol.</i> , 174(19):6076-6086 (1992)
X59404	gdh	Glutamate dehydrogenase	Bormann, E.R. et al. "Molecular analysis of the <i>Corynebacterium glutamicum</i> gdh gene encoding glutamate dehydrogenase," <i>Mol. Microbiol.</i> , 6(3):317-326 (1992)
X60312	lysI	L-lysine permease	Seep-Feldhaus, A.H. et al. "Molecular analysis of the <i>Corynebacterium glutamicum</i> lysI gene involved in lysine uptake," <i>Mol. Microbiol.</i> , 5(12):2995-3005 (1991)
X66078	copI	Psl protein	Joliff, G. et al. "Cloning and nucleotide sequence of the copI gene encoding Psl, one of the two major secreted proteins of <i>Corynebacterium glutamicum</i> : The deduced N-terminal region of Psl is similar to the <i>Mycobacterium</i> antigen 85 complex," <i>Mol. Microbiol.</i> , 6(16):2349-2362 (1992)
X66112	glt	Citrate synthase	Eikmanns, B.J. et al. "Cloning sequence, expression and transcriptional analysis of the <i>Corynebacterium glutamicum</i> gltA gene encoding citrate synthase," <i>Microbiol.</i> , 140:1817-1828 (1994)
X67737	dapB	Dihydrodipicolinate reductase	
X69103	csp2	Surface layer protein PS2	Peyret, J.L. et al. "Characterization of the cspB gene encoding PS2, an ordered surface-layer protein in <i>Corynebacterium glutamicum</i> ," <i>Mol. Microbiol.</i> , 9(1):97-109 (1993)
X69104		IS3 related insertion element	Bonamy, C. et al. "Identification of IS1206, a <i>Corynebacterium glutamicum</i> IS3-related insertion sequence and phylogenetic analysis," <i>Mol. Microbiol.</i> , 14(3):571-581 (1994)

Table 2 (continued)

X70959	leuA	Isopropylmalate synthase	Patek, M. et al. "Leucine synthesis in <i>Corynebacterium glutamicum</i> : enzyme activities, structure of leuA, and effect of leuA inactivation on lysine synthesis," <i>Appl. Environ. Microbiol.</i> , 60(1):133-140 (1994)
X71489	icd	Isocitrate dehydrogenase (NADP+)	Eikmanns, B.J. et al. "Cloning sequence analysis, expression, and inactivation of the <i>Corynebacterium glutamicum</i> icd gene encoding isocitrate dehydrogenase and biochemical characterization of the enzyme," <i>J. Bacteriol.</i> , 177(3):774-782 (1995)
X72855	GDHA	Glutamate dehydrogenase (NADP+)	
X75083, X70584	mirA	5-methyltryptophan resistance	Heery, D.M. et al. "A sequence from a tryptophan-hyperproducing strain of <i>Corynebacterium glutamicum</i> encoding resistance to 5-methyltryptophan," <i>Biochem. Biophys. Res. Commun.</i> , 201(3):1255-1262 (1994)
X75085	recA		Fitzpatrick, R. et al. "Construction and characterization of recA mutant strains of <i>Corynebacterium glutamicum</i> and <i>Brevibacterium lactofermentum</i> ," <i>Appl. Microbiol. Biotechnol.</i> , 42(4):575-580 (1994)
X75504	aceA, thiX	Partial Isocitrate lyase; ?	Reinscheid, D.J. et al. "Characterization of the isocitrate lyase gene from <i>Corynebacterium glutamicum</i> and biochemical analysis of the enzyme," <i>J. Bacteriol.</i> , 176(12):3474-3483 (1994)
X76875		ATPase beta-subunit	Ludwig, W. et al. "Phylogenetic relationships of bacteria based on comparative sequence analysis of elongation factor Tu and ATP-synthase beta-subunit genes," <i>Antonie Van Leeuwenhoek</i> , 64:285-305 (1993)
X77034	tuf	Elongation factor Tu	Ludwig, W. et al. "Phylogenetic relationships of bacteria based on comparative sequence analysis of elongation factor Tu and ATP-synthase beta-subunit genes," <i>Antonie Van Leeuwenhoek</i> , 64:285-305 (1993)
X77384	recA		Billman-Jacobe, H. "Nucleotide sequence of a recA gene from <i>Corynebacterium glutamicum</i> ," <i>DNA Seq.</i> , 4(6):403-404 (1994)
X78491	aceB	Malate synthase	Reinscheid, D.J. et al. "Malate synthase from <i>Corynebacterium glutamicum</i> pta-ack operon encoding phosphotransacetylase: sequence analysis," <i>Microbiology</i> , 140:3099-3108 (1994)
X80629	16S rDNA	16S ribosomal RNA	Rainey, F.A. et al. "Phylogenetic analysis of the genera <i>Rhodococcus</i> and <i>Norcardia</i> and evidence for the evolutionary origin of the genus <i>Norcardia</i> from within the radiation of <i>Rhodococcus</i> species," <i>Microbiol.</i> , 141:523-528 (1995)
X81191	gluA; gluB; gluC; gluD	Glutamate uptake system	Kronmeyer, W. et al. "Structure of the gluABCD cluster encoding the glutamate uptake system of <i>Corynebacterium glutamicum</i> ," <i>J. Bacteriol.</i> , 177(5):1152-1158 (1995)
X81379	dapE	Succinylidiaminopimelate desuccinylase	Wehrmann, A. et al. "Analysis of different DNA fragments of <i>Corynebacterium glutamicum</i> complementing dapE of <i>Escherichia coli</i> ," <i>Microbiology</i> , 40:3349-56 (1994)

Table 2 (continued)

	16S rDNA	16S ribosomal RNA	
X82061			Ruimy, R. et al. "Phylogeny of the genus <i>Corynebacterium</i> deduced from analyses of small-subunit ribosomal DNA sequences," <i>Int. J. Syst. Bacteriol.</i> , 45(4):740-746 (1995)
X82928	asd; lysC	Aspartate-semialdehyde dehydrogenase; ?	Serebrijski, I. et al. "Multicopy suppression by asd gene and osmotic stress-dependent complementation by heterologous proA in proA mutants," <i>J. Bacteriol.</i> , 177(24):7255-7260 (1995)
X82929	proA	Gamma-glutamyl phosphate reductase	Serebrijski, I. et al. "Multicopy suppression by asd gene and osmotic stress-dependent complementation by heterologous proA in proA mutants," <i>J. Bacteriol.</i> , 177(24):7255-7260 (1995)
X84257	16S rDNA	16S ribosomal RNA	Pascual, C. et al. "Phylogenetic analysis of the genus <i>Corynebacterium</i> based on 16S rRNA gene sequences," <i>Int. J. Syst. Bacteriol.</i> , 45(4):724-728 (1995)
X85965	aroP; dapE	Aromatic amino acid permease; ?	Wehrmann et al. "Functional analysis of sequences adjacent to dapE of <i>C. glutamicum</i> proline reveals the presence of aroP, which encodes the aromatic amino acid transporter," <i>J. Bacteriol.</i> , 177(20):5991-5993 (1995)
X86157	argB; argC; argD; argF; argJ	Acetylglutamate kinase; N-acetyl-gamma-glutamyl-phosphate reductase; acetylornithine aminotransferase; ornithine carbamoyltransferase; glutamate N-acetyltransferase	Sakanyan, V. et al. "Genes and enzymes of the acetyl cycle of arginine biosynthesis in <i>Corynebacterium glutamicum</i> : enzyme evolution in the early steps of the arginine pathway," <i>Microbiology</i> , 142:99-108 (1996)
X89084	pta; ackA	Phosphate acetyltransferase; acetate kinase	Reinscheid, D.J. et al. "Cloning, sequence analysis, expression and inactivation of the <i>Corynebacterium glutamicum</i> pta-ack operon encoding phosphotransacetylase and acetate kinase," <i>Microbiology</i> , 145:503-513 (1999)
X89850	attB	Attachment site	Le Marrec, C. et al. "Genetic characterization of site-specific integration functions of phi AAU2 infecting 'Arthrobacter aureus C70,' <i>J. Bacteriol.</i> , 178(7):1996-2004 (1996)
X90356		Promoter fragment F1	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90357		Promoter fragment F2	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90358		Promoter fragment F10	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90359		Promoter fragment F13	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)

Table 2 (continued)

X90360		Promoter fragment F22	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90361		Promoter fragment F34	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90362		Promoter fragment F37	Patek, M. et al. "Promoters from <i>C. glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90363		Promoter fragment F45	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90364		Promoter fragment F64	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90365		Promoter fragment F75	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90366		Promoter fragment PF101	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90367		Promoter fragment PF104	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X90368		Promoter fragment PF109	Patek, M. et al. "Promoters from <i>Corynebacterium glutamicum</i> : cloning, molecular analysis and search for a consensus motif," <i>Microbiology</i> , 142:1297-1309 (1996)
X93513	amt	Ammonium transport system	Siewe, R.M. et al. "Functional and genetic characterization of the (methyl) ammonium uptake carrier of <i>Corynebacterium glutamicum</i> ," <i>J. Biol. Chem.</i> , 271(10):5398-5403 (1996)
X93514	betP	Glycine betaine transport system	Peter, H. et al. "Isolation, characterization, and expression of the <i>Corynebacterium glutamicum</i> betP gene, encoding the transport system for the compatible solute glycine betaine," <i>J. Bacteriol.</i> , 178(17):5229-5234 (1996)
X95649	orf4		Patek, M. et al. "Identification and transcriptional analysis of the dapB-ORF2-dapA-ORF4 operon of <i>Corynebacterium glutamicum</i> , encoding two enzymes involved in L-lysine synthesis," <i>Biotechnol. Lett.</i> , 19:1113-1117 (1997)
X96471	lysE; lysG	Lysine exporter protein; Lysine export regulator protein	Vrijic, M. et al. "A new type of transporter with a new type of cellular function: L-lysine export from <i>Corynebacterium glutamicum</i> ," <i>Mol. Microbiol.</i> , 22(5):815-826 (1996)



Table 2 (continued)

X96580	panB; panC; xylB	3-methyl-2-oxobutanoate hydroxymethyltransferase; pantoate-beta-alanine ligase; xylulokinase	Sahm, H. et al. "D-pantothenate synthesis in <i>Corynebacterium glutamicum</i> and use of panBC and genes encoding L-valine synthesis for D-pantothenate overproduction," <i>Appl. Environ. Microbiol.</i> , 65(5):1973-1979 (1999)
X96962		Insertion sequence IS1207 and transposase	
X99289		Elongation factor P	Ramos, A. et al. "Cloning, sequencing and expression of the gene encoding elongation factor P in the amino-acid producer <i>Brevibacterium lactofermentum</i> ( <i>Corynebacterium glutamicum</i> ATCC 13869)," <i>Gene</i> , 198:217-222 (1997)
Y00140	thrB	Homoserine kinase	Mateos, L.M. et al. "Nucleotide sequence of the homoserine kinase (thrB) gene of the <i>Brevibacterium lactofermentum</i> ," <i>Nucleic Acids Res.</i> , 15(9):3922 (1987)
Y00151	ddh	Meso-diaminopimelate D-dehydrogenase (EC 1.4.1.16)	Ishino, S. et al. "Nucleotide sequence of the meso-diaminopimelate D-dehydrogenase gene from <i>Corynebacterium glutamicum</i> ," <i>Nucleic Acids Res.</i> , 15(9):3917 (1987)
Y00476	thrA	Homoserine dehydrogenase	Mateos, L.M. et al. "Nucleotide sequence of the homoserine dehydrogenase (thrA) gene of the <i>Brevibacterium lactofermentum</i> ," <i>Nucleic Acids Res.</i> , 15(24):10598 (1987)
Y00546	hom; thrB	Homoserine dehydrogenase; homoserine kinase	Peoples, O.P. et al. "Nucleotide sequence and fine structural analysis of the <i>Corynebacterium glutamicum</i> hom-thrB operon," <i>Mol. Microbiol.</i> , 2(1):63-72 (1988)
Y08964	murC; ftsQ/divD; ftsZ	UPD-N-acetylmuramate-alanine ligase; division initiation protein or cell division protein; cell division protein	Honrubia, M.P. et al. "Identification, characterization, and chromosomal organization of the ftsZ gene from <i>Brevibacterium lactofermentum</i> ," <i>Mol. Gen. Genet.</i> , 259(1):97-104 (1998)
Y09163	putP	High affinity proline transport system	Peter, H. et al. "Isolation of the putP gene of <i>Corynebacterium glutamicum</i> proline and characterization of a low-affinity uptake system for compatible solutes," <i>Arch. Microbiol.</i> , 168(2):143-151 (1997)
Y09548	pyc	Pyruvate carboxylase	Peters-Wendisch, P.G. et al. "Pyruvate carboxylase from <i>Corynebacterium glutamicum</i> : characterization, expression and inactivation of the pyc gene," <i>Microbiology</i> , 144:915-927 (1998)
Y09578	leuB	3-isopropylmalate dehydrogenase	Patek, M. et al. "Analysis of the leuB gene from <i>Corynebacterium glutamicum</i> ," <i>Appl. Microbiol. Biotechnol.</i> , 50(1):42-47 (1998)
Y12472		Attachment site bacteriophage Phi-16	Moreau, S. et al. "Site-specific integration of corynebacteriophage Phi-16: The construction of an integration vector," <i>Microbiol.</i> , 145:539-548 (1999)
Y12537	proP	Proline/ectoine uptake system protein	Peter, H. et al. "Corynebacterium glutamicum is equipped with four secondary carriers for compatible solutes: Identification, sequencing, and characterization of the proline/ectoine uptake system, ProP, and the ectoine/proline/glycine betaine carrier, EctP," <i>J. Bacteriol.</i> , 180(22):6005-6012 (1998)

Table 2 (continued)

Y13221	glnA	Glutamine synthetase I	Jakoby, M. et al. "Isolation of <i>Corynebacterium glutamicum</i> glnA gene encoding glutamine synthetase I," <i>FEBS Microbiol. Lett.</i> , 154(1):81-88 (1997)
Y16642	lpd	Dihydrolipoamide dehydrogenase	
Y18059		Attachment site <i>Corynebacterium</i> 304L	Moreau, S. et al. "Analysis of the integration functions of $\phi$ 304L: An integrase module among corynephages," <i>Virology</i> , 255(1):150-159 (1999)
Z21501	argS; lysA	Arginyl-tRNA synthetase; diaminopimelate decarboxylase (partial)	Oguiza, J.A. et al. "A gene encoding arginyl-tRNA synthetase is located in the upstream region of the lysA gene in <i>Brevibacterium lactofermentum</i> : Regulation of argS-lysA cluster expression by arginine," <i>J. Bacteriol.</i> , 175(22):7356-7362 (1993)
Z21502	dapA; dapB	Dihydrodipicolinate synthase; dihydrodipicolinate reductase	Pisabarro, A. et al. "A cluster of three genes (dapA, orf2, and dapB) of <i>Brevibacterium lactofermentum</i> encodes dihydrodipicolinate reductase, and a third polypeptide of unknown function," <i>J. Bacteriol.</i> , 175(9):2743-2749 (1993)
Z29563	thrC	Threonine synthase	Malumbres, M. et al. "Analysis and expression of the thrC gene of the encoded threonine synthase," <i>Appl. Environ. Microbiol.</i> , 60(7):2209-2219 (1994)
Z46753	16S rDNA	Gene for 16S ribosomal RNA	
Z49822	sigA	SigA sigma factor	Oguiza, J.A. et al. "Multiple sigma factor genes in <i>Brevibacterium lactofermentum</i> : Characterization of sigA and sigB," <i>J. Bacteriol.</i> , 178(2):550-553 (1996)
Z49823	galE; dtxR	Catalytic activity UDP-galactose 4-epimerase; diphtheria toxin regulatory protein	Oguiza, J.A. et al. "The galE gene encoding the UDP-galactose 4-epimerase of <i>Brevibacterium lactofermentum</i> is coupled transcriptionally to the dmdR gene," <i>Gene</i> , 177:103-107 (1996)
Z49824	orf1; sigB	?, SigB sigma factor	Oguiza, J.A. et al. "Multiple sigma factor genes in <i>Brevibacterium lactofermentum</i> : Characterization of sigA and sigB," <i>J. Bacteriol.</i> , 178(2):550-553 (1996)
Z66534		Transposase	Correia, A. et al. "Cloning and characterization of an IS-like element present in the genome of <i>Brevibacterium lactofermentum</i> ATCC 13869," <i>Gene</i> , 170(1):91-94 (1996)

A sequence for this gene was published in the indicated reference. However, the sequence obtained by the inventors of the present application is significantly longer than the published version. It is believed that the published version relied on an incorrect start codon, and thus represents only a fragment of the actual coding region.

TABLE 3: *Corynebacterium* and *Brevibacterium* Strains Which May be Used in the Practice of the Invention

Genus	species	ATCC	FERM	NRRL	CECT	NCIMB	CBS	NCTG	DSMZ
<i>Brevibacterium</i>	<i>ammoniagenes</i>	21054							
<i>Brevibacterium</i>	<i>ammoniagenes</i>	19350							
<i>Brevibacterium</i>	<i>ammoniagenes</i>	19351							
<i>Brevibacterium</i>	<i>ammoniagenes</i>	19352							
<i>Brevibacterium</i>	<i>ammoniagenes</i>	19353							
<i>Brevibacterium</i>	<i>ammoniagenes</i>	19354							
<i>Brevibacterium</i>	<i>ammoniagenes</i>	19355							
<i>Brevibacterium</i>	<i>ammoniagenes</i>	19356							
<i>Brevibacterium</i>	<i>ammoniagenes</i>	21055							
<i>Brevibacterium</i>	<i>ammoniagenes</i>	21077							
<i>Brevibacterium</i>	<i>ammoniagenes</i>	21553							
<i>Brevibacterium</i>	<i>ammoniagenes</i>	21580							
<i>Brevibacterium</i>	<i>ammoniagenes</i>	39101							
<i>Brevibacterium</i>	<i>butanicum</i>	21196							
<i>Brevibacterium</i>	<i>divaricatum</i>	21792	P928						
<i>Brevibacterium</i>	<i>flavum</i>	21474							
<i>Brevibacterium</i>	<i>flavum</i>	21129							
<i>Brevibacterium</i>	<i>flavum</i>	21518							
<i>Brevibacterium</i>	<i>flavum</i>			B11474					
<i>Brevibacterium</i>	<i>flavum</i>			B11472					
<i>Brevibacterium</i>	<i>flavum</i>	21127							
<i>Brevibacterium</i>	<i>flavum</i>	21128							
<i>Brevibacterium</i>	<i>flavum</i>	21427							
<i>Brevibacterium</i>	<i>flavum</i>	21475							
<i>Brevibacterium</i>	<i>flavum</i>	21517							
<i>Brevibacterium</i>	<i>flavum</i>	21528							
<i>Brevibacterium</i>	<i>flavum</i>	21529							
<i>Brevibacterium</i>	<i>flavum</i>			B11477					
<i>Brevibacterium</i>	<i>flavum</i>			B11478					
<i>Brevibacterium</i>	<i>flavum</i>	21127							
<i>Brevibacterium</i>	<i>flavum</i>			B11474					
<i>Brevibacterium</i>	<i>healii</i>	15527							
<i>Brevibacterium</i>	<i>ketoglutamicum</i>	21004							
<i>Brevibacterium</i>	<i>ketoglutamicum</i>	21089							
<i>Brevibacterium</i>	<i>ketosoreductum</i>	21914							
<i>Brevibacterium</i>	<i>lactofermentum</i>				70				
<i>Brevibacterium</i>	<i>lactofermentum</i>				74				
<i>Brevibacterium</i>	<i>lactofermentum</i>				77				
<i>Brevibacterium</i>	<i>lactofermentum</i>	21798							
<i>Brevibacterium</i>	<i>lactofermentum</i>	21799							
<i>Brevibacterium</i>	<i>lactofermentum</i>	21800							
<i>Brevibacterium</i>	<i>lactofermentum</i>	21801							
<i>Brevibacterium</i>	<i>lactofermentum</i>			B11470					
<i>Brevibacterium</i>	<i>lactofermentum</i>			B11471					

Brevibacterium	lactofermentum	21086							
Brevibacterium	lactofermentum	21420							
Brevibacterium	lactofermentum	21086							
Brevibacterium	lactofermentum	31269							
Brevibacterium	linens	9174							
Brevibacterium	linens	19391							
Brevibacterium	linens	8377							
Brevibacterium	paraffinolyticum					11160			
Brevibacterium	spec.						717.73		
Brevibacterium	spec.						717.73		
Brevibacterium	spec.	14604							
Brevibacterium	spec.	21860							
Brevibacterium	spec.	21864							
Brevibacterium	spec.	21865							
Brevibacterium	spec.	21866							
Brevibacterium	spec.	19240							
Corynebacterium	acetoacidophilum	21476							
Corynebacterium	acetoacidophilum	13870							
Corynebacterium	acetoglutamicum				B11473				
Corynebacterium	acetoglutamicum				B11475				
Corynebacterium	acetoglutamicum	15806							
Corynebacterium	acetoglutamicum	21491							
Corynebacterium	acetoglutamicum	31270							
Corynebacterium	acetophilum				B3671				
Corynebacterium	ammoniagenes	6872						2399	
Corynebacterium	ammoniagenes	15511							
Corynebacterium	fujikense	21496							
Corynebacterium	glutamicum	14067							
Corynebacterium	glutamicum	39137							
Corynebacterium	glutamicum	21254							
Corynebacterium	glutamicum	21255							
Corynebacterium	glutamicum	31830							
Corynebacterium	glutamicum	13032							
Corynebacterium	glutamicum	14305							
Corynebacterium	glutamicum	15455							
Corynebacterium	glutamicum	13058							
Corynebacterium	glutamicum	13059							
Corynebacterium	glutamicum	13060							
Corynebacterium	glutamicum	21492							
Corynebacterium	glutamicum	21513							
Corynebacterium	glutamicum	21526							
Corynebacterium	glutamicum	21543							
Corynebacterium	glutamicum	13287							
Corynebacterium	glutamicum	21851							
Corynebacterium	glutamicum	21253							
Corynebacterium	glutamicum	21514							
Corynebacterium	glutamicum	21516							
Corynebacterium	glutamicum	21299							
Corynebacterium	glutamicum	21300							

Corynebacterium	glutamicum	39684							
Corynebacterium	glutamicum	21488							
Corynebacterium	glutamicum	21649							
Corynebacterium	glutamicum	21650							
Corynebacterium	glutamicum	19223							
Corynebacterium	glutamicum	13869							
Corynebacterium	glutamicum	21157							
Corynebacterium	glutamicum	21158							
Corynebacterium	glutamicum	21159							
Corynebacterium	glutamicum	21355							
Corynebacterium	glutamicum	31808							
Corynebacterium	glutamicum	21674							
Corynebacterium	glutamicum	21562							
Corynebacterium	glutamicum	21563							
Corynebacterium	glutamicum	21564							
Corynebacterium	glutamicum	21565							
Corynebacterium	glutamicum	21566							
Corynebacterium	glutamicum	21567							
Corynebacterium	glutamicum	21568							
Corynebacterium	glutamicum	21569							
Corynebacterium	glutamicum	21570							
Corynebacterium	glutamicum	21571							
Corynebacterium	glutamicum	21572							
Corynebacterium	glutamicum	21573							
Corynebacterium	glutamicum	21579							
Corynebacterium	glutamicum	19049							
Corynebacterium	glutamicum	19050							
Corynebacterium	glutamicum	19051							
Corynebacterium	glutamicum	19052							
Corynebacterium	glutamicum	19053							
Corynebacterium	glutamicum	19054							
Corynebacterium	glutamicum	19055							
Corynebacterium	glutamicum	19056							
Corynebacterium	glutamicum	19057							
Corynebacterium	glutamicum	19058							
Corynebacterium	glutamicum	19059							
Corynebacterium	glutamicum	19060							
Corynebacterium	glutamicum	19185							
Corynebacterium	glutamicum	13286							
Corynebacterium	glutamicum	21515							
Corynebacterium	glutamicum	21527							
Corynebacterium	glutamicum	21544							
Corynebacterium	glutamicum	21492							
Corynebacterium	glutamicum				B8183				
Corynebacterium	glutamicum				B8182				
Corynebacterium	glutamicum				B12416				
Corynebacterium	glutamicum				B12417				
Corynebacterium	glutamicum				B12418				
Corynebacterium	glutamicum				B11476				

Corynebacterium	glutamicum	21608							
Corynebacterium	lilium		P973						
Corynebacterium	nitrilophilus	21419				11594			
Corynebacterium	spec.		P4445						
Corynebacterium	spec.		P4446						
Corynebacterium	spec.	31088							
Corynebacterium	spec.	31089							
Corynebacterium	spec.	31090							
Corynebacterium	spec.	31090							
Corynebacterium	spec.	31090							
Corynebacterium	spec.	15954							20145
Corynebacterium	spec.	21857							
Corynebacterium	spec.	21862							
Corynebacterium	spec.	21863							

ATCC: American Type Culture Collection, Rockville, MD, USA

FERM: Fermentation Research Institute, Chiba, Japan

NRRL: ARS Culture Collection, Northern Regional Research Laboratory, Peoria, IL, USA

CECT: Coleccion Espanola de Cultivos Tipo, Valencia, Spain

NCIMB: National Collection of Industrial and Marine Bacteria Ltd., Aberdeen, UK

CBS: Centraalbureau voor Schimmelcultures, Baarn, NL

NCTC: National Collection of Type Cultures, London, UK

DSMZ: Deutsche Sammlung von Mikroorganismen und Zellkulturen, Braunschweig, Germany

For reference see Sugawara, H. et al. (1993) World directory of collections of cultures of microorganisms: Bacteria, fungi and yeasts (4<sup>th</sup> edn), World federation for culture collections world data center on microorganisms, Saimata, Japan.

TABLE 4: ALIGNMENT RESULTS

ID #	length (NT)	Genbank Hit	Length	Accession	Name of Genbank Hit	Source of Genbank Hit	% homology (GAP)	Date of Deposit
rx00315	1527	GB_BA1:AB007125	4078	AB007125	Serratia marcescens slaA gene for surface layer protein, complete cds, isolate 8000.	Serratia marcescens	40,386	26-MAR-1998
		GB_IN1:CELC47D2	17381	U64861	Caenorhabditis elegans cosmid C47D2.	Caenorhabditis elegans	36,207	28-Jul-96
		GB_HTG2:AC006732	159453	AC006732	Caenorhabditis elegans clone Y32G9, *** SEQUENCING IN PROGRESS ***, 9 unordered pieces.	Caenorhabditis elegans	36,436	23-Feb-99
rx001503	372	GB_PR3:AC005019	188362	AC005019	Homo sapiens BAC clone GS250A16 from 7p21-p22, complete	Homo sapiens	39,722	27-Aug-98
		GB_GSS12:AQ390040	680	AQ390040	RPC111-157C9 TJ RPC1-11 Homo sapiens genomic clone RPC1-11-157C9 genomic survey sequence.	Homo sapiens	43,137	21-MAY-1999
		GB_GSS5:AQ784231	542	AQ784231	HS_3087_B1_C10_T7C CIT Approved Human Genomic Sperm Library D Homo sapiens genomic clone Plate=3087 Col=19 Row=F, genomic survey sequence.	Homo sapiens	37,643	3-Aug-99
rx001299	2187	GB_EST38:AW047296	614	AW047296	UI-M-BH1-amh-e-03-0-UI.s1 NIH_BMAP_M_S2 Mus musculus cDNA clone UI-M-BH1-amh-e-03-0-UI 3' mRNA sequence.	Mus musculus	41,475	18-Sep-99
		GB_RO:AB004056	1581	AB004056	Rattus norvegicus mRNA for BarH-class homeodomain transcription factor, complete cds.	Rattus norvegicus	41,031	2-Sep-98
		GB_RO:AB004056	1581	AB004056	Rattus norvegicus mRNA for BarH-class homeodomain transcription factor, complete cds.	Rattus norvegicus	40,717	2-Sep-98
rx000951	416	GB_BA1:SCJ21	31717	AL109747	Streptomyces coelicolor cosmid J21.	Streptomyces coelicolor A3(2)	34,913	5-Aug-99
		GB_VI:MCU68299	230278	U68299	Mouse cytomegalovirus 1 complete genomic sequence.	Mouse cytomegalovirus 1	40,097	04-DEC-1996
		GB_VI:U93872	133661	U93872	Kaposi's sarcoma-associated herpesvirus glycoprotein M, DNA replication protein, glycoprotein, DNA replication protein, FLICE inhibitory protein and v-cyclin genes, complete cds, and tegument	Kaposi's sarcoma-associated herpesvirus	36,029	9-Jul-97
rx001244	1827	GB_BA1:AFAPHBHI	4501	M69036	Alcaligenes eutrophus protein H (phbH) and protein I (phbI) genes, complete cds.	Ralstonia eutropha	45,624	26-Apr-93
		GB_PR3:HSJ836E13	78055	AL050326	Human DNA sequence from clone 836E13 on chromosome 20 Contains ESTs, STS and GSSs, complete sequence.	Homo sapiens	37,303	23-Nov-99
		GB_EST24:AI170227	409	AI170227	EST216152 Normalized rat lung, Bento Soares Rattus sp. cDNA clone RLUCF56 3' end, mRNA sequence.	Rattus sp.	39,098	20-Jan-99
rx001300	390	GB_PR3:HUMDDDDA	26764	L39874	Homo sapiens deoxycytidylate deaminase gene, complete cds.	Homo sapiens	37,644	11-Aug-95
		GB_PAT:I40899	26764	I40899	Sequence 1 from patent US 5622851.	Unknown.	37,644	13-MAY-1997
		GB_PAT:I40900	1317	I40900	Sequence 2 from patent US 5622851.	Unknown.	37,644	13-MAY-1997
rx000953	789	GB_BA1:SCJ21	31717	AL109747	Streptomyces coelicolor cosmid J21.	Streptomyces coelicolor A3(2)	39,398	5-Aug-99
		GB_BA1:BLTRP	7725	X04960	Brevibacterium lactofermentum tryptophan operon.	Corynebacterium glutamicum	39,610	10-Feb-99
		GB_PAT:E01375	7726	E01375	DNA sequence of tryptophan operon.	Corynebacterium glutamicum	46,753	29-Sep-97

- 75 -

**Table 4 (continued)**

rx01943	2172	GB_BA1:CORPTSMA	2656	L18874	Corynebacterium glutamicum phosphoenolpyruvate sugar phosphotransferase (ptsM) mRNA, complete cds.	Corynebacterium glutamicum	100,000	24-Nov-94
		GB_BA1:BRLPTSG	3163	L18875	Brevibacterium lactofermentum phosphoenolpyruvate sugar phosphotransferase (ptsG) gene, complete cds.	Brevibacterium lactofermentum	84,963	01-OCT-1993
		GB_BA2:AF045481	2841	AF045481	Corynebacterium ammoniagenes glucose permease (ptsG) gene, complete cds.	Corynebacterium ammoniagenes	53,558	29-Jul-98



### Exemplification

#### **Example 1: Preparation of total genomic DNA of *Corynebacterium glutamicum* ATCC 13032**

- 5 A culture of *Corynebacterium glutamicum* (ATCC 13032) was grown overnight at 30°C with vigorous shaking in BHI medium (Difco). The cells were harvested by centrifugation, the supernatant was discarded and the cells were resuspended in 5 ml buffer-I (5% of the original volume of the culture — all indicated volumes have been calculated for 100 ml of culture volume). Composition of buffer-I: 140.34 g/l sucrose,
- 10 2.46 g/l  $\text{MgSO}_4 \times 7\text{H}_2\text{O}$ , 10 ml/l  $\text{KH}_2\text{PO}_4$  solution (100 g/l, adjusted to pH 6.7 with KOH), 50 ml/l M12 concentrate (10 g/l  $(\text{NH}_4)_2\text{SO}_4$ , 1 g/l NaCl, 2 g/l  $\text{MgSO}_4 \times 7\text{H}_2\text{O}$ , 0.2 g/l  $\text{CaCl}_2$ , 0.5 g/l yeast extract (Difco), 10 ml/l trace-elements-mix (200 mg/l  $\text{FeSO}_4 \times \text{H}_2\text{O}$ , 10 mg/l  $\text{ZnSO}_4 \times 7\text{H}_2\text{O}$ , 3 mg/l  $\text{MnCl}_2 \times 4\text{H}_2\text{O}$ , 30 mg/l  $\text{H}_3\text{BO}_3$ , 20 mg/l  $\text{CoCl}_2 \times 6\text{H}_2\text{O}$ , 1 mg/l  $\text{NiCl}_2 \times 6\text{H}_2\text{O}$ , 3 mg/l  $\text{Na}_2\text{MoO}_4 \times 2\text{H}_2\text{O}$ , 500 mg/l complexing agent
- 15 (EDTA or citric acid), 100 ml/l vitamins-mix (0.2 mg/l biotin, 0.2 mg/l folic acid, 20 mg/l p-amino benzoic acid, 20 mg/l riboflavin, 40 mg/l ca-panthothenate, 140 mg/l nicotinic acid, 40 mg/l pyridoxole hydrochloride, 200 mg/l myo-inositol). Lysozyme was added to the suspension to a final concentration of 2.5 mg/ml. After an approximately 4 h incubation at 37°C, the cell wall was degraded and the resulting
- 20 protoplasts are harvested by centrifugation. The pellet was washed once with 5 ml buffer-I and once with 5 ml TE-buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8). The pellet was resuspended in 4 ml TE-buffer and 0.5 ml SDS solution (10%) and 0.5 ml NaCl solution (5 M) are added. After adding of proteinase K to a final concentration of 200 µg/ml, the suspension is incubated for ca. 18 h at 37°C. The DNA was purified by
- 25 extraction with phenol, phenol-chloroform-isoamylalcohol and chloroform-isoamylalcohol using standard procedures. Then, the DNA was precipitated by adding 1/50 volume of 3 M sodium acetate and 2 volumes of ethanol, followed by a 30 min incubation at -20°C and a 30 min centrifugation at 12,000 rpm in a high speed centrifuge using a SS34 rotor (Sorvall). The DNA was dissolved in 1 ml TE-buffer containing 20
- 30 µg/ml RNaseA and dialysed at 4°C against 1000 ml TE-buffer for at least 3 hours. During this time, the buffer was exchanged 3 times. To aliquots of 0.4 ml of the dialysed DNA solution, 0.4 ml of 2 M LiCl and 0.8 ml of ethanol are added. After a 30

- 77 -

min incubation at -20°C, the DNA was collected by centrifugation (13,000 rpm, Biofuge Fresco, Heraeus, Hanau, Germany). The DNA pellet was dissolved in TE-buffer. DNA prepared by this procedure could be used for all purposes, including southern blotting or construction of genomic libraries.

5

**Example 2: Construction of genomic libraries in *Escherichia coli* of *Corynebacterium glutamicum* ATCC13032.**

Using DNA prepared as described in Example 1, cosmid and plasmid libraries were constructed according to known and well established methods (see e.g., Sambrook, J. *et al.* 10 (1989) "Molecular Cloning : A Laboratory Manual", Cold Spring Harbor Laboratory Press, or Ausubel, F.M. *et al.* (1994) "Current Protocols in Molecular Biology", John Wiley & Sons.)

Any plasmid or cosmid could be used. Of particular use were the plasmids pBR322 (Sutcliffe, J.G. (1979) *Proc. Natl. Acad. Sci. USA*, 75:3737-3741); pACYC177 (Change & 15 Cohen (1978) *J. Bacteriol* 134:1141-1156), plasmids of the pBS series (pBSSK+, pBSSK- and others; Stratagene, LaJolla, USA), or cosmids as SuperCos1 (Stratagene, LaJolla, USA) or Lorist6 (Gibson, T.J., Rosenthal A. and Waterson, R.H. (1987) *Gene* 53:283-286. Gene libraries specifically for use in *C. glutamicum* may be constructed using plasmid pSL109 (Lee, H.-S. and A. J. Sinskey (1994) *J. Microbiol. Biotechnol.* 4: 256-263).

20

**Example 3: DNA Sequencing and Computational Functional Analysis**

Genomic libraries as described in Example 2 were used for DNA sequencing according to standard methods, in particular by the chain termination method using ABI377 sequencing machines (see e.g., Fleischman, R.D. *et al.* (1995) "Whole-genome 25 Random Sequencing and Assembly of Haemophilus Influenzae Rd., *Science*, 269:496-512). Sequencing primers with the following nucleotide sequences were used: 5'-GGAAACAGTATGACCATG-3' or 5'-GTAAAACGACGGCCAGT-3'.

**Example 4: *In vivo* Mutagenesis**

30 *In vivo* mutagenesis of *Corynebacterium glutamicum* can be performed by passage of plasmid (or other vector) DNA through *E. coli* or other microorganisms (e.g. *Bacillus* spp. or yeasts such as *Saccharomyces cerevisiae*) which are impaired in their capabilities to maintain

the integrity of their genetic information. Typical mutator strains have mutations in the genes for the DNA repair system (*e.g.*, mutHLS, mutD, mutT, etc.; for reference, see Rupp, W.D. (1996) DNA repair mechanisms, in: *Escherichia coli* and *Salmonella*, p. 2277-2294, ASM: Washington.) Such strains are well known to one of ordinary skill in the art. The use of such  
5 strains is illustrated, for example, in Greener, A. and Callahan, M. (1994) *Strategies* 7: 32-34.

**Example 5: DNA Transfer Between *Escherichia coli* and *Corynebacterium glutamicum***

Several *Corynebacterium* and *Brevibacterium* species contain endogenous  
10 plasmids (as *e.g.*, pHM1519 or pBL1) which replicate autonomously (for review see, *e.g.*, Martin, J.F. *et al.* (1987) *Biotechnology*, 5:137-146). Shuttle vectors for *Escherichia coli* and *Corynebacterium glutamicum* can be readily constructed by using standard vectors for *E. coli* (Sambrook, J. *et al.* (1989), "Molecular Cloning: A Laboratory Manual", Cold Spring Harbor Laboratory Press or Ausubel, F.M. *et al.* (1994) "Current Protocols in  
15 Molecular Biology", John Wiley & Sons) to which a origin or replication for and a suitable marker from *Corynebacterium glutamicum* is added. Such origins of replication are preferably taken from endogenous plasmids isolated from *Corynebacterium* and *Brevibacterium* species. Of particular use as transformation markers for these species are genes for kanamycin resistance (such as those derived from the Tn5 or Tn903  
20 transposons) or chloramphenicol (Winnacker, E.L. (1987) "From Genes to Clones — Introduction to Gene Technology, VCH, Weinheim). There are numerous examples in the literature of the construction of a wide variety of shuttle vectors which replicate in both *E. coli* and *C. glutamicum*, and which can be used for several purposes, including gene over-expression (for reference, see *e.g.*, Yoshihama, M. *et al.* (1985) *J. Bacteriol.* 162:591-597,  
25 Martin J.F. *et al.* (1987) *Biotechnology*, 5:137-146 and Eikmanns, B.J. *et al.* (1991) *Gene*, 102:93-98).

Using standard methods, it is possible to clone a gene of interest into one of the shuttle vectors described above and to introduce such a hybrid vectors into strains of *Corynebacterium glutamicum*. Transformation of *C. glutamicum* can be achieved by  
30 protoplast transformation (Kastsumata, R. *et al.* (1984) *J. Bacteriol.* 159:306-311), electroporation (Liebl, E. *et al.* (1989) *FEMS Microbiol. Letters*, 53:399-303) and in cases where special vectors are used, also by conjugation (as described *e.g.* in Schäfer, A *et al.*

(1990) *J. Bacteriol.* 172:1663-1666). It is also possible to transfer the shuttle vectors for *C. glutamicum* to *E. coli* by preparing plasmid DNA from *C. glutamicum* (using standard methods well-known in the art) and transforming it into *E. coli*. This transformation step can be performed using standard methods, but it is advantageous to use an *Mcr*-deficient *E. coli* strain, such as NM522 (Gough & Murray (1983) *J. Mol. Biol.* 166:1-19).

Genes may be overexpressed in *C. glutamicum* strains using plasmids which comprise pCG1 (U.S. Patent No. 4,617,267) or fragments thereof, and optionally the gene for kanamycin resistance from TN903 (Grindley, N.D. and Joyce, C.M. (1980) *Proc. Natl. Acad. Sci. USA* 77(12): 7176-7180). In addition, genes may be overexpressed in *C. glutamicum* strains using plasmid pSL109 (Lee, H.-S. and A. J. Sinskey (1994) *J. Microbiol. Biotechnol.* 4: 256-263).

Aside from the use of replicative plasmids, gene overexpression can also be achieved by integration into the genome. Genomic integration in *C. glutamicum* or other *Corynebacterium* or *Brevibacterium* species may be accomplished by well-known methods, such as homologous recombination with genomic region(s), restriction endonuclease mediated integration (REMI) (see, e.g., DE Patent 19823834), or through the use of transposons. It is also possible to modulate the activity of a gene of interest by modifying the regulatory regions (e.g., a promoter, a repressor, and/or an enhancer) by sequence modification, insertion, or deletion using site-directed methods (such as homologous recombination) or methods based on random events (such as transposon mutagenesis or REMI). Nucleic acid sequences which function as transcriptional terminators may also be inserted 3' to the coding region of one or more genes of the invention; such terminators are well-known in the art and are described, for example, in Winnacker, E.L. (1987) *From Genes to Clones – Introduction to Gene Technology*. VCH: Weinheim.

#### **Example 6: Assessment of the Expression of the Mutant Protein**

Observations of the activity of a mutated protein in a transformed host cell rely on the fact that the mutant protein is expressed in a similar fashion and in a similar quantity to that of the wild-type protein. A useful method to ascertain the level of transcription of the mutant gene (an indicator of the amount of mRNA available for translation to the gene product) is to perform a Northern blot (for reference see, for example, Ausubel *et al.*

(1988) Current Protocols in Molecular Biology, Wiley: New York), in which a primer designed to bind to the gene of interest is labeled with a detectable tag (usually radioactive or chemiluminescent), such that when the total RNA of a culture of the organism is extracted, run on gel, transferred to a stable matrix and incubated with this probe, the binding and quantity of binding of the probe indicates the presence and also the quantity of mRNA for this gene. This information is evidence of the degree of transcription of the mutant gene. Total cellular RNA can be prepared from *Corynebacterium glutamicum* by several methods, all well-known in the art, such as that described in Bormann, E.R. *et al.* (1992) *Mol. Microbiol.* 6: 317-326.

To assess the presence or relative quantity of protein translated from this mRNA, standard techniques, such as a Western blot, may be employed (see, for example, Ausubel *et al.* (1988) Current Protocols in Molecular Biology, Wiley: New York). In this process, total cellular proteins are extracted, separated by gel electrophoresis, transferred to a matrix such as nitrocellulose, and incubated with a probe, such as an antibody, which specifically binds to the desired protein. This probe is generally tagged with a chemiluminescent or colorimetric label which may be readily detected. The presence and quantity of label observed indicates the presence and quantity of the desired mutant protein present in the cell.

#### **Example 7: Growth of Genetically Modified *Corynebacterium glutamicum* — Media and Culture Conditions**

Genetically modified *Corynebacteria* are cultured in synthetic or natural growth media. A number of different growth media for *Corynebacteria* are both well-known and readily available (Lieb *et al.* (1989) *Appl. Microbiol. Biotechnol.*, 32:205-210; von der Osten *et al.* (1998) *Biotechnology Letters*, 11:11-16; Patent DE 4,120,867; Liebl (1992) "The Genus *Corynebacterium*, in: The Prokaryotes, Volume II, Balows, A. *et al.*, eds. Springer-Verlag). These media consist of one or more carbon sources, nitrogen sources, inorganic salts, vitamins and trace elements. Preferred carbon sources are sugars, such as mono-, di-, or polysaccharides. For example, glucose, fructose, mannose, galactose, ribose, sorbose, ribulose, lactose, maltose, sucrose, raffinose, starch or cellulose serve as very good carbon sources. It is also possible to supply sugar to the media via complex compounds such as molasses or other by-products from sugar refinement. It can also be

advantageous to supply mixtures of different carbon sources. Other possible carbon sources are alcohols and organic acids, such as methanol, ethanol, acetic acid or lactic acid. Nitrogen sources are usually organic or inorganic nitrogen compounds, or materials which contain these compounds. Exemplary nitrogen sources include ammonia gas or  
5 ammonia salts, such as  $\text{NH}_4\text{Cl}$  or  $(\text{NH}_4)_2\text{SO}_4$ ,  $\text{NH}_4\text{OH}$ , nitrates, urea, amino acids or complex nitrogen sources like corn steep liquor, soy bean flour, soy bean protein, yeast extract, meat extract and others.

Inorganic salt compounds which may be included in the media include the chloride-, phosphorous- or sulfate- salts of calcium, magnesium, sodium, cobalt,  
10 molybdenum, potassium, manganese, zinc, copper and iron. Chelating compounds can be added to the medium to keep the metal ions in solution. Particularly useful chelating compounds include dihydroxyphenols, like catechol or protocatechuate, or organic acids, such as citric acid. It is typical for the media to also contain other growth factors, such as vitamins or growth promoters, examples of which include biotin, riboflavin, thiamin, folic  
15 acid, nicotinic acid, pantothenate and pyridoxin. Growth factors and salts frequently originate from complex media components such as yeast extract, molasses, corn steep liquor and others. The exact composition of the media compounds depends strongly on the immediate experiment and is individually decided for each specific case. Information about media optimization is available in the textbook "Applied Microbiol. Physiology, A  
20 Practical Approach (*eds.* P.M. Rhodes, P.F. Stanbury, IRL Press (1997) pp. 53-73, ISBN 0 19 963577 3). It is also possible to select growth media from commercial suppliers, like standard 1 (Merck) or BHI (grain heart infusion, DIFCO) or others.

All medium components are sterilized, either by heat (20 minutes at 1.5 bar and 121°C) or by sterile filtration. The components can either be sterilized together or, if  
25 necessary, separately. All media components can be present at the beginning of growth, or they can optionally be added continuously or batchwise.

Culture conditions are defined separately for each experiment. The temperature should be in a range between 15°C and 45°C. The temperature can be kept constant or can be altered during the experiment. The pH of the medium should be in the range of 5 to  
30 8.5, preferably around 7.0, and can be maintained by the addition of buffers to the media. An exemplary buffer for this purpose is a potassium phosphate buffer. Synthetic buffers such as MOPS, HEPES, ACES and others can alternatively or simultaneously be used. It

is also possible to maintain a constant culture pH through the addition of NaOH or NH<sub>4</sub>OH during growth. If complex medium components such as yeast extract are utilized, the necessity for additional buffers may be reduced, due to the fact that many complex compounds have high buffer capacities. If a fermentor is utilized for culturing the micro-organisms, the pH can also be controlled using gaseous ammonia.

The incubation time is usually in a range from several hours to several days. This time is selected in order to permit the maximal amount of product to accumulate in the broth. The disclosed growth experiments can be carried out in a variety of vessels, such as microtiter plates, glass tubes, glass flasks or glass or metal fermentors of different sizes.

For screening a large number of clones, the microorganisms should be cultured in microtiter plates, glass tubes or shake flasks, either with or without baffles. Preferably 100 ml shake flasks are used, filled with 10% (by volume) of the required growth medium. The flasks should be shaken on a rotary shaker (amplitude 25 mm) using a speed-range of 100 – 300 rpm. Evaporation losses can be diminished by the maintenance of a humid atmosphere; alternatively, a mathematical correction for evaporation losses should be performed.

If genetically modified clones are tested, an unmodified control clone or a control clone containing the basic plasmid without any insert should also be tested. The medium is inoculated to an OD<sub>600</sub> of 0.5 – 1.5 using cells grown on agar plates, such as CM plates (10 g/l glucose, 2,5 g/l NaCl, 2 g/l urea, 10 g/l polypeptone, 5 g/l yeast extract, 5 g/l meat extract, 22 g/l NaCl, 2 g/l urea, 10 g/l polypeptone, 5 g/l yeast extract, 5 g/l meat extract, 22 g/l agar, pH 6.8 with 2M NaOH) that had been incubated at 30°C. Inoculation of the media is accomplished by either introduction of a saline suspension of *C. glutamicum* cells from CM plates or addition of a liquid preculture of this bacterium.

#### **Example 8 – *In vitro* Analysis of the Function of Mutant Proteins**

The determination of activities and kinetic parameters of enzymes is well established in the art. Experiments to determine the activity of any given altered enzyme must be tailored to the specific activity of the wild-type enzyme, which is well within the ability of one of ordinary skill in the art. Overviews about enzymes in general, as well as specific details concerning structure, kinetics, principles, methods, applications and examples for the determination of many enzyme activities may be

found, for example, in the following references: Dixon, M., and Webb, E.C., (1979) *Enzymes*. Longmans: London; Fersht, (1985) *Enzyme Structure and Mechanism*. Freeman: New York; Walsh, (1979) *Enzymatic Reaction Mechanisms*. Freeman: San Francisco; Price, N.C., Stevens, L. (1982) *Fundamentals of Enzymology*. Oxford Univ. Press: Oxford; Boyer, P.D., ed. (1983) *The Enzymes*, 3<sup>rd</sup> ed. Academic Press: New York; Bisswanger, H., (1994) *Enzymkinetik*, 2<sup>nd</sup> ed. VCH: Weinheim (ISBN 3527300325); Bergmeyer, H.U., Bergmeyer, J., Graßl, M., eds. (1983-1986) *Methods of Enzymatic Analysis*, 3<sup>rd</sup> ed., vol. I-XII, Verlag Chemie: Weinheim; and Ullmann's *Encyclopedia of Industrial Chemistry* (1987) vol. A9, "Enzymes". VCH: Weinheim, p. 352-363.

The activity of proteins which bind to DNA can be measured by several well-established methods, such as DNA band-shift assays (also called gel retardation assays). The effect of such proteins on the expression of other molecules can be measured using reporter gene assays (such as that described in Kolmar, H. *et al.* (1995) *EMBO J.* 14: 3895-3904 and references cited therein). Reporter gene test systems are well known and established for applications in both pro- and eukaryotic cells, using enzymes such as beta-galactosidase, green fluorescent protein, and several others.

The determination of activity of membrane-transport proteins can be performed according to techniques such as those described in Gennis, R.B. (1989) "Pores, Channels and Transporters", in *Biomembranes, Molecular Structure and Function*, Springer: Heidelberg, p. 85-137; 199-234; and 270-322.

#### **Example 9: Analysis of Impact of Mutant Protein on the Production of the Desired Product**

The effect of the genetic modification in *C. glutamicum* on production of a desired compound (such as an amino acid) can be assessed by growing the modified microorganism under suitable conditions (such as those described above) and analyzing the medium and/or the cellular component for increased production of the desired product (i.e., an amino acid). Such analysis techniques are well known to one of ordinary skill in the art, and include spectroscopy, thin layer chromatography, staining methods of various kinds, enzymatic and microbiological methods, and analytical chromatography such as high performance liquid chromatography (see, for example,



- Ullman, Encyclopedia of Industrial Chemistry, vol. A2, p. 89-90 and p. 443-613, VCH: Weinheim (1985); Fallon, A. *et al.*, (1987) "Applications of HPLC in Biochemistry" in: Laboratory Techniques in Biochemistry and Molecular Biology, vol. 17; Rehm *et al.* (1993) Biotechnology, vol. 3, Chapter III: "Product recovery and purification", page 469-714, VCH: Weinheim; Belter, P.A. *et al.* (1988) Bioseparations: downstream processing for biotechnology, John Wiley and Sons; Kennedy, J.F. and Cabral, J.M.S. (1992) Recovery processes for biological materials, John Wiley and Sons; Shaeiwitz, J.A. and Henry, J.D. (1988) Biochemical separations, in: Ulmann's Encyclopedia of Industrial Chemistry, vol. B3, Chapter 11, page 1-27, VCH: Weinheim; and Dechow, F.J. (1989) Separation and purification techniques in biotechnology, Noyes Publications.)

In addition to the measurement of the final product of fermentation, it is also possible to analyze other components of the metabolic pathways utilized for the production of the desired compound, such as intermediates and side-products, to determine the overall productivity of the organism, yield, and/or efficiency of production of the compound. Analysis methods include measurements of nutrient levels in the medium (*e.g.*, sugars, hydrocarbons, nitrogen sources, phosphate, and other ions), measurements of biomass composition and growth, analysis of the production of common metabolites of biosynthetic pathways, and measurement of gasses produced during fermentation. Standard methods for these measurements are outlined in Applied Microbial Physiology, A Practical Approach, P.M. Rhodes and P.F. Stanbury, eds., IRL Press, p. 103-129; 131-163; and 165-192 (ISBN: 0199635773) and references cited therein.

#### 25 **Example 10: Purification of the Desired Product from *C. glutamicum* Culture**

Recovery of the desired product from the *C. glutamicum* cells or supernatant of the above-described culture can be performed by various methods well known in the art. If the desired product is not secreted from the cells, the cells can be harvested from the culture by low-speed centrifugation, the cells can be lysed by standard techniques, such as mechanical force or sonication. The cellular debris is removed by centrifugation, and the supernatant fraction containing the soluble proteins is retained for further purification of the desired compound. If the product is secreted from the *C. glutamicum*

cells, then the cells are removed from the culture by low-speed centrifugation, and the supernate fraction is retained for further purification.

The supernatant fraction from either purification method is subjected to chromatography with a suitable resin, in which the desired molecule is either retained on a chromatography resin while many of the impurities in the sample are not, or where the impurities are retained by the resin while the sample is not. Such chromatography steps may be repeated as necessary, using the same or different chromatography resins. One of ordinary skill in the art would be well-versed in the selection of appropriate chromatography resins and in their most efficacious application for a particular molecule to be purified. The purified product may be concentrated by filtration or ultrafiltration, and stored at a temperature at which the stability of the product is maximized.

There are a wide array of purification methods known to the art and the preceding method of purification is not meant to be limiting. Such purification techniques are described, for example, in Bailey, J.E. & Ollis, D.F. *Biochemical Engineering Fundamentals*, McGraw-Hill: New York (1986).

The identity and purity of the isolated compounds may be assessed by techniques standard in the art. These include high-performance liquid chromatography (HPLC), spectroscopic methods, staining methods, thin layer chromatography, NIRS, enzymatic assay, or microbiologically. Such analysis methods are reviewed in: Patek *et al.* (1994) *Appl. Environ. Microbiol.* 60: 133-140; Malakhova *et al.* (1996) *Biotekhnologiya* 11: 27-32; and Schmidt *et al.* (1998) *Bioprocess Engineer.* 19: 67-70. Ulmann's Encyclopedia of Industrial Chemistry, (1996) vol. A27, VCH: Weinheim, p. 89-90, p. 521-540, p. 540-547, p. 559-566, 575-581 and p. 581-587; Michal, G. (1999) *Biochemical Pathways: An Atlas of Biochemistry and Molecular Biology*, John Wiley and Sons; Fallon, A. *et al.* (1987) *Applications of HPLC in Biochemistry in: Laboratory Techniques in Biochemistry and Molecular Biology*, vol. 17.

#### **Example 11: Analysis of the Gene Sequences of the Invention**

The comparison of sequences and determination of percent homology between two sequences are art-known techniques, and can be accomplished using a mathematical algorithm, such as the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264-68, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA*

90:5873-77. Such an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to PTS nucleic acid  
5 molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to PTS protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped  
10 BLAST programs, one of ordinary skill in the art will know how to optimize the parameters of the program (*e.g.*, XBLAST and NBLAST) for the specific sequence being analyzed.

Another example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Meyers and Miller ((1988) *Comput. Appl. Biosci.* 4: 11-  
15 17). Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used. Additional algorithms for sequence analysis are known in the art, and include ADVANCE and ADAM. described  
20 in Torelli and Robotti (1994) *Comput. Appl. Biosci.* 10:3-5; and FASTA, described in Pearson and Lipman (1988) *P.N.A.S.* 85:2444-8.

The percent homology between two amino acid sequences can also be accomplished using the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blosum 62 matrix or a PAM250 matrix, and a gap  
25 weight of 12, 10, 8, 6, or 4 and a length weight of 2, 3, or 4. The percent homology between two nucleic acid sequences can be accomplished using the GAP program in the GCG software package, using standard parameters, such as a gap weight of 50 and a length weight of 3.

A comparative analysis of the gene sequences of the invention with those present  
30 in Genbank has been performed using techniques known in the art (see, *e.g.*, Bexevanis and Ouellette, eds. (1998) *Bioinformatics: A Practical Guide to the Analysis of Genes and Proteins.* John Wiley and Sons: New York). The gene sequences of the invention

were compared to genes present in Genbank in a three-step process. In a first step, a BLASTN analysis (*e.g.*, a local alignment analysis) was performed for each of the sequences of the invention against the nucleotide sequences present in Genbank, and the top 500 hits were retained for further analysis. A subsequent FASTA search (*e.g.*, a  
5 combined local and global alignment analysis, in which limited regions of the sequences are aligned) was performed on these 500 hits. Each gene sequence of the invention was subsequently globally aligned to each of the top three FASTA hits, using the GAP program in the GCG software package (using standard parameters). In order to obtain correct results, the length of the sequences extracted from Genbank were adjusted to the  
10 length of the query sequences by methods well-known in the art. The results of this analysis are set forth in Table 4. The resulting data is identical to that which would have been obtained had a GAP (global) analysis alone been performed on each of the genes of the invention in comparison with each of the references in Genbank, but required significantly reduced computational time as compared to such a database-wide GAP  
15 (global) analysis. Sequences of the invention for which no alignments above the cutoff values were obtained are indicated on Table 4 by the absence of alignment information. It will further be understood by one of ordinary skill in the art that the GAP alignment homology percentages set forth in Table 4 under the heading "% homology (GAP)" are listed in the European numerical format, wherein a ',' represents a decimal point. For  
20 example, a value of "40,345" in this column represents "40.345%".

#### **Example 12: Construction and Operation of DNA Microarrays**

The sequences of the invention may additionally be used in the construction and application of DNA microarrays (the design, methodology, and uses of DNA arrays are  
25 well known in the art, and are described, for example, in Schena, M. *et al.* (1995) *Science* 270: 467-470; Wodicka, L. *et al.* (1997) *Nature Biotechnology* 15: 1359-1367; DeSaizieu, A. *et al.* (1998) *Nature Biotechnology* 16: 45-48; and DeRisi, J.L. *et al.* (1997) *Science* 278: 680-686).

DNA microarrays are solid or flexible supports consisting of nitrocellulose,  
30 nylon, glass, silicone, or other materials. Nucleic acid molecules may be attached to the surface in an ordered manner. After appropriate labeling, other nucleic acids or nucleic acid mixtures can be hybridized to the immobilized nucleic acid molecules, and the label

may be used to monitor and measure the individual signal intensities of the hybridized molecules at defined regions. This methodology allows the simultaneous quantification of the relative or absolute amount of all or selected nucleic acids in the applied nucleic acid sample or mixture. DNA microarrays, therefore, permit an analysis of the  
5 expression of multiple (as many as 6800 or more) nucleic acids in parallel (see, *e.g.*, Schena, M. (1996) *BioEssays* 18(5): 427-431).

The sequences of the invention may be used to design oligonucleotide primers which are able to amplify defined regions of one or more *C. glutamicum* genes by a nucleic acid amplification reaction such as the polymerase chain reaction. The choice  
10 and design of the 5' or 3' oligonucleotide primers or of appropriate linkers allows the covalent attachment of the resulting PCR products to the surface of a support medium described above (and also described, for example, Schena, M. *et al.* (1995) *Science* 270: 467-470).

Nucleic acid microarrays may also be constructed by *in situ* oligonucleotide  
15 synthesis as described by Wodicka, L. *et al.* (1997) *Nature Biotechnology* 15: 1359-1367. By photolithographic methods, precisely defined regions of the matrix are exposed to light. Protective groups which are photolabile are thereby activated and undergo nucleotide addition, whereas regions that are masked from light do not undergo any modification. Subsequent cycles of protection and light activation permit the  
20 synthesis of different oligonucleotides at defined positions. Small, defined regions of the genes of the invention may be synthesized on microarrays by solid phase oligonucleotide synthesis.

The nucleic acid molecules of the invention present in a sample or mixture of nucleotides may be hybridized to the microarrays. These nucleic acid molecules can be  
25 labeled according to standard methods. In brief, nucleic acid molecules (*e.g.*, mRNA molecules or DNA molecules) are labeled by the incorporation of isotopically or fluorescently labeled nucleotides, *e.g.*, during reverse transcription or DNA synthesis. Hybridization of labeled nucleic acids to microarrays is described (*e.g.*, in Schena, M. *et al.* (1995) *supra*; Wodicka, L. *et al.* (1997), *supra*; and DeSaizieu A. *et al.* (1998),  
30 *supra*). The detection and quantification of the hybridized molecule are tailored to the specific incorporated label. Radioactive labels can be detected, for example, as

described in Schena, M. *et al.* (1995) *supra*) and fluorescent labels may be detected, for example, by the method of Shalon *et al.* (1996) *Genome Research* 6: 639-645).

The application of the sequences of the invention to DNA microarray technology, as described above, permits comparative analyses of different strains of *C. glutamicum* or other Corynebacteria. For example, studies of inter-strain variations based on individual transcript profiles and the identification of genes that are important for specific and/or desired strain properties such as pathogenicity, productivity and stress tolerance are facilitated by nucleic acid array methodologies. Also, comparisons of the profile of expression of genes of the invention during the course of a fermentation reaction are possible using nucleic acid array technology.

### **Example 13: Analysis of the Dynamics of Cellular Protein Populations (Proteomics)**

The genes, compositions, and methods of the invention may be applied to study the interactions and dynamics of populations of proteins, termed 'proteomics'. Protein populations of interest include, but are not limited to, the total protein population of *C. glutamicum* (e.g., in comparison with the protein populations of other organisms), those proteins which are active under specific environmental or metabolic conditions (e.g., during fermentation, at high or low temperature, or at high or low pH), or those proteins which are active during specific phases of growth and development.

Protein populations can be analyzed by various well-known techniques, such as gel electrophoresis. Cellular proteins may be obtained, for example, by lysis or extraction, and may be separated from one another using a variety of electrophoretic techniques. Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) separates proteins largely on the basis of their molecular weight. Isoelectric focusing polyacrylamide gel electrophoresis (IEF-PAGE) separates proteins by their isoelectric point (which reflects not only the amino acid sequence but also posttranslational modifications of the protein). Another, more preferred method of protein analysis is the consecutive combination of both IEF-PAGE and SDS-PAGE, known as 2-D-gel electrophoresis (described, for example, in Hermann *et al.* (1998) *Electrophoresis* 19: 3217-3221; Fountoulakis *et al.* (1998) *Electrophoresis* 19: 1193-1202; Langen *et al.* (1997) *Electrophoresis* 18: 1184-1192; Antelmann *et al.* (1997) *Electrophoresis* 18:

1451-1463). Other separation techniques may also be utilized for protein separation, such as capillary gel electrophoresis; such techniques are well known in the art.

Proteins separated by these methodologies can be visualized by standard techniques, such as by staining or labeling. Suitable stains are known in the art, and  
5 include Coomassie Brilliant Blue, silver stain, or fluorescent dyes such as Sypro Ruby (Molecular Probes). The inclusion of radioactively labeled amino acids or other protein precursors (e.g.,  $^{35}\text{S}$ -methionine,  $^{35}\text{S}$ -cysteine,  $^{14}\text{C}$ -labelled amino acids,  $^{15}\text{N}$ -amino acids,  $^{15}\text{NO}_3$  or  $^{15}\text{NH}_4^+$  or  $^{13}\text{C}$ -labelled amino acids) in the medium of *C. glutamicum* permits the labeling of proteins from these cells prior to their separation. Similarly,  
10 fluorescent labels may be employed. These labeled proteins can be extracted, isolated and separated according to the previously described techniques.

Proteins visualized by these techniques can be further analyzed by measuring the amount of dye or label used. The amount of a given protein can be determined quantitatively using, for example, optical methods and can be compared to the amount  
15 of other proteins in the same gel or in other gels. Comparisons of proteins on gels can be made, for example, by optical comparison, by spectroscopy, by image scanning and analysis of gels, or through the use of photographic films and screens. Such techniques are well-known in the art.

To determine the identity of any given protein, direct sequencing or other  
20 standard techniques may be employed. For example, N- and/or C-terminal amino acid sequencing (such as Edman degradation) may be used, as may mass spectrometry (in particular MALDI or ESI techniques (see, e.g., Langen *et al.* (1997) *Electrophoresis* 18: 1184-1192)). The protein sequences provided herein can be used for the identification of *C. glutamicum* proteins by these techniques.

25 The information obtained by these methods can be used to compare patterns of protein presence, activity, or modification between different samples from various biological conditions (e.g., different organisms, time points of fermentation, media conditions, or different biotopes, among others). Data obtained from such experiments alone, or in combination with other techniques, can be used for various applications,  
30 such as to compare the behavior of various organisms in a given (e.g., metabolic) situation, to increase the productivity of strains which produce fine chemicals or to increase the efficiency of the production of fine chemicals.

**Equivalents**

Those of ordinary skill in the art will recognize, or will be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

5



What is claimed:

1. An isolated nucleic acid molecule from *Corynebacterium glutamicum* encoding a phosphoenolpyruvate: sugar phosphotransferase system protein, or a portion thereof, provided that the nucleic acid molecule does not consist of any of the F-designated genes set forth in Table 1.
2. The isolated nucleic acid molecule of claim 1, wherein said phosphoenolpyruvate: sugar phosphotransferase system protein is selected from the group consisting of proteins involved in the transport of glucose, sucrose, mannose, fructose, mannitol, raffinose, ribulose, ribose, lactose, maltose, sorbose, sorbitol, xylose, and galactose.
3. An isolated *Corynebacterium glutamicum* nucleic acid molecule selected from the group consisting of those sequences set forth as odd-numbered SEQ ID NOs of the Sequence Listing, or a portion thereof, provided that the nucleic acid molecule does not consist of any of the F-designated genes set forth in Table 1.
4. An isolated nucleic acid molecule which encodes a polypeptide sequence selected from the group consisting of those sequences set forth as even-numbered SEQ ID NOs of the Sequence Listing, provided that the nucleic acid molecule does not consist of any of the F-designated genes set forth in Table 1.
5. An isolated nucleic acid molecule which encodes a naturally occurring allelic variant of a polypeptide selected from the group of amino acid sequences consisting of those sequences set forth as even-numbered SEQ ID NOs of the Sequence Listing, provided that the nucleic acid molecule does not consist of any of the F-designated genes set forth in Table 1.
6. An isolated nucleic acid molecule comprising a nucleotide sequence which is at least 50% homologous to a nucleotide sequence selected from the group consisting of

those sequences set forth as odd-numbered SEQ ID NOs of the Sequence Listing , or a portion thereof, provided that the nucleic acid molecule does not consist of any of the F-designated genes set forth in Table 1.

5        7. An isolated nucleic acid molecule comprising a fragment of at least 15 nucleotides of a nucleic acid comprising a nucleotide sequence selected from the group consisting of those sequences set forth as odd-numbered SEQ ID NOs of the Sequence Listing , provided that the nucleic acid molecule does not consist of any of the F-designated genes set forth in Table 1.

10

8. An isolated nucleic acid molecule which hybridizes to the nucleic acid molecule of any one of claims 1-7 under stringent conditions.

9. An isolated nucleic acid molecule comprising the nucleic acid molecule of any  
15 one of claims 1-8 or a portion thereof and a nucleotide sequence encoding a heterologous polypeptide.

10. A vector comprising the nucleic acid molecule of any one of claims 1-9.

20        11. The vector of claim 10, which is an expression vector.

12. A host cell transfected with the expression vector of claim 11.

13. The host cell of claim 12, wherein said cell is a microorganism.

25

14. The host cell of claim 13, wherein said cell belongs to the genus *Corynebacterium* or *Brevibacterium*.

15. The host cell of claim 12, wherein the expression of said nucleic acid molecule  
30 results in the modulation in production of a fine chemical from said cell.

16. The host cell of claim 15, wherein said fine chemical is selected from the group consisting of: organic acids, proteinogenic amino acids, nonproteinogenic amino acids, purine and pyrimidine bases, nucleosides, nucleotides, lipids, saturated and unsaturated fatty acids, diols, carbohydrates, aromatic compounds, vitamins, cofactors, polyketides,  
5 and enzymes.

17. A method of producing a polypeptide comprising culturing the host cell of claim 12 in an appropriate culture medium to, thereby, produce the polypeptide.

10 18. An isolated phosphoenolpyruvate: sugar phosphotransferase system polypeptide from *Corynebacterium glutamicum*, or a portion thereof.

19. The protein of claim 18, wherein said phosphoenolpyruvate: sugar phosphotransferase system protein is selected from the group consisting of proteins  
15 involved in the transport of glucose, sucrose, mannose, fructose, mannitol, raffinose, ribulose, ribose, lactose, maltose, sorbose, and galactose.

20. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of those sequences set forth as even-numbered SEQ ID NOs of the  
20 Sequence Listing, provided that the amino acid sequence is not encoded by any of the F-designated genes set forth in Table 1.

21. An isolated polypeptide comprising a naturally occurring allelic variant of a polypeptide comprising an amino acid sequence selected from the group consisting of  
25 those sequences set forth as even-numbered SEQ ID NOs of the Sequence Listing, or a portion thereof, provided that the amino acid sequence is not encoded by any of the F-designated genes set forth in Table 1.

22. The isolated polypeptide of any of claims 18-21, further comprising heterologous  
30 amino acid sequences.

23. An isolated polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 50% homologous to a nucleic acid selected from the group consisting of those sequences set forth as odd-numbered SEQ ID NOs of the Sequence Listing, provided that the nucleic acid molecule does not consist of any of the F-designated nucleic acid molecules set forth in Table 1.

24. An isolated polypeptide comprising an amino acid sequence which is at least 50% homologous to an amino acid sequence selected from the group consisting of those sequences set forth as even-numbered SEQ ID NOs of the Sequence Listing, provided that the amino acid sequence is not encoded by any of the F-designated genes set forth in Table 1.

25. A method for producing a fine chemical, comprising culturing a cell containing a vector of claim 12 such that the fine chemical is produced.

26. The method of claim 25, wherein said method further comprises the step of recovering the fine chemical from said culture.

27. The method of claim 25, wherein said method further comprises the step of transfecting said cell with the vector of claim 11 to result in a cell containing said vector.

28. The method of claim 25, wherein said cell belongs to the genus *Corynebacterium* or *Brevibacterium*.

29. The method of claim 25, wherein said cell is selected from the group consisting of: *Corynebacterium glutamicum*, *Corynebacterium herculis*, *Corynebacterium lilium*, *Corynebacterium acetoacidophilum*, *Corynebacterium acetoglutamicum*, *Corynebacterium acetophilum*, *Corynebacterium ammoniagenes*, *Corynebacterium fujiokense*, *Corynebacterium nitrilophilus*, *Brevibacterium ammoniagenes*, *Brevibacterium butanicum*, *Brevibacterium divaricatum*, *Brevibacterium flavum*, *Brevibacterium healii*, *Brevibacterium ketoglutamicum*, *Brevibacterium*

*ketosoreductum*, *Brevibacterium lactofermentum*, *Brevibacterium linens*,  
*Brevibacterium paraffinolyticum*, and those strains set forth in Table 3.

30. The method of claim 25, wherein expression of the nucleic acid molecule from  
5 said vector results in modulation of production of said fine chemical.

31. The method of claim 25, wherein said fine chemical is selected from the group  
consisting of: organic acids, proteinogenic amino acids, nonproteinogenic amino acids,  
purine and pyrimidine bases, nucleosides, nucleotides, lipids, saturated and unsaturated  
10 fatty acids, diols, carbohydrates, aromatic compounds, vitamins, cofactors, polyketides,  
and enzymes.

32. The method of claim 25, wherein said fine chemical is an amino acid.

15 33. The method of claim 32, wherein said amino acid is drawn from the group  
consisting of: lysine, glutamate, glutamine, alanine, aspartate, glycine, serine, threonine,  
methionine, cysteine, valine, leucine, isoleucine, arginine, proline, histidine, tyrosine,  
phenylalanine, and tryptophan.

20 34. A method for producing a fine chemical, comprising culturing a cell whose  
genomic DNA has been altered by the inclusion of a nucleic acid molecule of any one of  
claims 1-9.

25 35. A method for diagnosing the presence or activity of *Corynebacterium*  
*diphtheriae* in a subject, comprising detecting the presence of one or more of SEQ ID  
NOs 1 through 34 of the Sequence Listing in the subject, provided that the sequences are  
not or are not encoded by any of the F-designated sequences set forth in Table 1, thereby  
diagnosing the presence or activity of *Corynebacterium diphtheriae* in the subject.

30 36. A host cell comprising a nucleic acid molecule selected from the group  
consisting of the nucleic acid molecules set forth as odd-numbered SEQ ID NOs of the  
Sequence Listing, wherein the nucleic acid molecule is disrupted.

37. A host cell comprising a nucleic acid molecule selected from the group consisting of the nucleic acid molecules set forth as odd-numbered SEQ ID NOs of the Sequence Listing , wherein the nucleic acid molecule comprises one or more nucleic  
5 acid modifications from the sequence set forth in as odd-numbered SEQ ID NOs of the Sequence Listing .

38. A host cell comprising a nucleic acid molecule selected from the group consisting of the nucleic acid molecules set forth as odd-numbered SEQ ID NOs of the  
10 Sequence Listing , wherein the regulatory region of the nucleic acid molecule is modified relative to the wild-type regulatory region of the molecule.

## SEQUENCE LISTING

<110> BASF Aktiengesellschaft  
 <120> CORYNEBACTERIUM GLUTAMICUM GENES ENCODING  
 PHOSPHOENOLPYRUVATE:SUGAR PHOSPHOTRANSFERASE  
 SYSTEM PROTEINS

<130> BGI-122CPPC

<140>

<141>

<160> 34

<210> 1

<211> 1527

<212> DNA

<213> Corynebacterium glutamicum

<220>

<221> CDS

<222> (101)..(1504)

<223> RXS00315

<400> 1

ctcatggcat ctgcgccgtt cgcgttcttg ccagtgttgg ttggtttcac cgcaaccaag 60

cgtttcggcg gcaatgagtt cctgggcgcc gcgtattggt atg gcg atg gtg ttc 115  
 Met Ala Met Val Phe  
 1 5

ccg agc ttg gtg aac ggc tac gac gtg gcc gcc acc atg gct gcg ggc 163  
 Pro Ser Leu Val Asn Gly Tyr Asp Val Ala Ala Thr Met Ala Ala Gly  
 10 15 20

gaa atg cca atg tgg tcc ctg ttt ggt tta gat gtt gcc caa gcc ggt 211  
 Glu Met Pro Met Trp Ser Leu Phe Gly Leu Asp Val Ala Gln Ala Gly  
 25 30 35

tac cag ggc acc gtg ctt cct gtg ctg gtg gtt tct tgg att ctg gca 259  
 Tyr Gln Gly Thr Val Leu Pro Val Leu Val Val Ser Trp Ile Leu Ala  
 40 45 50

acg atc gag aag ttc ctg cac aag cga ctc aag ggc act gca gac ttc 307  
 Thr Ile Glu Lys Phe Leu His Lys Arg Leu Lys Gly Thr Ala Asp Phe  
 55 60 65

ctg atc act cca gtg ctg acg ttg ctg ctc acc gga ttc ctt aca ttc 355  
 Leu Ile Thr Pro Val Leu Thr Leu Leu Leu Thr Gly Phe Leu Thr Phe  
 70 75 80 85

atc gcc att ggc cca gca atg cgc tgg gtg ggc gat gtg ctg gca cac 403  
 Ile Ala Ile Gly Pro Ala Met Arg Trp Val Gly Asp Val Leu Ala His  
 90 95 100

ggt cta cag gga ctt tat gat ttc ggt ggt cca gtc ggc ggt ctg ctc 451  
 Gly Leu Gln Gly Leu Tyr Asp Phe Gly Gly Pro Val Gly Gly Leu Leu  
 105 110 115

ttc ggt ctg gtc tac tca cca atc gtc atc act ggt ctg cac cag tcc 499  
 Phe Gly Leu Val Tyr Ser Pro Ile Val Ile Thr Gly Leu His Gln Ser  
 120 125 130

ttc ccg cca att gag ctg gag ctg ttt aac cag ggt gga tcc ttc atc 547

Phe	Pro	Pro	Ile	Glu	Leu	Glu	Leu	Phe	Asn	Gln	Gly	Gly	Ser	Phe	Ile		
135						140					145						
ttc	gca	acg	gca	tct	atg	gct	aat	atc	gcc	cag	ggg	gcg	gca	tgt	ttg	595	
Phe	Ala	Thr	Ala	Ser	Met	Ala	Asn	Ile	Ala	Gln	Gly	Ala	Ala	Cys	Leu		
150					155					160					165		
gca	gtg	ttc	ttc	ctg	gcg	aag	agt	gaa	aag	ctc	aag	ggc	ctt	gca	ggg	643	
Ala	Val	Phe	Phe	Leu	Ala	Lys	Ser	Glu	Lys	Leu	Lys	Gly	Leu	Ala	Gly		
				170					175					180			
gct	tca	ggg	gtc	tcc	gct	gtt	ctt	ggg	att	acg	gag	cct	gcg	atc	ttc	691	
Ala	Ser	Gly	Val	Ser	Ala	Val	Leu	Gly	Ile	Thr	Glu	Pro	Ala	Ile	Phe		
			185					190					195				
ggg	gtg	aac	ctt	cgc	ctg	cgc	tgg	ccg	ttc	ttc	atc	ggg	atc	ggg	acc	739	
Gly	Val	Asn	Leu	Arg	Leu	Arg	Trp	Pro	Phe	Phe	Ile	Gly	Ile	Gly	Thr		
		200					205					210					
gca	gct	atc	ggg	ggc	gct	ttg	att	gca	ctc	ttt	aat	atc	aag	gca	gtt	787	
Ala	Ala	Ile	Gly	Gly	Ala	Leu	Ile	Ala	Leu	Phe	Asn	Ile	Lys	Ala	Val		
	215					220					225						
gcg	ttg	ggc	gct	gca	ggg	ttc	ttg	ggg	gtt	gtt	tct	att	gat	gct	cca	835	
Ala	Leu	Gly	Ala	Ala	Gly	Phe	Leu	Gly	Val	Val	Ser	Ile	Asp	Ala	Pro		
230					235					240					245		
gat	atg	gtc	atg	ttc	ttg	gtg	tgt	gca	gtt	gtt	acc	ttc	ttc	atc	gca	883	
Asp	Met	Val	Met	Phe	Leu	Val	Cys	Ala	Val	Val	Thr	Phe	Phe	Ile	Ala		
				250					255					260			
ttc	ggc	gca	gcg	att	gct	tat	ggc	ctt	tac	ttg	gtt	cgc	cgc	aac	ggc	931	
Phe	Gly	Ala	Ala	Ile	Ala	Tyr	Gly	Leu	Tyr	Leu	Val	Arg	Arg	Asn	Gly		
			265					270					275				
agc	att	gat	cca	gat	gca	acc	gct	gct	cca	gtg	cct	gca	gga	acg	acc	979	
Ser	Ile	Asp	Pro	Asp	Ala	Thr	Ala	Ala	Pro	Val	Pro	Ala	Gly	Thr	Thr		
		280					285					290					
aaa	gcc	gaa	gca	gaa	gca	ccc	gca	gaa	ttt	tca	aac	gat	tcc	acc	atc	1027	
Lys	Ala	Glu	Ala	Glu	Ala	Pro	Ala	Glu	Phe	Ser	Asn	Asp	Ser	Thr	Ile		
	295					300					305						
atc	cag	gca	cct	ttg	acc	ggg	gaa	gct	att	gca	ctg	agc	agc	gtc	agc	1075	
Ile	Gln	Ala	Pro	Leu	Thr	Gly	Glu	Ala	Ile	Ala	Leu	Ser	Ser	Val	Ser		
310					315					320					325		
gat	gcc	atg	ttt	gcc	agc	gga	aag	ctt	ggc	tcg	ggc	gtt	gcc	atc	gtc	1123	
Asp	Ala	Met	Phe	Ala	Ser	Gly	Lys	Leu	Gly	Ser	Gly	Val	Ala	Ile	Val		
				330					335					340			
cca	acc	aag	ggg	cag	tta	gtt	tct	ccg	gtg	agt	gga	aag	att	gtg	gtg	1171	
Pro	Thr	Lys	Gly	Gln	Leu	Val	Ser	Pro	Val	Ser	Gly	Lys	Ile	Val	Val		
			345					350					355				
gca	ttc	cca	tct	ggc	cat	gct	ttc	gca	gtt	cgc	acc	aag	gct	gag	gat	1219	
Ala	Phe	Pro	Ser	Gly	His	Ala	Phe	Ala	Val	Arg	Thr	Lys	Ala	Glu	Asp		
		360					365						370				
ggg	tcc	aat	gtg	gat	atc	ttg	atg	cac	att	ggg	ttc	gac	aca	gta	aac	1267	
Gly	Ser	Asn	Val	Asp	Ile	Leu	Met	His	Ile	Gly	Phe	Asp	Thr	Val	Asn		



375                      380                      385

ctc aac ggc acg cac ttt aac ccg ctg aag aag cag ggc gat gaa gtc 1315  
 Leu Asn Gly Thr His Phe Asn Pro Leu Lys Lys Gln Gly Asp Glu Val  
 390                      395                      400                      405

aaa gca ggg gag ctg ctg tgt gaa ttc gat att gat gcc att aag gct 1363  
 Lys Ala Gly Glu Leu Leu Cys Glu Phe Asp Ile Asp Ala Ile Lys Ala  
                     410                      415                      420

gca ggt tat gag gta acc acg ccg att gtt gtt tcg aat tac aag aaa 1411  
 Ala Gly Tyr Glu Val Thr Thr Pro Ile Val Val Ser Asn Tyr Lys Lys  
                     425                      430                      435

acc gga cct gta aac act tac ggt ttg ggc gaa att gaa gcg gga gcc 1459  
 Thr Gly Pro Val Asn Thr Tyr Gly Leu Gly Glu Ile Glu Ala Gly Ala  
                     440                      445                      450

aac ctg ctc aac gtc gca aag aaa gaa gcg gtg cca gca aca cca 1504  
 Asn Leu Leu Asn Val Ala Lys Lys Glu Ala Val Pro Ala Thr Pro  
                     455                      460                      465

taagttgaaa ccttgagtgt tcg 1527

<210> 2  
 <211> 468  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 2  
 Met Ala Met Val Phe Pro Ser Leu Val Asn Gly Tyr Asp Val Ala Ala  
   1                      5                      10                      15

Thr Met Ala Ala Gly Glu Met Pro Met Trp Ser Leu Phe Gly Leu Asp  
                     20                      25                      30

Val Ala Gln Ala Gly Tyr Gln Gly Thr Val Leu Pro Val Leu Val Val  
                     35                      40                      45

Ser Trp Ile Leu Ala Thr Ile Glu Lys Phe Leu His Lys Arg Leu Lys  
                     50                      55                      60

Gly Thr Ala Asp Phe Leu Ile Thr Pro Val Leu Thr Leu Leu Leu Thr  
   65                      70                      75                      80

Gly Phe Leu Thr Phe Ile Ala Ile Gly Pro Ala Met Arg Trp Val Gly  
                     85                      90                      95

Asp Val Leu Ala His Gly Leu Gln Gly Leu Tyr Asp Phe Gly Gly Pro  
                     100                      105                      110

Val Gly Gly Leu Leu Phe Gly Leu Val Tyr Ser Pro Ile Val Ile Thr  
                     115                      120                      125

Gly Leu His Gln Ser Phe Pro Pro Ile Glu Leu Glu Leu Phe Asn Gln  
   130                      135                      140

Gly Gly Ser Phe Ile Phe Ala Thr Ala Ser Met Ala Asn Ile Ala Gln  
   145                      150                      155                      160

Gly Ala Ala Cys Leu Ala Val Phe Phe Leu Ala Lys Ser Glu Lys Leu  
 165 170 175  
 Lys Gly Leu Ala Gly Ala Ser Gly Val Ser Ala Val Leu Gly Ile Thr  
 180 185 190  
 Glu Pro Ala Ile Phe Gly Val Asn Leu Arg Leu Arg Trp Pro Phe Phe  
 195 200 205  
 Ile Gly Ile Gly Thr Ala Ala Ile Gly Gly Ala Leu Ile Ala Leu Phe  
 210 215 220  
 Asn Ile Lys Ala Val Ala Leu Gly Ala Ala Gly Phe Leu Gly Val Val  
 225 230 235 240  
 Ser Ile Asp Ala Pro Asp Met Val Met Phe Leu Val Cys Ala Val Val  
 245 250 255  
 Thr Phe Phe Ile Ala Phe Gly Ala Ala Ile Ala Tyr Gly Leu Tyr Leu  
 260 265 270  
 Val Arg Arg Asn Gly Ser Ile Asp Pro Asp Ala Thr Ala Ala Pro Val  
 275 280 285  
 Pro Ala Gly Thr Thr Lys Ala Glu Ala Glu Ala Pro Ala Glu Phe Ser  
 290 295 300  
 Asn Asp Ser Thr Ile Ile Gln Ala Pro Leu Thr Gly Glu Ala Ile Ala  
 305 310 315 320  
 Leu Ser Ser Val Ser Asp Ala Met Phe Ala Ser Gly Lys Leu Gly Ser  
 325 330 335  
 Gly Val Ala Ile Val Pro Thr Lys Gly Gln Leu Val Ser Pro Val Ser  
 340 345 350  
 Gly Lys Ile Val Val Ala Phe Pro Ser Gly His Ala Phe Ala Val Arg  
 355 360 365  
 Thr Lys Ala Glu Asp Gly Ser Asn Val Asp Ile Leu Met His Ile Gly  
 370 375 380  
 Phe Asp Thr Val Asn Leu Asn Gly Thr His Phe Asn Pro Leu Lys Lys  
 385 390 395 400  
 Gln Gly Asp Glu Val Lys Ala Gly Glu Leu Leu Cys Glu Phe Asp Ile  
 405 410 415  
 Asp Ala Ile Lys Ala Ala Gly Tyr Glu Val Thr Thr Pro Ile Val Val  
 420 425 430  
 Ser Asn Tyr Lys Lys Thr Gly Pro Val Asn Thr Tyr Gly Leu Gly Glu  
 435 440 445  
 Ile Glu Ala Gly Ala Asn Leu Leu Asn Val Ala Lys Lys Glu Ala Val  
 450 455 460  
 Pro Ala Thr Pro  
 465

<210> 3  
 <211> 1109  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (1)..(1086)  
 <223> FRXA00315

<400> 3  
 tat gat ttc ggc ggt cca gtc ggc ggt ctg ctc ttc ggt ctg gtc tac 48  
 Tyr Asp Phe Gly Gly Pro Val Gly Gly Leu Leu Phe Gly Leu Val Tyr  
 1 5 10 15  
 tca cca atc gtc atc act ggt ctg cac cag tcc ttc ccg cca att gag 96  
 Ser Pro Ile Val Ile Thr Gly Leu His Gln Ser Phe Pro Pro Ile Glu  
 20 25 30  
 ctg gag ctg ttt aac cag ggt gga tcc ttc atc ttc gca acg gca tct 144  
 Leu Glu Leu Phe Asn Gln Gly Gly Ser Phe Ile Phe Ala Thr Ala Ser  
 35 40 45  
 atg gct aat atc gcc cag ggt gcg gca tgt ttg gca gtg ttc ttc ctg 192  
 Met Ala Asn Ile Ala Gln Gly Ala Ala Cys Leu Ala Val Phe Phe Leu  
 50 55 60  
 gcg aag agt gaa aag ctc aag ggc ctt gca ggt gct tca ggt gtc tcc 240  
 Ala Lys Ser Glu Lys Leu Lys Gly Leu Ala Gly Ala Ser Gly Val Ser  
 65 70 75 80  
 gct gtt ctt ggt att acg gag cct gcg atc ttc ggt gtg aac ctt cgc 288  
 Ala Val Leu Gly Ile Thr Glu Pro Ala Ile Phe Gly Val Asn Leu Arg  
 85 90 95  
 ctg cgc tgg ccg ttc ttc atc ggt atc ggt acc gca gct atc ggt ggc 336  
 Leu Arg Trp Pro Phe Phe Ile Gly Ile Gly Thr Ala Ala Ile Gly Gly  
 100 105 110  
 gct ttg att gca ctc ttt aat atc aag gca gtt gcg ttg ggc gct gca 384  
 Ala Leu Ile Ala Leu Phe Asn Ile Lys Ala Val Ala Leu Gly Ala Ala  
 115 120 125  
 ggt ttc ttg ggt gtt gtt tct att gat gct cca gat atg gtc atg ttc 432  
 Gly Phe Leu Gly Val Val Ser Ile Asp Ala Pro Asp Met Val Met Phe  
 130 135 140  
 ttg gtg tgt gca gtt gtt acc ttc ttc atc gca ttc ggc gca gcg att 480  
 Leu Val Cys Ala Val Val Thr Phe Phe Ile Ala Phe Gly Ala Ala Ile  
 145 150 155 160  
 gct tat ggc ctt tac ttg gtt cgc cgc aac ggc agc att gat cca gat 528  
 Ala Tyr Gly Leu Tyr Leu Val Arg Arg Asn Gly Ser Ile Asp Pro Asp  
 165 170 175  
 gca acc gct gct cca gtg cct gca gga acg acc aaa gcc gaa gca gaa 576  
 Ala Thr Ala Ala Pro Val Pro Ala Gly Thr Thr Lys Ala Glu Ala Glu  
 180 185 190  
 gca ccc gca gaa ttt tca aac gat tcc acc atc atc cag gca cct ttg 624  
 Ala Pro Ala Glu Phe Ser Asn Asp Ser Thr Ile Ile Gln Ala Pro Leu

195	200	205	
acc ggt gaa gct att gca ctg agc agc gtc agc gat gcc atg ttt gcc Thr Gly Glu Ala Ile Ala Leu Ser Ser Val Ser Asp Ala Met Phe Ala 210 215 220			672
agc gga aag ctt ggc tcg ggc gtt gcc atc gtc cca acc aag ggg cag Ser Gly Lys Leu Gly Ser Gly Val Ala Ile Val Pro Thr Lys Gly Gln 225 230 235 240			720
tta gtt tct ccg gtg agt gga aag att gtg gtg gca ttc cca tct ggc Leu Val Ser Pro Val Ser Gly Lys Ile Val Val Ala Phe Pro Ser Gly 245 250 255			768
cat gct ttc gca gtt cgc acc aag gct gag gat ggt tcc aat gtg gat His Ala Phe Ala Val Arg Thr Lys Ala Glu Asp Gly Ser Asn Val Asp 260 265 270			816
atc ttg atg cac att ggt ttc gac aca gta aac ctc aac ggc acg cac Ile Leu Met His Ile Gly Phe Asp Thr Val Asn Leu Asn Gly Thr His 275 280 285			864
ttt aac ccg ctg aag aag cag ggc gat gaa gtc aaa gca ggg gag ctg Phe Asn Pro Leu Lys Lys Gln Gly Asp Glu Val Lys Ala Gly Glu Leu 290 295 300			912
ctg tgt gaa ttc gat att gat gcc att aag gct gca ggt tat gag gta Leu Cys Glu Phe Asp Ile Asp Ala Ile Lys Ala Ala Gly Tyr Glu Val 305 310 315 320			960
acc acg ccg att gtt gtt tcg aat tac aag aaa acc gga cct gta aac Thr Thr Pro Ile Val Val Ser Asn Tyr Lys Lys Thr Gly Pro Val Asn 325 330 335			1008
act tac ggt ttg ggc gaa att gaa gcg gga gcc aac ctg ctc aac gtc Thr Tyr Gly Leu Gly Glu Ile Glu Ala Gly Ala Asn Leu Leu Asn Val 340 345 350			1056
gca aag aaa gaa gcg gtg cca gca aca cca taagttgaaa ccttgagtgt Ala Lys Lys Glu Ala Val Pro Ala Thr Pro 355 360			1106
tcg			1109

&lt;210&gt; 4

&lt;211&gt; 362

&lt;212&gt; PRT

&lt;213&gt; Corynebacterium glutamicum

&lt;400&gt; 4

Tyr Asp Phe Gly Gly Pro Val Gly Gly Leu Leu Phe Gly Leu Val Tyr  
1 5 10 15

Ser Pro Ile Val Ile Thr Gly Leu His Gln Ser Phe Pro Pro Ile Glu  
20 25 30

Leu Glu Leu Phe Asn Gln Gly Gly Ser Phe Ile Phe Ala Thr Ala Ser  
35 40 45

Met Ala Asn Ile Ala Gln Gly Ala Ala Cys Leu Ala Val Phe Phe Leu

50					55					60					
Ala	Lys	Ser	Glu	Lys	Leu	Lys	Gly	Leu	Ala	Gly	Ala	Ser	Gly	Val	Ser
65					70					75					80
Ala	Val	Leu	Gly	Ile	Thr	Glu	Pro	Ala	Ile	Phe	Gly	Val	Asn	Leu	Arg
				85					90					95	
Leu	Arg	Trp	Pro	Phe	Phe	Ile	Gly	Ile	Gly	Thr	Ala	Ala	Ile	Gly	Gly
			100					105					110		
Ala	Leu	Ile	Ala	Leu	Phe	Asn	Ile	Lys	Ala	Val	Ala	Leu	Gly	Ala	Ala
		115					120					125			
Gly	Phe	Leu	Gly	Val	Val	Ser	Ile	Asp	Ala	Pro	Asp	Met	Val	Met	Phe
	130					135					140				
Leu	Val	Cys	Ala	Val	Val	Thr	Phe	Phe	Ile	Ala	Phe	Gly	Ala	Ala	Ile
145					150					155					160
Ala	Tyr	Gly	Leu	Tyr	Leu	Val	Arg	Arg	Asn	Gly	Ser	Ile	Asp	Pro	Asp
				165					170					175	
Ala	Thr	Ala	Ala	Pro	Val	Pro	Ala	Gly	Thr	Thr	Lys	Ala	Glu	Ala	Glu
			180					185					190		
Ala	Pro	Ala	Glu	Phe	Ser	Asn	Asp	Ser	Thr	Ile	Ile	Gln	Ala	Pro	Leu
		195					200					205			
Thr	Gly	Glu	Ala	Ile	Ala	Leu	Ser	Ser	Val	Ser	Asp	Ala	Met	Phe	Ala
	210					215					220				
Ser	Gly	Lys	Leu	Gly	Ser	Gly	Val	Ala	Ile	Val	Pro	Thr	Lys	Gly	Gln
225					230					235					240
Leu	Val	Ser	Pro	Val	Ser	Gly	Lys	Ile	Val	Val	Ala	Phe	Pro	Ser	Gly
				245					250					255	
His	Ala	Phe	Ala	Val	Arg	Thr	Lys	Ala	Glu	Asp	Gly	Ser	Asn	Val	Asp
			260					265					270		
Ile	Leu	Met	His	Ile	Gly	Phe	Asp	Thr	Val	Asn	Leu	Asn	Gly	Thr	His
		275					280					285			
Phe	Asn	Pro	Leu	Lys	Lys	Gln	Gly	Asp	Glu	Val	Lys	Ala	Gly	Glu	Leu
	290					295					300				
Leu	Cys	Glu	Phe	Asp	Ile	Asp	Ala	Ile	Lys	Ala	Ala	Gly	Tyr	Glu	Val
305					310					315				320	
Thr	Thr	Pro	Ile	Val	Val	Ser	Asn	Tyr	Lys	Lys	Thr	Gly	Pro	Val	Asn
				325					330					335	
Thr	Tyr	Gly	Leu	Gly	Glu	Ile	Glu	Ala	Gly	Ala	Asn	Leu	Leu	Asn	Val
			340					345					350		
Ala	Lys	Lys	Glu	Ala	Val	Pro	Ala	Thr	Pro						
		355					360								

&lt;210&gt; 5

<211> 372  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (101)..(349)  
 <223> RXA01503

<400> 5  
 gtatcctcaa aggccttcta gctgttgacg ctgcagcgca ctcggtggat acgacatcca 60  
 cgacctatca aattctttat gctgcaggcg atgccttttc atg ttc ttg gca gtc 115  
 Met Phe Leu Ala Val  
 1 5  
 att ttg gcg att act gcg gct cgt aaa ttc ggt gcc aat gtc ttt aca 163  
 Ile Leu Ala Ile Thr Ala Ala Arg Lys Phe Gly Ala Asn Val Phe Thr  
 10 15 20  
 tca gtc gca ctc gct ggt gca ttg ctg cac aca cag ctt cag gca gta 211  
 Ser Val Ala Leu Ala Gly Ala Leu Leu His Thr Gln Leu Gln Ala Val  
 25 30 35  
 acc gtg ttg gtt gac ggt gaa ctc cag tcg atg act ctg gtg gct ttc 259  
 Thr Val Leu Val Asp Gly Glu Leu Gln Ser Met Thr Leu Val Ala Phe  
 40 45 50  
 caa aag gct ggt aat gac gtc acc ttc ctg ggc att cca gtg gtg ctg 307  
 Gln Lys Ala Gly Asn Asp Val Thr Phe Leu Gly Ile Pro Val Val Leu  
 55 60 65  
 cag ttg gcg ttg cat gta gcg agt ttg atg aag ttg tcg cga 349  
 Gln Leu Ala Leu His Val Ala Ser Leu Met Lys Leu Ser Arg  
 70 75 80  
 taagaggagg ggcgtgtcgg tct 372

<210> 6  
 <211> 83  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 6  
 Met Phe Leu Ala Val Ile Leu Ala Ile Thr Ala Ala Arg Lys Phe Gly  
 1 5 10 15  
 Ala Asn Val Phe Thr Ser Val Ala Leu Ala Gly Ala Leu Leu His Thr  
 20 25 30  
 Gln Leu Gln Ala Val Thr Val Leu Val Asp Gly Glu Leu Gln Ser Met  
 35 40 45  
 Thr Leu Val Ala Phe Gln Lys Ala Gly Asn Asp Val Thr Phe Leu Gly  
 50 55 60  
 Ile Pro Val Val Leu Gln Leu Ala Leu His Val Ala Ser Leu Met Lys  
 65 70 75 80  
 Leu Ser Arg

<210> 7  
 <211> 2187  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (101)..(2164)  
 <223> RXN01299

<400> 7

```
cgactgcggc gtctcttcct ggcactacca ttcctcgctc tgaccaactc gccacagctg 60
gtgcaacggt cacccaagtc aaaggattga aagaatcagc atg aat agc gta aat 115
                                         Met Asn Ser Val Asn
                                         1 5

aat tcc tcg ctt gtc cgg ctg gat gtc gat ttc ggc gac tcc acc acg 163
Asn Ser Ser Leu Val Arg Leu Asp Val Asp Phe Gly Asp Ser Thr Thr
                        10 15 20

gat gtc atc aac aac ctt gcc act gtt att ttc gac gct ggc cga gct 211
Asp Val Ile Asn Asn Leu Ala Thr Val Ile Phe Asp Ala Gly Arg Ala
                        25 30 35

tcc tcc gcc gac gcc ctt gcc aaa gac gcg ctg gat cgt gaa gca aag 259
Ser Ser Ala Asp Ala Leu Ala Lys Asp Ala Leu Asp Arg Glu Ala Lys
                        40 45 50

tcc ggc acc ggc gtt cct ggt caa gtt gct atc ccc cac tgc cgt tcc 307
Ser Gly Thr Gly Val Pro Gly Gln Val Ala Ile Pro His Cys Arg Ser
                        55 60 65

gaa gcc gta tct gtc cct acc ttg ggc ttt gct cgc ctg agc aag ggt 355
Glu Ala Val Ser Val Pro Thr Leu Gly Phe Ala Arg Leu Ser Lys Gly
                        70 75 80 85

gtg gac ttc agc gga cct gat ggc gat gcc aac ttg gtg ttc ctc att 403
Val Asp Phe Ser Gly Pro Asp Gly Asp Ala Asn Leu Val Phe Leu Ile
                        90 95 100

gca gca cct gct ggc ggc ggc aaa gag cac ctg aag atc ctg tcc aag 451
Ala Ala Pro Ala Gly Gly Gly Lys Glu His Leu Lys Ile Leu Ser Lys
                        105 110 115

ctt gct cgc tcc ttg gtg aag aag gat ttc atc aag gct ctg cag gaa 499
Leu Ala Arg Ser Leu Val Lys Lys Asp Phe Ile Lys Ala Leu Gln Glu
                        120 125 130

gcc acc acc gag cag gaa atc gtc gac gtt gtc gat gcc gtg ctc aac 547
Ala Thr Thr Glu Gln Glu Ile Val Asp Val Val Asp Ala Val Leu Asn
                        135 140 145

cca gca cca aaa acc acc gag cca gct gca gct ccg gct gcg gcg gcg 595
Pro Ala Pro Lys Thr Thr Glu Pro Ala Ala Ala Pro Ala Ala Ala Ala
                        150 155 160 165

gtt gct gag agt ggg gcg gcg tcg aca agc gtt act cgt atc gtg gca 643
```

Val	Ala	Glu	Ser	Gly	Ala	Ala	Ser	Thr	Ser	Val	Thr	Arg	Ile	Val	Ala		
				170					175					180			
atc	acc	gca	tgc	cca	acc	ggc	atc	gca	cac	acc	tac	atg	gct	gcg	gat	691	
Ile	Thr	Ala	Cys	Pro	Thr	Gly	Ile	Ala	His	Thr	Tyr	Met	Ala	Ala	Asp		
			185					190					195				
tcc	ctg	acg	caa	aac	gcg	gaa	ggc	cgc	gat	gat	gtg	gaa	ctc	gtt	gtg	739	
Ser	Leu	Thr	Gln	Asn	Ala	Glu	Gly	Arg	Asp	Asp	Val	Glu	Leu	Val	Val		
			200				205					210					
gag	act	cag	ggc	tct	tcc	gct	gtc	acc	cca	gtc	gat	ccg	aag	atc	atc	787	
Glu	Thr	Gln	Gly	Ser	Ser	Ala	Val	Thr	Pro	Val	Asp	Pro	Lys	Ile	Ile		
	215					220					225						
gaa	gct	gcc	gac	gcc	gtc	atc	ttc	gcc	acc	gac	gtg	gga	gtt	aaa	gac	835	
Glu	Ala	Ala	Asp	Ala	Val	Ile	Phe	Ala	Thr	Asp	Val	Gly	Val	Lys	Asp		
	230				235					240					245		
cgc	gag	cgt	ttc	gct	ggc	aag	cca	gtc	att	gaa	tcc	ggc	gtc	aag	cgc	883	
Arg	Glu	Arg	Phe	Ala	Gly	Lys	Pro	Val	Ile	Glu	Ser	Gly	Val	Lys	Arg		
			250					255						260			
gcg	atc	aat	gag	cca	gcc	aag	atg	atc	gac	gag	gcc	atc	gca	gcc	tcc	931	
Ala	Ile	Asn	Glu	Pro	Ala	Lys	Met	Ile	Asp	Glu	Ala	Ile	Ala	Ala	Ser		
			265					270					275				
aag	aac	cca	aac	gcc	cgc	aag	gtt	tcc	ggc	tcc	ggc	gtc	gcg	gca	tct	979	
Lys	Asn	Pro	Asn	Ala	Arg	Lys	Val	Ser	Gly	Ser	Gly	Val	Ala	Ala	Ser		
		280					285					290					
gct	gaa	acc	acc	ggc	gag	aag	ctc	ggc	tgg	ggc	aag	cgc	atc	cag	cag	1027	
Ala	Glu	Thr	Thr	Gly	Glu	Lys	Leu	Gly	Trp	Gly	Lys	Arg	Ile	Gln	Gln		
	295					300					305						
gca	gtc	atg	acc	ggc	gtg	tcc	tac	atg	gtt	cca	ttc	gta	gct	gcc	ggc	1075	
Ala	Val	Met	Thr	Gly	Val	Ser	Tyr	Met	Val	Pro	Phe	Val	Ala	Ala	Gly		
	310				315					320					325		
ggc	ctc	ctg	ttg	gct	ctc	ggc	ttc	gca	ttc	ggc	gga	tac	gac	atg	gcg	1123	
Gly	Leu	Leu	Leu	Ala	Leu	Gly	Phe	Ala	Phe	Gly	Gly	Tyr	Asp	Met	Ala		
				330				335						340			
aac	ggc	tgg	caa	gca	atc	gcc	acc	cag	ttc	tct	ctg	acc	aac	ctg	cca	1171	
Asn	Gly	Trp	Gln	Ala	Ile	Ala	Thr	Gln	Phe	Ser	Leu	Thr	Asn	Leu	Pro		
			345					350					355				
ggc	aac	acc	gtc	gat	gtt	gac	ggc	gtg	gcc	atg	acc	ttc	gag	cgt	tca	1219	
Gly	Asn	Thr	Val	Asp	Val	Asp	Gly	Val	Ala	Met	Thr	Phe	Glu	Arg	Ser		
		360					365					370					
ggc	ttc	ctg	ttg	tac	ttc	ggc	gca	gtc	ctg	ttc	gcc	acc	ggc	caa	gca	1267	
Gly	Phe	Leu	Leu	Tyr	Phe	Gly	Ala	Val	Leu	Phe	Ala	Thr	Gly	Gln	Ala		
	375					380					385						
gcc	atg	ggc	ttc	atc	gtg	gca	gcc	ctg	tct	ggc	tac	acc	gca	tac	gca	1315	
Ala	Met	Gly	Phe	Ile	Val	Ala	Ala	Leu	Ser	Gly	Tyr	Thr	Ala	Tyr	Ala		
	390				395				400						405		
ctt	gct	gga	cgc	cca	ggc	atc	gcg	ccg	ggc	ttc	gtc	ggc	ggc	gcc	atc	1363	
Leu	Ala	Gly	Arg	Pro	Gly	Ile	Ala	Pro	Gly	Phe	Val	Gly	Gly	Ala	Ile		



410				415				420				
tcc gtc acc atc ggc gct ggc ttc att ggt ggt ctg gtt acc ggt atc	1411											
Ser Val Thr Ile Gly Ala Gly Phe Ile Gly Gly Leu Val Thr Gly Ile												
425	430				435							
ttg gct ggt ctc att gcc ctg tgg att ggc tcc tgg aag gtg cca cgc	1459											
Leu Ala Gly Leu Ile Ala Leu Trp Ile Gly Ser Trp Lys Val Pro Arg												
440	445				450							
gtg gtg cag tca ctg atg cct gtg gtc atc atc ccg cta ctt acc tca	1507											
Val Val Gln Ser Leu Met Pro Val Val Ile Ile Ile Pro Leu Leu Thr Ser												
455	460				465							
gtg gtt gtt ggt ctc gtc atg tac ctc ctg ctg ggt cgc cca ctc gca	1555											
Val Val Val Gly Leu Val Met Tyr Leu Leu Leu Gly Arg Pro Leu Ala												
470	475				480				485			
tcc atc atg act ggt ttg cag gac tgg cta tcg tca atg tcc gga agc	1603											
Ser Ile Met Thr Gly Leu Gln Asp Trp Leu Ser Ser Met Ser Gly Ser												
490	495				500							
tcc gcc atc ttg ctg ggt atc atc ttg ggc ctc atg atg tgt ttc gac	1651											
Ser Ala Ile Leu Leu Gly Ile Ile Leu Gly Leu Met Met Cys Phe Asp												
505	510				515							
ctc ggc gga cca gta aac aag gca gcc tac ctc ttt ggt acc gca ggc	1699											
Leu Gly Gly Pro Val Asn Lys Ala Ala Tyr Leu Phe Gly Thr Ala Gly												
520	525				530							
ctg tct acc ggc gac caa gct tcc atg gaa atc atg gcc gcg atc atg	1747											
Leu Ser Thr Gly Asp Gln Ala Ser Met Glu Ile Met Ala Ala Ile Met												
535	540				545							
gca gct ggc atg gtc cca cca atc gcg ttg tcc att gct acc ctg ctg	1795											
Ala Ala Gly Met Val Pro Pro Ile Ala Leu Ser Ile Ala Thr Leu Leu												
550	555				560				565			
cgc aag aag ctg ttc acc cca gca gag caa gaa aac ggc aag tct tcc	1843											
Arg Lys Lys Leu Phe Thr Pro Ala Glu Gln Glu Asn Gly Lys Ser Ser												
570	575				580							
tgg ctg ctt ggc ctg gca ttc gtc tcc gaa ggt gcc atc cca ttc gcc	1891											
Trp Leu Leu Gly Leu Ala Phe Val Ser Glu Gly Ala Ile Pro Phe Ala												
585	590				595							
gca gct gac cca ttc cgt gtg atc cca gca atg atg gct ggc ggt gca	1939											
Ala Ala Asp Pro Phe Arg Val Ile Pro Ala Met Met Ala Gly Gly Ala												
600	605				610							
acc act ggt gca atc tcc atg gca ctg ggc gtc ggc tct cgg gct cca	1987											
Thr Thr Gly Ala Ile Ser Met Ala Leu Gly Val Gly Ser Arg Ala Pro												
615	620				625							
cac ggc ggt atc ttc gtg gtc tgg gca atc gaa cca tgg tgg ggc tgg	2035											
His Gly Gly Ile Phe Val Val Trp Ala Ile Glu Pro Trp Trp Gly Trp												
630	635				640				645			
ctc atc gca ctt gca gca ggc acc atc gtg tcc acc atc gtt gtc atc	2083											
Leu Ile Ala Leu Ala Ala Gly Thr Ile Val Ser Thr Ile Val Val Ile												
650	655				660							

gca ctg aag cag ttc tgg cca aac aag gcc gtc gct gca gaa gtc gcg 2131  
 Ala Leu Lys Gln Phe Trp Pro Asn Lys Ala Val Ala Ala Glu Val Ala  
           665                                670                                675

aag caa gaa gca caa caa gca gct gta aac gca taatcggacc ttgacccgat 2184  
 Lys Gln Glu Ala Gln Gln Ala Ala Val Asn Ala  
           680                                685

gtc 2187

<210> 8

<211> 688

<212> PRT

<213> Corynebacterium glutamicum

<400> 8

Met Asn Ser Val Asn Asn Ser Ser Leu Val Arg Leu Asp Val Asp Phe  
   1                                5                                10                                15

Gly Asp Ser Thr Thr Asp Val Ile Asn Asn Leu Ala Thr Val Ile Phe  
           20                                25                                30

Asp Ala Gly Arg Ala Ser Ser Ala Asp Ala Leu Ala Lys Asp Ala Leu  
           35                                40                                45

Asp Arg Glu Ala Lys Ser Gly Thr Gly Val Pro Gly Gln Val Ala Ile  
           50                                55                                60

Pro His Cys Arg Ser Glu Ala Val Ser Val Pro Thr Leu Gly Phe Ala  
           65                                70                                75                                80

Arg Leu Ser Lys Gly Val Asp Phe Ser Gly Pro Asp Gly Asp Ala Asn  
           85                                90                                95

Leu Val Phe Leu Ile Ala Ala Pro Ala Gly Gly Gly Lys Glu His Leu  
           100                                105                                110

Lys Ile Leu Ser Lys Leu Ala Arg Ser Leu Val Lys Lys Asp Phe Ile  
           115                                120                                125

Lys Ala Leu Gln Glu Ala Thr Thr Glu Gln Glu Ile Val Asp Val Val  
           130                                135                                140

Asp Ala Val Leu Asn Pro Ala Pro Lys Thr Thr Glu Pro Ala Ala Ala  
           145                                150                                155                                160

Pro Ala Ala Ala Ala Val Ala Glu Ser Gly Ala Ala Ser Thr Ser Val  
           165                                170                                175

Thr Arg Ile Val Ala Ile Thr Ala Cys Pro Thr Gly Ile Ala His Thr  
           180                                185                                190

Tyr Met Ala Ala Asp Ser Leu Thr Gln Asn Ala Glu Gly Arg Asp Asp  
           195                                200                                205

Val Glu Leu Val Val Glu Thr Gln Gly Ser Ser Ala Val Thr Pro Val  
           210                                215                                220

Asp Pro Lys Ile Ile Glu Ala Ala Asp Ala Val Ile Phe Ala Thr Asp

225						230						235						240
Val Gly Val Lys Asp	Arg Glu Arg Phe	Ala Gly Lys Pro Val	Ile Glu															
		245						250								255		
Ser Gly Val Lys Arg	Ala Ile Asn Glu	Pro Ala Lys Met	Ile Asp Glu															
		260						265								270		
Ala Ile Ala Ala Ser	Lys Asn Pro Asn	Ala Arg Lys Val	Ser Gly Ser															
		275						280								285		
Gly Val Ala Ala Ser	Ala Glu Thr Thr	Gly Glu Lys Leu	Gly Trp Gly															
		290						295								300		
Lys Arg Ile Gln Gln	Ala Val Met Thr	Gly Val Ser Tyr	Met Val Pro															
		305						310								315		
Phe Val Ala Ala Gly	Gly Leu Leu Leu	Ala Leu Gly Phe	Ala Phe Gly															
								325								330		
Gly Tyr Asp Met Ala	Asn Gly Trp Gln	Ala Ile Ala Thr	Gln Phe Ser															
		340						345								350		
Leu Thr Asn Leu Pro	Gly Asn Thr Val	Asp Val Asp Gly	Val Ala Met															
		355						360								365		
Thr Phe Glu Arg Ser	Gly Phe Leu Leu	Tyr Phe Gly Ala	Val Leu Phe															
		370						375								380		
Ala Thr Gly Gln Ala	Ala Met Gly Phe	Ile Val Ala Ala	Leu Ser Gly															
		385						390								395		
Tyr Thr Ala Tyr Ala	Leu Ala Gly Arg	Pro Gly Ile Ala	Pro Gly Phe															
								405								410		
Val Gly Gly Ala Ile	Ser Val Thr Ile	Gly Ala Gly Phe	Ile Gly Gly															
								420								425		
Leu Val Thr Gly Ile	Leu Ala Gly Leu	Ile Ala Leu Trp	Ile Gly Ser															
		435						440								445		
Trp Lys Val Pro Arg	Val Val Gln Ser	Leu Met Pro Val	Val Ile Ile															
								450								455		
Pro Leu Leu Thr Ser	Val Val Val Gly	Leu Val Met Tyr	Leu Leu Leu															
								465								470		
Gly Arg Pro Leu Ala	Ser Ile Met Thr	Gly Leu Gln Asp	Trp Leu Ser															
								485								490		
Ser Met Ser Gly Ser	Ser Ala Ile Leu	Leu Leu Gly Ile	Ile Leu Gly															
								500								505		
Met Met Cys Phe Asp	Leu Gly Gly Pro	Val Asn Lys Ala	Ala Tyr Leu															
								515								520		
Phe Gly Thr Ala Gly	Leu Ser Thr Gly	Asp Gln Ala Ser	Met Glu Ile															
								530								535		
Met Ala Ala Ile Met	Ala Ala Gly Met	Val Pro Pro Ile	Ala Leu Ser															
								545								550		

Ile Ala Thr Leu Leu Arg Lys Lys Leu Phe Thr Pro Ala Glu Gln Glu  
565 570 575

Asn Gly Lys Ser Ser Trp Leu Leu Gly Leu Ala Phe Val Ser Glu Gly  
580 585 590

Ala Ile Pro Phe Ala Ala Ala Asp Pro Phe Arg Val Ile Pro Ala Met  
595 600 605

Met Ala Gly Gly Ala Thr Thr Gly Ala Ile Ser Met Ala Leu Gly Val  
610 615 620

Gly Ser Arg Ala Pro His Gly Gly Ile Phe Val Val Trp Ala Ile Glu  
625 630 635 640

Pro Trp Trp Gly Trp Leu Ile Ala Leu Ala Ala Gly Thr Ile Val Ser  
645 650 655

Thr Ile Val Val Ile Ala Leu Lys Gln Phe Trp Pro Asn Lys Ala Val  
660 665 670

Ala Ala Glu Val Ala Lys Gln Glu Ala Gln Gln Ala Ala Val Asn Ala  
675 680 685

<210> 9

<211> 464

<212> DNA

<213> Corynebacterium glutamicum

<220>

<221> CDS

<222> (1)..(441)

<223> FRXA01299

<400> 9

atg gaa atc atg gcc gcg atc atg gca gct ggc atg gtc cca cca atc 48  
Met Glu Ile Met Ala Ala Ile Met Ala Ala Gly Met Val Pro Pro Ile  
1 5 10 15

gcg ttg tcc att gct acc ctg ctg cgc aag aag ctg ttc acc cca gca 96  
Ala Leu Ser Ile Ala Thr Leu Leu Arg Lys Lys Leu Phe Thr Pro Ala  
20 25 30

gag caa gaa aac ggc aag tct tcc tgg ctg ctt ggc ctg gca ttc gtc 144  
Glu Gln Glu Asn Gly Lys Ser Ser Trp Leu Leu Gly Leu Ala Phe Val  
35 40 45

tcc gaa ggt gcc atc cca ttc gcc gca gct gac cca ttc cgt gtg atc 192  
Ser Glu Gly Ala Ile Pro Phe Ala Ala Ala Asp Pro Phe Arg Val Ile  
50 55 60

cca gca atg atg gct ggc ggt gca acc act ggt gca atc tcc atg gca 240  
Pro Ala Met Met Ala Gly Gly Ala Thr Thr Gly Ala Ile Ser Met Ala  
65 70 75 80

ctg ggc gtc ggc tct cgg gct cca cac ggc ggt atc ttc gtg gtc tgg 288

```
<210> 10
<211> 147
<212> PRT
<213> Corynebacterium glutamicum
```

```
<210> 11
<211> 580
<212> DNA
<213> Corynebacterium glutamicum
```

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (101)..(580)

&lt;223&gt; FRXA01883

&lt;400&gt; 11

```

cgactgcggc gtctcttctt ggcactacca ttctctgtcc tgaccaactc gccacagctg 60

gtgcaacggt cacccaagtc aaaggattga aagaatcagc atg aat agc gta aat 115
                               Met Asn Ser Val Asn
                               1      5

aat tcc tcg ctt gtc cgg ctg gat gtc gat ttc ggc gac tcc acc acg 163
Asn Ser Ser Leu Val Arg Leu Asp Val Asp Phe Gly Asp Ser Thr Thr
                10                15                20

gat gtc atc aac aac ctt gcc act gtt att ttc gac gct ggc cga gct 211
Asp Val Ile Asn Asn Leu Ala Thr Val Ile Phe Asp Ala Gly Arg Ala
                25                30                35

tcc tcc gcc gac gcc ctt gcc aaa gac gcg ctg gat cgt gaa gca aag 259
Ser Ser Ala Asp Ala Leu Ala Lys Asp Ala Leu Asp Arg Glu Ala Lys
                40                45                50

tcc ggc acc ggc gtt cct ggt caa gtt gct atc ccc cac tgc cgt tcc 307
Ser Gly Thr Gly Val Pro Gly Gln Val Ala Ile Pro His Cys Arg Ser
                55                60                65

gaa gcc gta tct gtc cct acc ttg ggc ttt gct cgc ctg agc aag ggt 355
Glu Ala Val Ser Val Pro Thr Leu Gly Phe Ala Arg Leu Ser Lys Gly
                70                75                80                85

gtg gac ttc agc gga cct gat ggc gat gcc aac ttg gtg ttc ctc att 403
Val Asp Phe Ser Gly Pro Asp Gly Asp Ala Asn Leu Val Phe Leu Ile
                90                95                100

gca gca cct gct ggc ggc ggc aaa gag cac ctg aag atc ctg tcc aag 451
Ala Ala Pro Ala Gly Gly Gly Lys Glu His Leu Lys Ile Leu Ser Lys
                105                110                115

ctt gct cgc tcc ttg gtg aag aag gat ttc atc aag gct ctg cag gaa 499
Leu Ala Arg Ser Leu Val Lys Lys Asp Phe Ile Lys Ala Leu Gln Glu
                120                125                130

gcc acc acc gag cag gaa atc gtc gac gtt gtc gat gcc gtg ctc aac 547
Ala Thr Thr Glu Gln Glu Ile Val Asp Val Val Asp Ala Val Leu Asn
                135                140                145

cca gca cca aaa aac cac cga gcc agc tgc agc 580
Pro Ala Pro Lys Asn His Arg Ala Ser Cys Ser
150                155                160

```

&lt;210&gt; 12

&lt;211&gt; 160

&lt;212&gt; PRT

&lt;213&gt; Corynebacterium glutamicum

&lt;400&gt; 12

Met Asn Ser Val Asn Asn Ser Ser Leu Val Arg Leu Asp Val Asp Phe

1	5	10	15
Gly Asp Ser Thr Thr Asp Val Ile Asn Asn Leu Ala Thr Val Ile Phe	20	25	30
Asp Ala Gly Arg Ala Ser Ser Ala Asp Ala Leu Ala Lys Asp Ala Leu	35	40	45
Asp Arg Glu Ala Lys Ser Gly Thr Gly Val Pro Gly Gln Val Ala Ile	50	55	60
Pro His Cys Arg Ser Glu Ala Val Ser Val Pro Thr Leu Gly Phe Ala	65	70	75
Arg Leu Ser Lys Gly Val Asp Phe Ser Gly Pro Asp Gly Asp Ala Asn	85	90	95
Leu Val Phe Leu Ile Ala Ala Pro Ala Gly Gly Gly Lys Glu His Leu	100	105	110
Lys Ile Leu Ser Lys Leu Ala Arg Ser Leu Val Lys Lys Asp Phe Ile	115	120	125
Lys Ala Leu Gln Glu Ala Thr Thr Glu Gln Glu Ile Val Asp Val Val	130	135	140
Asp Ala Val Leu Asn Pro Ala Pro Lys Asn His Arg Ala Ser Cys Ser	145	150	155
			160

<210> 13  
 <211> 631  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (77)..(631)  
 <223> FRXA01889

<400> 13  
 accgagccag ctgcagctcc ggctgcggcg gccggttggt aagagtgggg cggcgtcgac 60  
 aagcgttact cgtatcgtg gca atc acc gca tgc cca acc ggt atc gca cac 112  
                   Val Ala Ile Thr Ala Cys Pro Thr Gly Ile Ala His  
                   1                  5                  10  
 acc tac atg gct gcg gat tcc ctg acg caa aac gcg gaa ggc cgc gat 160  
 Thr Tyr Met Ala Ala Asp Ser Leu Thr Gln Asn Ala Glu Gly Arg Asp  
                   15                  20                  25  
 gat gtg gaa ctc gtt gtg gag act cag ggc tct tcc gct gtc acc cca 208  
 Asp Val Glu Leu Val Val Glu Thr Gln Gly Ser Ser Ala Val Thr Pro  
                   30                  35                  40  
 gtc gat ccg aag atc atc gaa gct gcc gac gcc gtc atc ttc gcc acc 256  
 Val Asp Pro Lys Ile Ile Glu Ala Ala Asp Ala Val Ile Phe Ala Thr  
                   45                  50                  55                  60

gac gtg gga gtt aaa gac cgc gag cgt ttc gct ggc aag cca gtc att 304  
 Asp Val Gly Val Lys Asp Arg Glu Arg Phe Ala Gly Lys Pro Val Ile  
 65 70 75

gaa tcc ggc gtc aag cgc gcg atc aat gag cca gcc aag atg atc gac 352  
 Glu Ser Gly Val Lys Arg Ala Ile Asn Glu Pro Ala Lys Met Ile Asp  
 80 85 90

gag gcc atc gca gcc tcc aag aac cca aac gcc cgc aag gtt tcc ggt 400  
 Glu Ala Ile Ala Ala Ser Lys Asn Pro Asn Ala Arg Lys Val Ser Gly  
 95 100 105

tcc ggt gtc gcg gca tct gct gaa acc acc ggc gag aag ctc ggc tgg 448  
 Ser Gly Val Ala Ala Ser Ala Glu Thr Thr Gly Glu Lys Leu Gly Trp  
 110 115 120

ggc aag cgc atc cag cag gca gtc atg acc ggc gtg tcc tac atg gtt 496  
 Gly Lys Arg Ile Gln Gln Ala Val Met Thr Gly Val Ser Tyr Met Val  
 125 130 135 140

cca ttc gta gct gcc ggc ggc ctc ctg ttg gct ctc ggc ttc gca ttc 544  
 Pro Phe Val Ala Ala Gly Gly Leu Leu Leu Ala Leu Gly Phe Ala Phe  
 145 150 155

ggt gga tac gac atg gcg aac ggc tgg caa gca atc gcc acc cag ttc 592  
 Gly Gly Tyr Asp Met Ala Asn Gly Trp Gln Ala Ile Ala Thr Gln Phe  
 160 165 170

tct ctg acc aac ctg cca ggc aac acc gtc gat gtt gac 631  
 Ser Leu Thr Asn Leu Pro Gly Asn Thr Val Asp Val Asp  
 175 180 185

&lt;210&gt; 14

&lt;211&gt; 185

&lt;212&gt; PRT

&lt;213&gt; Corynebacterium glutamicum

&lt;400&gt; 14

Val Ala Ile Thr Ala Cys Pro Thr Gly Ile Ala His Thr Tyr Met Ala  
 1 5 10 15

Ala Asp Ser Leu Thr Gln Asn Ala Glu Gly Arg Asp Asp Val Glu Leu  
 20 25 30

Val Val Glu Thr Gln Gly Ser Ser Ala Val Thr Pro Val Asp Pro Lys  
 35 40 45

Ile Ile Glu Ala Ala Asp Ala Val Ile Phe Ala Thr Asp Val Gly Val  
 50 55 60

Lys Asp Arg Glu Arg Phe Ala Gly Lys Pro Val Ile Glu Ser Gly Val  
 65 70 75 80

Lys Arg Ala Ile Asn Glu Pro Ala Lys Met Ile Asp Glu Ala Ile Ala  
 85 90 95

Ala Ser Lys Asn Pro Asn Ala Arg Lys Val Ser Gly Ser Gly Val Ala  
 100 105 110



Ala Ser Ala Glu Thr Thr Gly Glu Lys Leu Gly Trp Gly Lys Arg Ile  
 115 120 125

Gln Gln Ala Val Met Thr Gly Val Ser Tyr Met Val Pro Phe Val Ala  
 130 135 140

Ala Gly Gly Leu Leu Leu Ala Leu Gly Phe Ala Phe Gly Gly Tyr Asp  
 145 150 155 160

Met Ala Asn Gly Trp Gln Ala Ile Ala Thr Gln Phe Ser Leu Thr Asn  
 165 170 175

Leu Pro Gly Asn Thr Val Asp Val Asp  
 180 185

<210> 15  
 <211> 416  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (1)..(393)  
 <223> RXA00951

<400> 15  
 atc caa gca atc tta gag aag gca gca gcg ccg gcg aag cag aag gct 48  
 Ile Gln Ala Ile Leu Glu Lys Ala Ala Ala Pro Ala Lys Gln Lys Ala  
 1 5 10 15

cct gct gtg gct cct gct gta aca ccc act gac gct cct gca gcc tca 96  
 Pro Ala Val Ala Pro Ala Val Thr Pro Thr Asp Ala Pro Ala Ala Ser  
 20 25 30

gtc caa tcc aaa acc cac gac aag atc ctc acc gtc tgt ggc aac ggc 144  
 Val Gln Ser Lys Thr His Asp Lys Ile Leu Thr Val Cys Gly Asn Gly  
 35 40 45

ttg ggt acc tcc ctc ttc ctc aaa aac acc ctt gag caa gtt ttc gac 192  
 Leu Gly Thr Ser Leu Phe Leu Lys Asn Thr Leu Glu Gln Val Phe Asp  
 50 55 60

acc tgg ggt tgg ggt cca tac atg acg gtg gag gca acc gac act atc 240  
 Thr Trp Gly Trp Gly Pro Tyr Met Thr Val Glu Ala Thr Asp Thr Ile  
 65 70 75 80

tcc gcc aag ggc aaa gcc aag gaa gct gat ctc atc atg acc tct ggt 288  
 Ser Ala Lys Gly Lys Ala Lys Glu Ala Asp Leu Ile Met Thr Ser Gly  
 85 90 95

gaa atc gcc cgc acg ttg ggt gat gtt gga atc ccg gtt cac gtg atc 336  
 Glu Ile Ala Arg Thr Leu Gly Asp Val Gly Ile Pro Val His Val Ile  
 100 105 110

aat gac ttc acg agc acc gat gaa atc gat gct gcg ctt cgt gaa cgc 384  
 Asn Asp Phe Thr Ser Thr Asp Glu Ile Asp Ala Ala Leu Arg Glu Arg  
 115 120 125

tac gac atc taactacttt aaaaggacga aaa 416  
 Tyr Asp Ile

130

<210> 16  
 <211> 131  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 16  
 Ile Gln Ala Ile Leu Glu Lys Ala Ala Ala Pro Ala Lys Gln Lys Ala  
 1 5 10 15  
 Pro Ala Val Ala Pro Ala Val Thr Pro Thr Asp Ala Pro Ala Ala Ser  
 20 25 30  
 Val Gln Ser Lys Thr His Asp Lys Ile Leu Thr Val Cys Gly Asn Gly  
 35 40 45  
 Leu Gly Thr Ser Leu Phe Leu Lys Asn Thr Leu Glu Gln Val Phe Asp  
 50 55 60  
 Thr Trp Gly Trp Gly Pro Tyr Met Thr Val Glu Ala Thr Asp Thr Ile  
 65 70 75 80  
 Ser Ala Lys Gly Lys Ala Lys Glu Ala Asp Leu Ile Met Thr Ser Gly  
 85 90 95  
 Glu Ile Ala Arg Thr Leu Gly Asp Val Gly Ile Pro Val His Val Ile  
 100 105 110  
 Asn Asp Phe Thr Ser Thr Asp Glu Ile Asp Ala Ala Leu Arg Glu Arg  
 115 120 125  
 Tyr Asp Ile  
 130

<210> 17  
 <211> 1827  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (101)..(1804)  
 <223> RXN01244

<400> 17  
 gatatgtggt tgtttgtcaa tatccaaatg tttgaatagt tgcacaactg ttggttttgt 60  
 ggtgatcttg aggaaattaa ctcaatgatt gtgaggatgg gtg gct act gtg gct 115  
 Val Ala Thr Val Ala  
 1 5  
 gat gtg aat caa gac act gta ctg aag ggc acc ggc gtt gtc ggt gga 163  
 Asp Val Asn Gln Asp Thr Val Leu Lys Gly Thr Gly Val Val Gly Gly  
 10 15 20  
 gtc cgt tat gca agc gcg gtg tgg att acc cca cgc ccc gaa cta ccc 211  
 Val Arg Tyr Ala Ser Ala Val Trp Ile Thr Pro Arg Pro Glu Leu Pro  
 25 30 35

caa gca ggc gaa gtc gtc gcc gaa gaa aac cgt gaa gca gag cag gag	259
Gln Ala Gly Glu Val Val Ala Glu Glu Asn Arg Glu Ala Glu Gln Glu	
40 45 50	
cgt ttc gac gcc gct gca gcc aca gtc tct tct cgt ttg ctt gag cgc	307
Arg Phe Asp Ala Ala Ala Thr Val Ser Ser Arg Leu Leu Glu Arg	
55 60 65	
tcc gaa gct gct gaa gga cca gca gct gag gtg ctt aaa gct act gct	355
Ser Glu Ala Ala Glu Gly Pro Ala Ala Glu Val Leu Lys Ala Thr Ala	
70 75 80 85	
ggc atg gtc aat gac cgt ggc tgg cgt aag gct gtc atc aag ggt gtc	403
Gly Met Val Asn Asp Arg Gly Trp Arg Lys Ala Val Ile Lys Gly Val	
90 95 100	
aag ggt ggt cac cct gcg gaa tac gcc gtg gtt gca gca aca acc aag	451
Lys Gly Gly His Pro Ala Glu Tyr Ala Val Val Ala Ala Thr Thr Lys	
105 110 115	
ttc atc tcc atg ttc gaa gcc gca ggc ggc ctg atc gcg gag cgc acc	499
Phe Ile Ser Met Phe Glu Ala Ala Gly Gly Leu Ile Ala Glu Arg Thr	
120 125 130	
aca gac ttg cgc gac atc cgc gac cgc gtc atc gca gaa ctt cgt ggc	547
Thr Asp Leu Arg Asp Ile Arg Asp Arg Val Ile Ala Glu Leu Arg Gly	
135 140 145	
gat gaa gag cca ggt ctg cca gct gtt tcc gga cag gtc att ctc ttt	595
Asp Glu Glu Pro Gly Leu Pro Ala Val Ser Gly Gln Val Ile Leu Phe	
150 155 160 165	
gca gat gac ctc tcc cca gca gac acc gcg gca cta gac aca gat ctc	643
Ala Asp Asp Leu Ser Pro Ala Asp Thr Ala Ala Leu Asp Thr Asp Leu	
170 175 180	
ttt gtg gga ctt gtc act gag ctg ggt ggc cca acg agc cac acc gcg	691
Phe Val Gly Leu Val Thr Glu Leu Gly Gly Pro Thr Ser His Thr Ala	
185 190 195	
atc atc gca cgc cag ctc aac gtg cct tgc atc gtc gca tcc ggc gcc	739
Ile Ile Ala Arg Gln Leu Asn Val Pro Cys Ile Val Ala Ser Gly Ala	
200 205 210	
ggc atc aag gac atc aag tcc ggc gaa aag gtg ctt atc gac ggc agc	787
Gly Ile Lys Asp Ile Lys Ser Gly Glu Lys Val Leu Ile Asp Gly Ser	
215 220 225	
ctc ggc acc att gac cgc aac gcg gac gaa gct gaa gca acc aag ctc	835
Leu Gly Thr Ile Asp Arg Asn Ala Asp Glu Ala Glu Ala Thr Lys Leu	
230 235 240 245	
gtc tcc gag tcc ctc gag cgc gct gct cgc atc gcc gag tgg aag ggt	883
Val Ser Glu Ser Leu Glu Arg Ala Ala Arg Ile Ala Glu Trp Lys Gly	
250 255 260	
cct gca caa acc aag gac ggc tac cgc gtt cag ctg ttg gcc aac gtc	931
Pro Ala Gln Thr Lys Asp Gly Tyr Arg Val Gln Leu Leu Ala Asn Val	
265 270 275	

caa gac ggc aac tct gca cag cag gct gca cag acc gaa gca gaa ggc	979
Gln Asp Gly Asn Ser Ala Gln Gln Ala Ala Gln Thr Glu Ala Glu Gly	
280 285 290	
atc ggc ctg ttc cgc acc gaa ctg tgc ttc ctt tcc gcc acc gaa gag	1027
Ile Gly Leu Phe Arg Thr Glu Leu Cys Phe Leu Ser Ala Thr Glu Glu	
295 300 305	
cca agc gtt gat gag cag gct gcg gtc tac tca aag gtg ctt gaa gca	1075
Pro Ser Val Asp Glu Gln Ala Ala Val Tyr Ser Lys Val Leu Glu Ala	
310 315 320 325	
ttc cca gag tcc aag gtc gtt gtc cgc tcc ctc gac gca ggt tct gac	1123
Phe Pro Glu Ser Lys Val Val Val Arg Ser Leu Asp Ala Gly Ser Asp	
330 335 340	
aag cca gtt cca ttc gca tcg atg gct gat gag atg aac cca gca ctg	1171
Lys Pro Val Pro Phe Ala Ser Met Ala Asp Glu Met Asn Pro Ala Leu	
345 350 355	
ggt gtt cgt ggc ctg cgt atc gca cgt gga cag gtt gat ctg ctg act	1219
Gly Val Arg Gly Leu Arg Ile Ala Arg Gly Gln Val Asp Leu Leu Thr	
360 365 370	
cgc cag ctc gac gca att gcg aag gcc agc gaa gaa ctc ggc cgt ggc	1267
Arg Gln Leu Asp Ala Ile Ala Lys Ala Ser Glu Glu Leu Gly Arg Gly	
375 380 385	
gac gac gcc cca acc tgg gtt atg gct cca atg gtg gct acc gct tat	1315
Asp Asp Ala Pro Thr Trp Val Met Ala Pro Met Val Ala Thr Ala Tyr	
390 395 400 405	
gaa gca aag tgg ttt gct gac atg tgc cgt gag cgt ggc cta atc gcc	1363
Glu Ala Lys Trp Phe Ala Asp Met Cys Arg Glu Arg Gly Leu Ile Ala	
410 415 420	
ggc gcc atg atc gaa gtt cca gca gca tcc ctg atg gca gac aag atc	1411
Gly Ala Met Ile Glu Val Pro Ala Ala Ser Leu Met Ala Asp Lys Ile	
425 430 435	
atg cct cac ctg gac ttt gtt tcc atc ggt acc aac gac ctg acc cag	1459
Met Pro His Leu Asp Phe Val Ser Ile Gly Thr Asn Asp Leu Thr Gln	
440 445 450	
tac acc atg gca gcg gac cgc atg tct cct gag ctt gcc tac ctg acc	1507
Tyr Thr Met Ala Ala Asp Arg Met Ser Pro Glu Leu Ala Tyr Leu Thr	
455 460 465	
gat cct tgg cag cca gca gtc ctg cgc ctg atc aag cac acc tgt gac	1555
Asp Pro Trp Gln Pro Ala Val Leu Arg Leu Ile Lys His Thr Cys Asp	
470 475 480 485	
gaa ggt gct cgc ttt aac acc ccg gtc ggt gtt tgt ggt gaa gca gca	1603
Glu Gly Ala Arg Phe Asn Thr Pro Val Gly Val Cys Gly Glu Ala Ala	
490 495 500	
gca gac cca ctg ttg gca act gtc ctc acc ggt ctt ggc gtg aac tcc	1651
Ala Asp Pro Leu Leu Ala Thr Val Leu Thr Gly Leu Gly Val Asn Ser	
505 510 515	
ctg tcc gca gca tcc act gct ctc gca gca gtc ggt gca aag ctg tca	1699

Leu Ser Ala Ala Ser Thr Ala Leu Ala Ala Val Gly Ala Lys Leu Ser  
 520 525 530  
 gag gtc acc ctg gaa acc tgt aag aag gca gca gaa gca gca ctt gac 1747  
 Glu Val Thr Leu Glu Thr Cys Lys Lys Ala Ala Glu Ala Ala Leu Asp  
 535 540 545  
 gct gaa ggt gca act gaa gca cgc gat gct gta cgc gca gtg atc gac 1795  
 Ala Glu Gly Ala Thr Glu Ala Arg Asp Ala Val Arg Ala Val Ile Asp  
 550 555 560 565  
 gca gca gtc taaaccactg ttgagctaaa aag 1827  
 Ala Ala Val

<210> 18  
 <211> 568  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 18  
 Val Ala Thr Val Ala Asp Val Asn Gln Asp Thr Val Leu Lys Gly Thr  
 1 5 10 15  
 Gly Val Val Gly Gly Val Arg Tyr Ala Ser Ala Val Trp Ile Thr Pro  
 20 25 30  
 Arg Pro Glu Leu Pro Gln Ala Gly Glu Val Val Ala Glu Glu Asn Arg  
 35 40 45  
 Glu Ala Glu Gln Glu Arg Phe Asp Ala Ala Ala Thr Val Ser Ser  
 50 55 60  
 Arg Leu Leu Glu Arg Ser Glu Ala Ala Glu Gly Pro Ala Ala Glu Val  
 65 70 75 80  
 Leu Lys Ala Thr Ala Gly Met Val Asn Asp Arg Gly Trp Arg Lys Ala  
 85 90 95  
 Val Ile Lys Gly Val Lys Gly Gly His Pro Ala Glu Tyr Ala Val Val  
 100 105 110  
 Ala Ala Thr Thr Lys Phe Ile Ser Met Phe Glu Ala Ala Gly Gly Leu  
 115 120 125  
 Ile Ala Glu Arg Thr Thr Asp Leu Arg Asp Ile Arg Asp Arg Val Ile  
 130 135 140  
 Ala Glu Leu Arg Gly Asp Glu Glu Pro Gly Leu Pro Ala Val Ser Gly  
 145 150 155 160  
 Gln Val Ile Leu Phe Ala Asp Asp Leu Ser Pro Ala Asp Thr Ala Ala  
 165 170 175  
 Leu Asp Thr Asp Leu Phe Val Gly Leu Val Thr Glu Leu Gly Gly Pro  
 180 185 190  
 Thr Ser His Thr Ala Ile Ile Ala Arg Gln Leu Asn Val Pro Cys Ile  
 195 200 205

Val Ala Ser Gly Ala Gly Ile Lys Asp Ile Lys Ser Gly Glu Lys Val  
 210 215 220  
 Leu Ile Asp Gly Ser Leu Gly Thr Ile Asp Arg Asn Ala Asp Glu Ala  
 225 230 235 240  
 Glu Ala Thr Lys Leu Val Ser Glu Ser Leu Glu Arg Ala Ala Arg Ile  
 245 250 255  
 Ala Glu Trp Lys Gly Pro Ala Gln Thr Lys Asp Gly Tyr Arg Val Gln  
 260 265 270  
 Leu Leu Ala Asn Val Gln Asp Gly Asn Ser Ala Gln Gln Ala Ala Gln  
 275 280 285  
 Thr Glu Ala Glu Gly Ile Gly Leu Phe Arg Thr Glu Leu Cys Phe Leu  
 290 295 300  
 Ser Ala Thr Glu Glu Pro Ser Val Asp Glu Gln Ala Ala Val Tyr Ser  
 305 310 315 320  
 Lys Val Leu Glu Ala Phe Pro Glu Ser Lys Val Val Val Arg Ser Leu  
 325 330 335  
 Asp Ala Gly Ser Asp Lys Pro Val Pro Phe Ala Ser Met Ala Asp Glu  
 340 345 350  
 Met Asn Pro Ala Leu Gly Val Arg Gly Leu Arg Ile Ala Arg Gly Gln  
 355 360 365  
 Val Asp Leu Leu Thr Arg Gln Leu Asp Ala Ile Ala Lys Ala Ser Glu  
 370 375 380  
 Glu Leu Gly Arg Gly Asp Asp Ala Pro Thr Trp Val Met Ala Pro Met  
 385 390 395 400  
 Val Ala Thr Ala Tyr Glu Ala Lys Trp Phe Ala Asp Met Cys Arg Glu  
 405 410 415  
 Arg Gly Leu Ile Ala Gly Ala Met Ile Glu Val Pro Ala Ala Ser Leu  
 420 425 430  
 Met Ala Asp Lys Ile Met Pro His Leu Asp Phe Val Ser Ile Gly Thr  
 435 440 445  
 Asn Asp Leu Thr Gln Tyr Thr Met Ala Ala Asp Arg Met Ser Pro Glu  
 450 455 460  
 Leu Ala Tyr Leu Thr Asp Pro Trp Gln Pro Ala Val Leu Arg Leu Ile  
 465 470 475 480  
 Lys His Thr Cys Asp Glu Gly Ala Arg Phe Asn Thr Pro Val Gly Val  
 485 490 495  
 Cys Gly Glu Ala Ala Ala Asp Pro Leu Leu Ala Thr Val Leu Thr Gly  
 500 505 510  
 Leu Gly Val Asn Ser Leu Ser Ala Ala Ser Thr Ala Leu Ala Ala Val  
 515 520 525  
 Gly Ala Lys Leu Ser Glu Val Thr Leu Glu Thr Cys Lys Lys Ala Ala

530

535

540

Glu Ala Ala Leu Asp Ala Glu Gly Ala Thr Glu Ala Arg Asp Ala Val  
 545 550 555 560

Arg Ala Val Ile Asp Ala Ala Val  
 565

&lt;210&gt; 19

&lt;211&gt; 1629

&lt;212&gt; DNA

<213> *Corynebacterium glutamicum*

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (98)..(1606)

&lt;223&gt; FRXA01244

&lt;400&gt; 19

agatgtcgat ttctcgagga agaagttaac gccgaagaaa accgtgaatc agagcaggag 60

cgcttcgacg ccgctgcagc cacagtctct tcttcgtttg ctt gag cgc tcc gaa 115  
 Leu Leu Glu Arg Ser Glu  
 1 5

gct gct gaa gga cca gca gct gag gtg ctt aaa gct act gct ggc atg 163  
 Ala Ala Glu Gly Pro Ala Ala Glu Val Leu Lys Ala Thr Ala Gly Met  
 10 15 20

gtc aat gac cgt ggc tgg cgt aag gct gtc atc aag ggt gtc aag ggt 211  
 Val Asn Asp Arg Gly Trp Arg Lys Ala Val Ile Lys Gly Val Lys Gly  
 25 30 35

ggt cac cct gcg gaa tac gcc gtg gtt gca gca aca acc aag ttc atc 259  
 Gly His Pro Ala Glu Tyr Ala Val Val Ala Ala Thr Thr Lys Phe Ile  
 40 45 50

tcc atg ttc gaa gcc gca ggc ggc ctg atc gcg gag cgc acc aca gac 307  
 Ser Met Phe Glu Ala Ala Gly Gly Leu Ile Ala Glu Arg Thr Thr Asp  
 55 60 65 70

ttg cgc gac atc cgc gac cgc gtc atc gca gaa ctt cgt ggc gat gaa 355  
 Leu Arg Asp Ile Arg Asp Arg Val Ile Ala Glu Leu Arg Gly Asp Glu  
 75 80 85

gag cca ggt ctg cca gct gtt tcc gga cag gtc att ctc ttt gca gat 403  
 Glu Pro Gly Leu Pro Ala Val Ser Gly Gln Val Ile Leu Phe Ala Asp  
 90 95 100

gac ctc tcc cca gca gac acc gcg gca cta gac aca gat ctc ttt gtg 451  
 Asp Leu Ser Pro Ala Asp Thr Ala Ala Leu Asp Thr Asp Leu Phe Val  
 105 110 115

gga ctt gtc act gag ctg ggt ggc cca acg agc cac acc gcg atc atc 499  
 Gly Leu Val Thr Glu Leu Gly Gly Pro Thr Ser His Thr Ala Ile Ile  
 120 125 130

gca cgc cag ctc aac gtg cct tgc atc gtc gca tcc ggc gcc ggc atc 547  
 Ala Arg Gln Leu Asn Val Pro Cys Ile Val Ala Ser Gly Ala Gly Ile  
 135 140 145 150

aag gac atc aag tcc ggc gaa aag gtg ctt atc gac ggc agc ctc ggc	595
Lys Asp Ile Lys Ser Gly Glu Lys Val Leu Ile Asp Gly Ser Leu Gly	
155 160 165	
acc att gac cgc aac gcg gac gaa gct gaa gca acc aag ctc gtc tcc	643
Thr Ile Asp Arg Asn Ala Asp Glu Ala Glu Ala Thr Lys Leu Val Ser	
170 175 180	
gag tcc ctc gag cgc gct gct cgc atc gcc gag tgg aag ggt cct gca	691
Glu Ser Leu Glu Arg Ala Ala Arg Ile Ala Glu Trp Lys Gly Pro Ala	
185 190 195	
caa acc aag gac ggc tac cgc gtt cag ctg ttg gcc aac gtc caa gac	739
Gln Thr Lys Asp Gly Tyr Arg Val Gln Leu Leu Ala Asn Val Gln Asp	
200 205 210	
ggc aac tct gca cag cag gct gca cag acc gaa gca gaa ggc atc ggc	787
Gly Asn Ser Ala Gln Gln Ala Ala Gln Thr Glu Ala Glu Gly Ile Gly	
215 220 225 230	
ctg ttc cgc acc gaa ctg tgc ttc ctt tcc gcc acc gaa gag cca agc	835
Leu Phe Arg Thr Glu Leu Cys Phe Leu Ser Ala Thr Glu Glu Pro Ser	
235 240 245	
gtt gat gag cag gct gcg gtc tac tca aag gtg ctt gaa gca ttc cca	883
Val Asp Glu Gln Ala Ala Val Tyr Ser Lys Val Leu Glu Ala Phe Pro	
250 255 260	
gag tcc aag gtc gtt gtc cgc tcc ctc gac gca ggt tct gac aag cca	931
Glu Ser Lys Val Val Val Arg Ser Leu Asp Ala Gly Ser Asp Lys Pro	
265 270 275	
gtt cca ttc gca tgc atg gct gat gag atg aac cca gca ctg ggt gtt	979
Val Pro Phe Ala Ser Met Ala Asp Glu Met Asn Pro Ala Leu Gly Val	
280 285 290	
cgt ggc ctg cgt atc gca cgt gga cag gtt gat ctg ctg act cgc cag	1027
Arg Gly Leu Arg Ile Ala Arg Gly Gln Val Asp Leu Leu Thr Arg Gln	
295 300 305 310	
ctc gac gca att gcg aag gcc agc gaa gaa ctc ggc cgt ggc gac gac	1075
Leu Asp Ala Ile Ala Lys Ala Ser Glu Glu Leu Gly Arg Gly Asp Asp	
315 320 325	
gcc cca acc tgg gtt atg gct cca atg gtg gct acc gct tat gaa gca	1123
Ala Pro Thr Trp Val Met Ala Pro Met Val Ala Thr Ala Tyr Glu Ala	
330 335 340	
aag tgg ttt gct gac atg tgc cgt gag cgt ggc cta atc gcc ggc gcc	1171
Lys Trp Phe Ala Asp Met Cys Arg Glu Arg Gly Leu Ile Ala Gly Ala	
345 350 355	
atg atc gaa gtt cca gca gca tcc ctg atg gca gac aag atc atg cct	1219
Met Ile Glu Val Pro Ala Ala Ser Leu Met Ala Asp Lys Ile Met Pro	
360 365 370	
cac ctg gac ttt gtt tcc atc ggt acc aac gac ctg acc cag tac acc	1267
His Leu Asp Phe Val Ser Ile Gly Thr Asn Asp Leu Thr Gln Tyr Thr	
375 380 385 390	



atg gca gcg gac cgc atg tct cct gag ctt gcc tac ctg acc gat cct 1315  
 Met Ala Ala Asp Arg Met Ser Pro Glu Leu Ala Tyr Leu Thr Asp Pro  
 395 400 405

tgg cag cca gca gtc ctg cgc ctg atc aag cac acc tgt gac gaa ggt 1363  
 Trp Gln Pro Ala Val Leu Arg Leu Ile Lys His Thr Cys Asp Glu Gly  
 410 415 420

gct cgc ttt aac acc ccg gtc ggt gtt tgt ggt gaa gca gca gca gac 1411  
 Ala Arg Phe Asn Thr Pro Val Gly Val Cys Gly Glu Ala Ala Ala Asp  
 425 430 435

cca ctg ttg gca act gtc ctc acc ggt ctt ggc gtg aac tcc ctg tcc 1459  
 Pro Leu Leu Ala Thr Val Leu Thr Gly Leu Gly Val Asn Ser Leu Ser  
 440 445 450

gca gca tcc act gct ctc gca gca gtc ggt gca aag ctg tca gag gtc 1507  
 Ala Ala Ser Thr Ala Leu Ala Ala Val Gly Ala Lys Leu Ser Glu Val  
 455 460 465 470

acc ctg gaa acc tgt aag aag gca gca gaa gca gca ctt gac gct gaa 1555  
 Thr Leu Glu Thr Cys Lys Lys Ala Ala Glu Ala Ala Leu Asp Ala Glu  
 475 480 485

ggt gca act gaa gca cgc gat gct gta cgc gca gtg atc gac gca gca 1603  
 Gly Ala Thr Glu Ala Arg Asp Ala Val Arg Ala Val Ile Asp Ala Ala  
 490 495 500

gtc taaaccactg ttgagctaaa aag 1629  
 Val

&lt;210&gt; 20

&lt;211&gt; 503

&lt;212&gt; PRT

&lt;213&gt; Corynebacterium glutamicum

&lt;400&gt; 20

Leu Leu Glu Arg Ser Glu Ala Ala Glu Gly Pro Ala Ala Glu Val Leu  
 1 5 10 15

Lys Ala Thr Ala Gly Met Val Asn Asp Arg Gly Trp Arg Lys Ala Val  
 20 25 30

Ile Lys Gly Val Lys Gly Gly His Pro Ala Glu Tyr Ala Val Val Ala  
 35 40 45

Ala Thr Thr Lys Phe Ile Ser Met Phe Glu Ala Ala Gly Gly Leu Ile  
 50 55 60

Ala Glu Arg Thr Thr Asp Leu Arg Asp Ile Arg Asp Arg Val Ile Ala  
 65 70 75 80

Glu Leu Arg Gly Asp Glu Glu Pro Gly Leu Pro Ala Val Ser Gly Gln  
 85 90 95

Val Ile Leu Phe Ala Asp Asp Leu Ser Pro Ala Asp Thr Ala Ala Leu  
 100 105 110

Asp Thr Asp Leu Phe Val Gly Leu Val Thr Glu Leu Gly Gly Pro Thr

115					120					125					
Ser	His	Thr	Ala	Ile	Ile	Ala	Arg	Gln	Leu	Asn	Val	Pro	Cys	Ile	Val
130						135					140				
Ala	Ser	Gly	Ala	Gly	Ile	Lys	Asp	Ile	Lys	Ser	Gly	Glu	Lys	Val	Leu
145					150					155					160
Ile	Asp	Gly	Ser	Leu	Gly	Thr	Ile	Asp	Arg	Asn	Ala	Asp	Glu	Ala	Glu
				165					170					175	
Ala	Thr	Lys	Leu	Val	Ser	Glu	Ser	Leu	Glu	Arg	Ala	Ala	Arg	Ile	Ala
			180					185					190		
Glu	Trp	Lys	Gly	Pro	Ala	Gln	Thr	Lys	Asp	Gly	Tyr	Arg	Val	Gln	Leu
	195						200					205			
Leu	Ala	Asn	Val	Gln	Asp	Gly	Asn	Ser	Ala	Gln	Gln	Ala	Ala	Gln	Thr
210						215					220				
Glu	Ala	Glu	Gly	Ile	Gly	Leu	Phe	Arg	Thr	Glu	Leu	Cys	Phe	Leu	Ser
225					230					235					240
Ala	Thr	Glu	Glu	Pro	Ser	Val	Asp	Glu	Gln	Ala	Ala	Val	Tyr	Ser	Lys
				245					250					255	
Val	Leu	Glu	Ala	Phe	Pro	Glu	Ser	Lys	Val	Val	Val	Arg	Ser	Leu	Asp
			260					265					270		
Ala	Gly	Ser	Asp	Lys	Pro	Val	Pro	Phe	Ala	Ser	Met	Ala	Asp	Glu	Met
		275					280					285			
Asn	Pro	Ala	Leu	Gly	Val	Arg	Gly	Leu	Arg	Ile	Ala	Arg	Gly	Gln	Val
290						295					300				
Asp	Leu	Leu	Thr	Arg	Gln	Leu	Asp	Ala	Ile	Ala	Lys	Ala	Ser	Glu	Glu
305					310					315					320
Leu	Gly	Arg	Gly	Asp	Asp	Ala	Pro	Thr	Trp	Val	Met	Ala	Pro	Met	Val
				325					330				335		
Ala	Thr	Ala	Tyr	Glu	Ala	Lys	Trp	Phe	Ala	Asp	Met	Cys	Arg	Glu	Arg
			340					345					350		
Gly	Leu	Ile	Ala	Gly	Ala	Met	Ile	Glu	Val	Pro	Ala	Ala	Ser	Leu	Met
		355					360					365			
Ala	Asp	Lys	Ile	Met	Pro	His	Leu	Asp	Phe	Val	Ser	Ile	Gly	Thr	Asn
		370				375					380				
Asp	Leu	Thr	Gln	Tyr	Thr	Met	Ala	Ala	Asp	Arg	Met	Ser	Pro	Glu	Leu
385					390					395					400
Ala	Tyr	Leu	Thr	Asp	Pro	Trp	Gln	Pro	Ala	Val	Leu	Arg	Leu	Ile	Lys
				405					410					415	
His	Thr	Cys	Asp	Glu	Gly	Ala	Arg	Phe	Asn	Thr	Pro	Val	Gly	Val	Cys
			420					425					430		
Gly	Glu	Ala	Ala	Ala	Asp	Pro	Leu	Leu	Ala	Thr	Val	Leu	Thr	Gly	Leu
		435					440					445			

Gly Val Asn Ser Leu Ser Ala Ala Ser Thr Ala Leu Ala Ala Val Gly  
 450 455 460

Ala Lys Leu Ser Glu Val Thr Leu Glu Thr Cys Lys Lys Ala Ala Glu  
 465 470 475 480

Ala Ala Leu Asp Ala Glu Gly Ala Thr Glu Ala Arg Asp Ala Val Arg  
 485 490 495

Ala Val Ile Asp Ala Ala Val  
 500

<210> 21  
 <211> 390  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (101)..(367)  
 <223> RXA01300

<400> 21  
 gatcgacatt aaatcccctc ccttgggggg tttaactaac aaatcgctgc gccctaattcc 60

gttcggatta acggcgtagc aacacgaaag gacactttcc atg gct tcc aag act 115  
 Met Ala Ser Lys Thr  
 1 5

gta acc gtc ggt tcc tcc gtt ggc ctg cac gca cgt cca gca tcc atc 163  
 Val Thr Val Gly Ser Ser Val Gly Leu His Ala Arg Pro Ala Ser Ile  
 10 15 20

atc gct gaa gcg gct gct gag tac gac gac gaa atc ttg ctg acc ctg 211  
 Ile Ala Glu Ala Ala Ala Glu Tyr Asp Asp Glu Ile Leu Leu Thr Leu  
 25 30 35

gtt ggc tcc gat gat gac gaa gag acc gac gcg tcc tct tcc ctc atg 259  
 Val Gly Ser Asp Asp Asp Glu Glu Thr Asp Ala Ser Ser Ser Leu Met  
 40 45 50

atc atg gcg ctg ggc gca gag cac ggc aac gaa gtt acc gtc acc tcc 307  
 Ile Met Ala Leu Gly Ala Glu His Gly Asn Glu Val Thr Val Thr Ser  
 55 60 65

gac aac gct gaa gct gtt gag aag atc gct gcg ctt atc gca cag gac 355  
 Asp Asn Ala Glu Ala Val Glu Lys Ile Ala Ala Leu Ile Ala Gln Asp  
 70 75 80 85

ctt gac gct gag taaacaacgc tctgcttggt aaa 390  
 Leu Asp Ala Glu

<210> 22  
 <211> 89  
 <212> PRT  
 <213> Corynebacterium glutamicum

&lt;400&gt; 22

Met Ala Ser Lys Thr Val Thr Val Gly Ser Ser Val Gly Leu His Ala  
 1 5 10 15

Arg Pro Ala Ser Ile Ile Ala Glu Ala Ala Glu Tyr Asp Asp Glu  
 20 25 30

Ile Leu Leu Thr Leu Val Gly Ser Asp Asp Asp Glu Glu Thr Asp Ala  
 35 40 45

Ser Ser Ser Leu Met Ile Met Ala Leu Gly Ala Glu His Gly Asn Glu  
 50 55 60

Val Thr Val Thr Ser Asp Asn Ala Glu Ala Val Glu Lys Ile Ala Ala  
 65 70 75 80

Leu Ile Ala Gln Asp Leu Asp Ala Glu  
 85

&lt;210&gt; 23

&lt;211&gt; 508

&lt;212&gt; DNA

&lt;213&gt; Corynebacterium glutamicum

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (101)..(508)

&lt;223&gt; RXN03002

&lt;400&gt; 23

ggaacttcga ggtgtcttcg tggggcggtac ggagatctag caagtgtggc tttatgtttg 60

accctatccg aatcaacatg cagtgaatta acatctactt atg ttt gta ctc aaa 115  
 Met Phe Val Leu Lys  
 1 5

gat ctg cta aag gca gaa cgc ata gaa ctc gac cgc acg gtc acc gat 163  
 Asp Leu Leu Lys Ala Glu Arg Ile Glu Leu Asp Arg Thr Val Thr Asp  
 10 15 20

tgg cgt gaa ggc atc cgc gcc gca ggt gta ctc cta gaa aag aca aac 211  
 Trp Arg Glu Gly Ile Arg Ala Ala Gly Val Leu Leu Glu Lys Thr Asn  
 25 30 35

agc att gat tcc gcc tac acc gat gcc atg atc gcc agc gtg gaa gaa 259  
 Ser Ile Asp Ser Ala Tyr Thr Asp Ala Met Ile Ala Ser Val Glu Glu  
 40 45 50

aaa ggc ccc tac att gtg gtc gct cca ggt ttc gct ttc gcg cac gcc 307  
 Lys Gly Pro Tyr Ile Val Val Ala Pro Gly Phe Ala Phe Ala His Ala  
 55 60 65

cgc ccc agc aga gca gtc cgc gag acc gct atg tcg tgg gtg cgc ctg 355  
 Arg Pro Ser Arg Ala Val Arg Glu Thr Ala Met Ser Trp Val Arg Leu  
 70 75 80 85

gcc tcc cct gtt tcc ttc ggt cac agt aag aat gat ccc ctc aat ctc 403  
 Ala Ser Pro Val Ser Phe Gly His Ser Lys Asn Asp Pro Leu Asn Leu  
 90 95 100

atc gtt gct ctc gct gcc aaa gat gcc acc gca cat acc caa gcg atg 451  
 Ile Val Ala Leu Ala Ala Lys Asp Ala Thr Ala His Thr Gln Ala Met  
                   105                  110                  115

gcg gca ttg gct aaa gct tta gga aaa tac cga aag gat ctc gac gag 499  
 Ala Ala Leu Ala Lys Ala Leu Gly Lys Tyr Arg Lys Asp Leu Asp Glu  
                   120                  125                  130

gca caa agt 508  
 Ala Gln Ser  
                   135

<210> 24

<211> 136

<212> PRT

<213> Corynebacterium glutamicum

<400> 24

Met Phe Val Leu Lys Asp Leu Leu Lys Ala Glu Arg Ile Glu Leu Asp  
           1                  5                  10                  15

Arg Thr Val Thr Asp Trp Arg Glu Gly Ile Arg Ala Ala Gly Val Leu  
                   20                  25                  30

Leu Glu Lys Thr Asn Ser Ile Asp Ser Ala Tyr Thr Asp Ala Met Ile  
                   35                  40                  45

Ala Ser Val Glu Glu Lys Gly Pro Tyr Ile Val Val Ala Pro Gly Phe  
           50                  55                  60

Ala Phe Ala His Ala Arg Pro Ser Arg Ala Val Arg Glu Thr Ala Met  
           65                  70                  75                  80

Ser Trp Val Arg Leu Ala Ser Pro Val Ser Phe Gly His Ser Lys Asn  
                   85                  90                  95

Asp Pro Leu Asn Leu Ile Val Ala Leu Ala Ala Lys Asp Ala Thr Ala  
                   100                  105                  110

His Thr Gln Ala Met Ala Ala Leu Ala Lys Ala Leu Gly Lys Tyr Arg  
                   115                  120                  125

Lys Asp Leu Asp Glu Ala Gln Ser  
           130                  135

<210> 25

<211> 789

<212> DNA

<213> Corynebacterium glutamicum

<220>

<221> CDS

<222> (14)..(766)

<223> RXC00953

<400> 25

cttgcatgcc ccaatg gcg cca cca acg gta ggc aac tac atc atg cag tcc 52  
                   Met Ala Pro Pro Thr Val Gly Asn Tyr Ile Met Gln Ser  
                   1                  5                  10

ttc act caa ggt ctg cag ttc ggc gtt gca gtt gcc gtg att ctc ttt	100
Phe Thr Gln Gly Leu Gln Phe Gly Val Ala Val Ala Val Ile Leu Phe	
15 20 25	
ggt gtc cgc acc att ctt ggt gaa ctg gtc ccc gca ttc caa ggt att	148
Gly Val Arg Thr Ile Leu Gly Glu Leu Val Pro Ala Phe Gln Gly Ile	
30 35 40 45	
gct gcg aag gtt gtt ccc gga gct atc ccc gca ttg gat gca ccg atc	196
Ala Ala Lys Val Val Pro Gly Ala Ile Pro Ala Leu Asp Ala Pro Ile	
50 55 60	
gtg ttc ccc tac gcg cag aac gcc gtt ctc att ggt ttc ttg tct tcc	244
Val Phe Pro Tyr Ala Gln Asn Ala Val Leu Ile Gly Phe Leu Ser Ser	
65 70 75	
ttc gtc ggt ggc ttg gtt ggc ctg act gtt ctt gca tcg tgg ctg aac	292
Phe Val Gly Gly Leu Val Gly Leu Thr Val Leu Ala Ser Trp Leu Asn	
80 85 90	
cca gct ttt ggt gtc gcg ttg att ctg cct ggt ttg gtc ccc cac ttc	340
Pro Ala Phe Gly Val Ala Leu Ile Leu Pro Gly Leu Val Pro His Phe	
95 100 105	
ttc act ggt ggc gcg gcg ggc gtt tac ggt aat gcc acg ggt ggt cgt	388
Phe Thr Gly Gly Ala Ala Gly Val Tyr Gly Asn Ala Thr Gly Gly Arg	
110 115 120 125	
cga gga gca gta ttt ggc gcc ttt gcc aac ggt ctt ctg att acc ttc	436
Arg Gly Ala Val Phe Gly Ala Phe Ala Asn Gly Leu Leu Ile Thr Phe	
130 135 140	
ctc cct gct ttc ctg ctt ggt gtg ctt ggt tcc ttc ggg tca gag aac	484
Leu Pro Ala Phe Leu Leu Gly Val Leu Gly Ser Phe Gly Ser Glu Asn	
145 150 155	
acc act ttc ggt gat gcg gac ttt ggt tgg ttc gga atc gtt gtt ggt	532
Thr Thr Phe Gly Asp Ala Asp Phe Gly Trp Phe Gly Ile Val Val Gly	
160 165 170	
tct gca gcc aag gtg gaa ggt gct ggc ggg ctc atc ttg ttg ctc atc	580
Ser Ala Ala Lys Val Glu Gly Ala Gly Leu Ile Leu Leu Leu Ile	
175 180 185	
atc gca gcg gtt ctt ctg ggt ggc gcg atg gtc ttc cag aag cgc gtc	628
Ile Ala Ala Val Leu Leu Gly Gly Ala Met Val Phe Gln Lys Arg Val	
190 195 200 205	
gtg aat ggg cac tgg gat cca gct ccc aac cgt gag cgc gtg gag aag	676
Val Asn Gly His Trp Asp Pro Ala Pro Asn Arg Glu Arg Val Glu Lys	
210 215 220	
gcg gaa gct gat gcc act cca acg gct ggg gct cgg acc tac cct aag	724
Ala Glu Ala Asp Ala Thr Pro Thr Ala Gly Ala Arg Thr Tyr Pro Lys	
225 230 235	
att gct cct ccg gcg ggc gct cct acc cca ccg gct cga agc	766
Ile Ala Pro Pro Ala Gly Ala Pro Thr Pro Pro Ala Arg Ser	
240 245 250	

taagatctcc aaaaccctga gat

789

&lt;210&gt; 26

&lt;211&gt; 251

&lt;212&gt; PRT

&lt;213&gt; Corynebacterium glutamicum

&lt;400&gt; 26

```

Met Ala Pro Pro Thr Val Gly Asn Tyr Ile Met Gln Ser Phe Thr Gln
  1           5           10           15

Gly Leu Gln Phe Gly Val Ala Val Ala Val Ile Leu Phe Gly Val Arg
          20           25           30

Thr Ile Leu Gly Glu Leu Val Pro Ala Phe Gln Gly Ile Ala Ala Lys
      35           40           45

Val Val Pro Gly Ala Ile Pro Ala Leu Asp Ala Pro Ile Val Phe Pro
      50           55           60

Tyr Ala Gln Asn Ala Val Leu Ile Gly Phe Leu Ser Ser Phe Val Gly
      65           70           75           80

Gly Leu Val Gly Leu Thr Val Leu Ala Ser Trp Leu Asn Pro Ala Phe
          85           90           95

Gly Val Ala Leu Ile Leu Pro Gly Leu Val Pro His Phe Phe Thr Gly
          100          105          110

Gly Ala Ala Gly Val Tyr Gly Asn Ala Thr Gly Gly Arg Arg Gly Ala
          115          120          125

Val Phe Gly Ala Phe Ala Asn Gly Leu Leu Ile Thr Phe Leu Pro Ala
      130           135           140

Phe Leu Leu Gly Val Leu Gly Ser Phe Gly Ser Glu Asn Thr Thr Phe
      145           150           155           160

Gly Asp Ala Asp Phe Gly Trp Phe Gly Ile Val Val Gly Ser Ala Ala
          165          170          175

Lys Val Glu Gly Ala Gly Gly Leu Ile Leu Leu Leu Ile Ile Ala Ala
          180          185          190

Val Leu Leu Gly Gly Ala Met Val Phe Gln Lys Arg Val Val Asn Gly
      195           200           205

His Trp Asp Pro Ala Pro Asn Arg Glu Arg Val Glu Lys Ala Glu Ala
      210           215           220

Asp Ala Thr Pro Thr Ala Gly Ala Arg Thr Tyr Pro Lys Ile Ala Pro
      225           230           235           240

Pro Ala Gly Ala Pro Thr Pro Pro Ala Arg Ser
          245          250

```

&lt;210&gt; 27

&lt;211&gt; 553

&lt;212&gt; DNA

&lt;213&gt; Corynebacterium glutamicum

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (101)..(553)

&lt;223&gt; RXC03001

&lt;400&gt; 27

```

cccggttcac gtgatcaatg acttcacgag caccgatgaa atcgatgctg cgcttcgtga 60
acgctacgac atctaactac tttaaaagga cgaaaatatt atg gac tgg tta acc 115
                                         Met Asp Trp Leu Thr
                                         1 5

att cct ctt ttc ctc gtt aat gaa atc ctt gcg gtt ccg gct ttc ctc 163
Ile Pro Leu Phe Leu Val Asn Glu Ile Leu Ala Val Pro Ala Phe Leu
                        10 15 20

atc ggt atc atc acc gcc gtg gga ttg ggt gcc atg ggg cgt tcc gtc 211
Ile Gly Ile Ile Thr Ala Val Gly Leu Gly Ala Met Gly Arg Ser Val
                        25 30 35

ggt cag gtt atc ggt gga gca atc aaa gca acg ttg ggc ttt ttg ctc 259
Gly Gln Val Ile Gly Gly Ala Ile Lys Ala Thr Leu Gly Phe Leu Leu
                        40 45 50

att ggt gcg ggt gcc acg ttg gtc act gcc tcc ctg gag cca ctg ggt 307
Ile Gly Ala Gly Ala Thr Leu Val Thr Ala Ser Leu Glu Pro Leu Gly
                        55 60 65

gcg atg atc atg ggt gcc aca ggc atg cgt ggt gtt gtc cca acg aat 355
Ala Met Ile Met Gly Ala Thr Gly Met Arg Gly Val Val Pro Thr Asn
                        70 75 80 85

gaa gcc atc gcc gga atc gca cag gct gaa tac ggc gcg cag gtg gcg 403
Glu Ala Ile Ala Gly Ile Ala Gln Ala Glu Tyr Gly Ala Gln Val Ala
                        90 95 100

tgg ctg atg att ctg ggc ttc gcc atc tct ttg gtg ttg gct cgt ttc 451
Trp Leu Met Ile Leu Gly Phe Ala Ile Ser Leu Val Leu Ala Arg Phe
                        105 110 115

acc aac ctg cgt tat gtc ttg ctc aac gga cac cac gtg ctg ttg atg 499
Thr Asn Leu Arg Tyr Val Leu Leu Asn Gly His His Val Leu Leu Met
                        120 125 130

tgc acc atg ctc acc atg gtc ttg gcc acc gga aga gtt gat gcg tgg 547
Cys Thr Met Leu Thr Met Val Leu Ala Thr Gly Arg Val Asp Ala Trp
                        135 140 145

atc ttc
Ile Phe
150
553

```

&lt;210&gt; 28

&lt;211&gt; 151

&lt;212&gt; PRT

&lt;213&gt; Corynebacterium glutamicum

&lt;400&gt; 28



Met Asp Trp Leu Thr Ile Pro Leu Phe Leu Val Asn Glu Ile Leu Ala  
 1 5 10 15  
 Val Pro Ala Phe Leu Ile Gly Ile Ile Thr Ala Val Gly Leu Gly Ala  
 20 25 30  
 Met Gly Arg Ser Val Gly Gln Val Ile Gly Gly Ala Ile Lys Ala Thr  
 35 40 45  
 Leu Gly Phe Leu Leu Ile Gly Ala Gly Ala Thr Leu Val Thr Ala Ser  
 50 55 60  
 Leu Glu Pro Leu Gly Ala Met Ile Met Gly Ala Thr Gly Met Arg Gly  
 65 70 75 80  
 Val Val Pro Thr Asn Glu Ala Ile Ala Gly Ile Ala Gln Ala Glu Tyr  
 85 90 95  
 Gly Ala Gln Val Ala Trp Leu Met Ile Leu Gly Phe Ala Ile Ser Leu  
 100 105 110  
 Val Leu Ala Arg Phe Thr Asn Leu Arg Tyr Val Leu Leu Asn Gly His  
 115 120 125  
 His Val Leu Leu Met Cys Thr Met Leu Thr Met Val Leu Ala Thr Gly  
 130 135 140  
 Arg Val Asp Ala Trp Ile Phe  
 145 150

<210> 29  
 <211> 2172  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (101)..(2149)  
 <223> RXN01943

<400> 29  
 ccgattcttt ttcggcccaa ttcgtaacgg cgatcctctt aagtggacaa gaaagtctct. 60  
 tgcccgcggg agacagaccc tacgtttaga aaggtttgac atg gcg tcc aaa ctg 115  
 Met Ala Ser Lys Leu  
 1 5  
 acg acg aca tcg caa cat att ctg gaa aac ctt ggt gga cca gac aat 163  
 Thr Thr Thr Ser Gln His Ile Leu Glu Asn Leu Gly Gly Pro Asp Asn  
 10 15 20  
 att act tcg atg act cac tgt gcg act cgc ctt cgc ttc caa gtg aag 211  
 Ile Thr Ser Met Thr His Cys Ala Thr Arg Leu Arg Phe Gln Val Lys  
 25 30 35  
 gat caa tcc att gtt gat caa caa gaa att gac tcc gac cca tca gtt 259  
 Asp Gln Ser Ile Val Asp Gln Gln Glu Ile Asp Ser Asp Pro Ser Val  
 40 45 50  
 ctt ggc gta gta ccc caa gga tcc acc ggt atg cag gtg gtg atg ggt 307

Leu	Gly	Val	Val	Pro	Gln	Gly	Ser	Thr	Gly	Met	Gln	Val	Val	Met	Gly		
55						60					65						
gga	tct	gtt	gca	aac	tat	tac	caa	gaa	atc	ctc	aaa	ctt	gat	gga	atg	355	
Gly	Ser	Val	Ala	Asn	Tyr	Tyr	Gln	Glu	Ile	Leu	Lys	Leu	Asp	Gly	Met	85	
70					75					80							
aag	cac	ttc	gcc	gac	ggg	gaa	gct	aca	gag	agt	tca	tcc	aag	aag	gaa	403	
Lys	His	Phe	Ala	Asp	Gly	Glu	Ala	Thr	Glu	Ser	Ser	Ser	Lys	Lys	Glu	100	
				90					95								
tac	ggc	gga	gtc	cgt	ggc	aag	tac	tcg	tgg	att	gac	tac	gcc	ttc	gag	451	
Tyr	Gly	Gly	Val	Arg	Gly	Lys	Tyr	Ser	Trp	Ile	Asp	Tyr	Ala	Phe	Glu	115	
			105					110					115				
ttc	ttg	tct	gat	act	ttc	cga	cca	atc	ctg	tgg	gcc	ctg	ctt	ggg	gcc	499	
Phe	Leu	Ser	Asp	Thr	Phe	Arg	Pro	Ile	Leu	Trp	Ala	Leu	Leu	Gly	Ala	130	
			120				125										
tca	ctg	att	att	acc	ttg	ttg	gtt	ctt	gcg	gat	act	ttc	ggg	ttg	caa	547	
Ser	Leu	Ile	Ile	Thr	Leu	Leu	Val	Leu	Ala	Asp	Thr	Phe	Gly	Leu	Gln	145	
			135				140										
gac	ttc	cgc	gct	cca	atg	gat	gag	cag	cct	gat	act	tat	gta	ttc	ctg	595	
Asp	Phe	Arg	Ala	Pro	Met	Asp	Glu	Gln	Pro	Asp	Thr	Tyr	Val	Phe	Leu	165	
150					155					160							
cac	tcc	atg	tgg	cgc	tcg	gtc	ttc	tac	ttc	ctg	cca	att	atg	gtt	ggg	643	
His	Ser	Met	Trp	Arg	Ser	Val	Phe	Tyr	Phe	Leu	Pro	Ile	Met	Val	Gly	180	
				170					175								
gcc	acc	gca	gct	cga	aag	ctc	ggc	gca	aac	gag	tgg	att	ggg	gca	gct	691	
Ala	Thr	Ala	Ala	Arg	Lys	Leu	Gly	Ala	Asn	Glu	Trp	Ile	Gly	Ala	Ala	195	
			185					190									
att	cca	gcc	gca	ctt	ctt	act	cca	gaa	ttc	ttg	gca	ctg	ggg	tct	gcc	739	
Ile	Pro	Ala	Ala	Leu	Leu	Thr	Pro	Glu	Phe	Leu	Ala	Leu	Gly	Ser	Ala	210	
			200				205										
ggc	gat	acc	gtc	aca	gtc	ttt	ggc	ctg	cca	atg	gtt	ctg	aat	gac	tac	787	
Gly	Asp	Thr	Val	Thr	Val	Phe	Gly	Leu	Pro	Met	Val	Leu	Asn	Asp	Tyr	225	
			215			220											
tcc	gga	cag	gta	ttc	cca	ccg	ctg	att	gca	gca	att	ggg	ctg	tac	tgg	835	
Ser	Gly	Gln	Val	Phe	Pro	Pro	Leu	Ile	Ala	Ala	Ile	Gly	Leu	Tyr	Trp	245	
230					235					240							
gtg	gaa	aag	gga	ctg	aag	aag	atc	atc	cct	gaa	gca	gtc	caa	atg	gtg	883	
Val	Glu	Lys	Gly	Leu	Lys	Lys	Ile	Ile	Pro	Glu	Ala	Val	Gln	Met	Val	260	
				250					255								
ttc	gtc	cca	ttc	ttc	tcc	ctg	ctg	att	atg	atc	cca	gcg	acc	gca	ttc	931	
Phe	Val	Pro	Phe	Phe	Ser	Leu	Leu	Ile	Met	Ile	Pro	Ala	Thr	Ala	Phe	275	
			265					270									
ctg	ctt	gga	cct	ttc	ggc	atc	ggg	gtt	ggg	aac	gga	att	tcc	aac	ctg	979	
Leu	Leu	Gly	Pro	Phe	Gly	Ile	Gly	Val	Gly	Asn	Gly	Ile	Ser	Asn	Leu	290	
			280				285					290					
ctt	gaa	gcg	att	aac	aac	ttc	agc	cca	ttt	att	ctt	tcc	atc	gtt	atc	1027	
Leu	Glu	Ala	Ile	Asn	Asn	Phe	Ser	Pro	Phe	Ile	Leu	Ser	Ile	Val	Ile		

295	300	305	
cca ttg ctc tac cca ttc ttg gtt cca ctt gga ttg cac tgg cca cta			1075
Pro Leu Leu Tyr Pro Phe Leu Val Pro Leu Gly Leu His Trp Pro Leu			
310	315	320	325
aac gcc atc atg atc cag aac atc aac acc ctg ggt tac gac ttc att			1123
Asn Ala Ile Met Ile Gln Asn Ile Asn Thr Leu Gly Tyr Asp Phe Ile			
	330	335	340
cag gga cca atg ggt gcc tgg aac ttc gcc tgc ttc ggc ctg gtc acc			1171
Gln Gly Pro Met Gly Ala Trp Asn Phe Ala Cys Phe Gly Leu Val Thr			
	345	350	355
ggc gtg ttc ttg ctc tcc att aag gaa cga aac aag gcc atg cgt cag			1219
Gly Val Phe Leu Leu Ser Ile Lys Glu Arg Asn Lys Ala Met Arg Gln			
	360	365	370
gtt tcc ctg ggt ggc atg ttg gct ggt ttg ctc ggc ggc att tcc gag			1267
Val Ser Leu Gly Gly Met Leu Ala Gly Leu Leu Gly Gly Ile Ser Glu			
	375	380	385
cct tcc ctc tac ggt gtt ctg ctc cga ttc aag aag acc tac ttc cgc			1315
Pro Ser Leu Tyr Gly Val Leu Leu Arg Phe Lys Lys Thr Tyr Phe Arg			
	390	395	400
ctc ctg ccg ggt tgt ttg gca ggc ggt atc gtg atg ggc atc ttc gac			1363
Leu Leu Pro Gly Cys Leu Ala Gly Gly Ile Val Met Gly Ile Phe Asp			
	410	415	420
atc aag gcg tac gct ttc gtg ttc acc tcc ttg ctt acc atc cca gca			1411
Ile Lys Ala Tyr Ala Phe Val Phe Thr Ser Leu Leu Thr Ile Pro Ala			
	425	430	435
atg gac cca tgg ttg ggc tac acc att ggt atc gca gtt gca ttc ttc			1459
Met Asp Pro Trp Leu Gly Tyr Thr Ile Gly Ile Ala Val Ala Phe Phe			
	440	445	450
gtt tcc atg ttc ctt gtt ctc gca ctg gac tac cgt tcc aac gaa gag			1507
Val Ser Met Phe Leu Val Leu Ala Leu Asp Tyr Arg Ser Asn Glu Glu			
	455	460	465
cgc gat gag gca cgt gca aag gtt gct gct gac aag cag gca gaa gaa			1555
Arg Asp Glu Ala Arg Ala Lys Val Ala Ala Asp Lys Gln Ala Glu Glu			
	470	475	480
gat ctg aag gca gaa gct aat gca act cct gca gct cca gta gct gct			1603
Asp Leu Lys Ala Glu Ala Asn Ala Thr Pro Ala Ala Pro Val Ala Ala			
	490	495	500
gca ggt gcg gga gcc ggt gca ggt gca gga gcc gct gct ggc gct gca			1651
Ala Gly Ala Gly Ala Gly Ala Gly Ala Gly Ala Ala Ala Gly Ala Ala			
	505	510	515
acc gcc gtg gca gct aag ccg aag ctg gcc gct ggg gaa gta gtg gac			1699
Thr Ala Val Ala Ala Lys Pro Lys Leu Ala Ala Gly Glu Val Val Asp			
	520	525	530
att gtt tcc cca ctc gaa ggc aag gca att cca ctt tct gaa gta cct			1747
Ile Val Ser Pro Leu Glu Gly Lys Ala Ile Pro Leu Ser Glu Val Pro			
	535	540	545

gac cca atc ttt gca gca ggc aag ctt gga cca ggc att gca atc caa 1795  
 Asp Pro Ile Phe Ala Ala Gly Lys Leu Gly Pro Gly Ile Ala Ile Gln  
 550 555 560 565

cca act gga aac acc gtt gtt gct cca gca gac gct act gtc atc ctt 1843  
 Pro Thr Gly Asn Thr Val Val Ala Pro Ala Asp Ala Thr Val Ile Leu  
 570 575 580

gtc cag aaa tct gga cac gca gtg gca ttg cgc tta gat agc gga gtt 1891  
 Val Gln Lys Ser Gly His Ala Val Ala Leu Arg Leu Asp Ser Gly Val  
 585 590 595

gaa atc ctt gtc cac gtt gga ttg gac acc gtg caa ttg ggc ggc gaa 1939  
 Glu Ile Leu Val His Val Gly Leu Asp Thr Val Gln Leu Gly Gly Glu  
 600 605 610

ggc ttc acc gtt cac gtt gag cgc agg cag caa gtc aag gcg ggg gat 1987  
 Gly Phe Thr Val His Val Glu Arg Arg Gln Gln Val Lys Ala Gly Asp  
 615 620 625

cca ctg atc act ttt gac gct gac ttc att cga tcc aag gat cta cct 2035  
 Pro Leu Ile Thr Phe Asp Ala Asp Phe Ile Arg Ser Lys Asp Leu Pro  
 630 635 640 645

ttg atc acc cca gtt gtg gtg tct aac gcc gcg aaa ttc ggt gaa att 2083  
 Leu Ile Thr Pro Val Val Val Ser Asn Ala Ala Lys Phe Gly Glu Ile  
 650 655 660

gaa ggt att cct gca gat cag gca aat tct tcc acg act gtg atc aag 2131  
 Glu Gly Ile Pro Ala Asp Gln Ala Asn Ser Ser Thr Thr Val Ile Lys  
 665 670 675

gtc aac ggc aag aac gag taacctggga tccatgttgc gca 2172  
 Val Asn Gly Lys Asn Glu  
 680

&lt;210&gt; 30

&lt;211&gt; 683

&lt;212&gt; PRT

<213> *Corynebacterium glutamicum*

&lt;400&gt; 30

Met Ala Ser Lys Leu Thr Thr Thr Ser Gln His Ile Leu Glu Asn Leu  
 1 5 10 15

Gly Gly Pro Asp Asn Ile Thr Ser Met Thr His Cys Ala Thr Arg Leu  
 20 25 30

Arg Phe Gln Val Lys Asp Gln Ser Ile Val Asp Gln Gln Glu Ile Asp  
 35 40 45

Ser Asp Pro Ser Val Leu Gly Val Val Pro Gln Gly Ser Thr Gly Met  
 50 55 60

Gln Val Val Met Gly Gly Ser Val Ala Asn Tyr Tyr Gln Glu Ile Leu  
 65 70 75 80

Lys Leu Asp Gly Met Lys His Phe Ala Asp Gly Glu Ala Thr Glu Ser  
 85 90 95

Ser Ser Lys Lys Glu Tyr Gly Gly Val Arg Gly Lys Tyr Ser Trp Ile  
 100 105 110  
 Asp Tyr Ala Phe Glu Phe Leu Ser Asp Thr Phe Arg Pro Ile Leu Trp  
 115 120 125  
 Ala Leu Leu Gly Ala Ser Leu Ile Ile Thr Leu Leu Val Leu Ala Asp  
 130 135 140  
 Thr Phe Gly Leu Gln Asp Phe Arg Ala Pro Met Asp Glu Gln Pro Asp  
 145 150 155 160  
 Thr Tyr Val Phe Leu His Ser Met Trp Arg Ser Val Phe Tyr Phe Leu  
 165 170 175  
 Pro Ile Met Val Gly Ala Thr Ala Ala Arg Lys Leu Gly Ala Asn Glu  
 180 185 190  
 Trp Ile Gly Ala Ala Ile Pro Ala Ala Leu Leu Thr Pro Glu Phe Leu  
 195 200 205  
 Ala Leu Gly Ser Ala Gly Asp Thr Val Thr Val Phe Gly Leu Pro Met  
 210 215 220  
 Val Leu Asn Asp Tyr Ser Gly Gln Val Phe Pro Pro Leu Ile Ala Ala  
 225 230 235 240  
 Ile Gly Leu Tyr Trp Val Glu Lys Gly Leu Lys Lys Ile Ile Pro Glu  
 245 250 255  
 Ala Val Gln Met Val Phe Val Pro Phe Phe Ser Leu Leu Ile Met Ile  
 260 265 270  
 Pro Ala Thr Ala Phe Leu Leu Gly Pro Phe Gly Ile Gly Val Gly Asn  
 275 280 285  
 Gly Ile Ser Asn Leu Leu Glu Ala Ile Asn Asn Phe Ser Pro Phe Ile  
 290 295 300  
 Leu Ser Ile Val Ile Pro Leu Leu Tyr Pro Phe Leu Val Pro Leu Gly  
 305 310 315 320  
 Leu His Trp Pro Leu Asn Ala Ile Met Ile Gln Asn Ile Asn Thr Leu  
 325 330 335  
 Gly Tyr Asp Phe Ile Gln Gly Pro Met Gly Ala Trp Asn Phe Ala Cys  
 340 345 350  
 Phe Gly Leu Val Thr Gly Val Phe Leu Leu Ser Ile Lys Glu Arg Asn  
 355 360 365  
 Lys Ala Met Arg Gln Val Ser Leu Gly Gly Met Leu Ala Gly Leu Leu  
 370 375 380  
 Gly Gly Ile Ser Glu Pro Ser Leu Tyr Gly Val Leu Leu Arg Phe Lys  
 385 390 395 400  
 Lys Thr Tyr Phe Arg Leu Leu Pro Gly Cys Leu Ala Gly Gly Ile Val  
 405 410 415

Met Gly Ile Phe Asp Ile Lys Ala Tyr Ala Phe Val Phe Thr Ser Leu  
                   420                                  425                                  430  
 Leu Thr Ile Pro Ala Met Asp Pro Trp Leu Gly Tyr Thr Ile Gly Ile  
                   435                                  440                                  445  
 Ala Val Ala Phe Phe Val Ser Met Phe Leu Val Leu Ala Leu Asp Tyr  
                   450                                  455                                  460  
 Arg Ser Asn Glu Glu Arg Asp Glu Ala Arg Ala Lys Val Ala Ala Asp  
                   465                                  470                                  475                                  480  
 Lys Gln Ala Glu Glu Asp Leu Lys Ala Glu Ala Asn Ala Thr Pro Ala  
                                   485                                  490                                  495  
 Ala Pro Val Ala Ala Ala Gly Ala Gly Ala Gly Ala Gly Ala Gly Ala  
                                   500                                  505                                  510  
 Ala Ala Gly Ala Ala Thr Ala Val Ala Ala Lys Pro Lys Leu Ala Ala  
                   515                                  520                                  525  
 Gly Glu Val Val Asp Ile Val Ser Pro Leu Glu Gly Lys Ala Ile Pro  
                   530                                  535                                  540  
 Leu Ser Glu Val Pro Asp Pro Ile Phe Ala Ala Gly Lys Leu Gly Pro  
                   545                                  550                                  555                                  560  
 Gly Ile Ala Ile Gln Pro Thr Gly Asn Thr Val Val Ala Pro Ala Asp  
                                   565                                  570                                  575  
 Ala Thr Val Ile Leu Val Gln Lys Ser Gly His Ala Val Ala Leu Arg  
                                   580                                  585                                  590  
 Leu Asp Ser Gly Val Glu Ile Leu Val His Val Gly Leu Asp Thr Val  
                   595                                  600                                  605  
 Gln Leu Gly Gly Glu Gly Phe Thr Val His Val Glu Arg Arg Gln Gln  
                   610                                  615                                  620  
 Val Lys Ala Gly Asp Pro Leu Ile Thr Phe Asp Ala Asp Phe Ile Arg  
                   625                                  630                                  635                                  640  
 Ser Lys Asp Leu Pro Leu Ile Thr Pro Val Val Val Ser Asn Ala Ala  
                                   645                                  650                                  655  
 Lys Phe Gly Glu Ile Glu Gly Ile Pro Ala Asp Gln Ala Asn Ser Ser  
                   660                                  665                                  670  
 Thr Thr Val Ile Lys Val Asn Gly Lys Asn Glu  
                   675                                  680

<210> 31  
 <211> 1339  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (101)..(1339)  
 <223> FRXA02191

&lt;400&gt; 31

```

ccgattcttt ttcggcccaa ttcgtaacgg cgatcctctt aagtggacaa gaaagtctct 60

tgcccgcggg agacagaccc tacgtttaga aaggtttgac atg gcg tcc aaa ctg 115
                                         Met Ala Ser Lys Leu
                                         1           5

acg acg aca tcg caa cat att ctg gaa aac ctt ggt gga cca gac aat 163
Thr Thr Thr Ser Gln His Ile Leu Glu Asn Leu Gly Gly Pro Asp Asn
                        10                        15                        20

att act tcg atg act cac tgt gcg act cgc ctt cgc ttc caa gtg aag 211
Ile Thr Ser Met Thr His Cys Ala Thr Arg Leu Arg Phe Gln Val Lys
                        25                        30                        35

gat caa tcc att gtt gat caa caa gaa att gac tcc gac cca tca gtt 259
Asp Gln Ser Ile Val Asp Gln Gln Glu Ile Asp Ser Asp Pro Ser Val
                        40                        45                        50

ctt ggc gta gta ccc caa gga tcc acc ggt atg cag gtg gtg atg ggt 307
Leu Gly Val Val Pro Gln Gly Ser Thr Gly Met Gln Val Val Met Gly
                        55                        60                        65

gga tct gtt gca aac tat tac caa gaa atc ctc aaa ctt gat gga atg 355
Gly Ser Val Ala Asn Tyr Tyr Gln Glu Ile Leu Lys Leu Asp Gly Met
                        70                        75                        80                        85

aag cac ttc gcc gac ggt gaa gct aca gag agt tca tcc aag aag gaa 403
Lys His Phe Ala Asp Gly Glu Ala Thr Glu Ser Ser Ser Lys Lys Glu
                        90                        95                        100

tac ggc gga gtc cgt ggc aag tac tcg tgg att gac tac gcc ttc gag 451
Tyr Gly Gly Val Arg Gly Lys Tyr Ser Trp Ile Asp Tyr Ala Phe Glu
                        105                        110                        115

ttc ttg tct gat act ttc cga cca atc ctg tgg gcc ctg ctt ggt gcc 499
Phe Leu Ser Asp Thr Phe Arg Pro Ile Leu Trp Ala Leu Leu Gly Ala
                        120                        125                        130

tca ctg att att acc ttg ttg gtt ctt gcg gat act ttc ggt ttg caa 547
Ser Leu Ile Ile Thr Leu Leu Val Leu Ala Asp Thr Phe Gly Leu Gln
                        135                        140                        145

gac ttc cgc gct cca atg gat gag cag cct gat act tat gta ttc ctg 595
Asp Phe Arg Ala Pro Met Asp Glu Gln Pro Asp Thr Tyr Val Phe Leu
                        150                        155                        160                        165

cac tcc atg tgg cgc tcg gtc ttc tac ttc ctg cca att atg gtt ggt 643
His Ser Met Trp Arg Ser Val Phe Tyr Phe Leu Pro Ile Met Val Gly
                        170                        175                        180

gcc acc gca gct cga aag ctc ggc gca aac gag tgg att ggt gca gct 691
Ala Thr Ala Ala Arg Lys Leu Gly Ala Asn Glu Trp Ile Gly Ala Ala
                        185                        190                        195

att cca gcc gca ctt ctt act cca gaa ttc ttg gca ctg ggt tct gcc 739
Ile Pro Ala Ala Leu Leu Thr Pro Glu Phe Leu Ala Leu Gly Ser Ala
                        200                        205                        210

ggc gat acc gtc aca gtc ttt ggc ctg cca atg gtt ctg aat gac tac 787

```

Gly	Asp	Thr	Val	Thr	Val	Phe	Gly	Leu	Pro	Met	Val	Leu	Asn	Asp	Tyr	
215						220					225					
tcc	gga	cag	gta	ttc	cca	ccg	ctg	att	gca	gca	att	ggg	ctg	tac	tgg	835
Ser	Gly	Gln	Val	Phe	Pro	Pro	Leu	Ile	Ala	Ala	Ile	Gly	Leu	Tyr	Trp	
230					235				240						245	
gtg	gaa	aag	gga	ctg	aag	aag	atc	atc	cct	gaa	gca	gtc	caa	atg	gtg	883
Val	Glu	Lys	Gly	Leu	Lys	Lys	Ile	Ile	Pro	Glu	Ala	Val	Gln	Met	Val	
				250					255						260	
ttc	gtc	cca	ttc	ttc	tcc	ctg	ctg	att	atg	atc	cca	gcg	acc	gca	ttc	931
Phe	Val	Pro	Phe	Phe	Ser	Leu	Leu	Ile	Met	Ile	Pro	Ala	Thr	Ala	Phe	
			265					270						275		
ctg	ctt	gga	cct	ttc	ggc	atc	ggg	gtt	ggg	aac	gga	att	tcc	aac	ctg	979
Leu	Leu	Gly	Pro	Phe	Gly	Ile	Gly	Val	Gly	Asn	Gly	Ile	Ser	Asn	Leu	
		280					285					290				
ctt	gaa	gcg	att	aac	aac	ttc	agc	cca	ttt	att	ctt	tcc	atc	gtt	atc	1027
Leu	Glu	Ala	Ile	Asn	Asn	Phe	Ser	Pro	Phe	Ile	Leu	Ser	Ile	Val	Ile	
	295					300					305					
cca	ttg	ctc	tac	cca	ttc	ttg	gtt	cca	ctt	gga	ttg	cac	tggt	cca	cta	1075
Pro	Leu	Leu	Tyr	Pro	Phe	Leu	Val	Pro	Leu	Gly	Leu	His	Trp	Pro	Leu	
310					315					320					325	
aac	gcc	atc	atg	atc	cag	aac	atc	aac	acc	ctg	ggg	tac	gac	ttc	att	1123
Asn	Ala	Ile	Met	Ile	Gln	Asn	Ile	Asn	Thr	Leu	Gly	Tyr	Asp	Phe	Ile	
				330					335						340	
cag	gga	cca	atg	ggg	gcc	tggt	aac	ttc	gcc	tgc	ttc	ggc	ctg	gtc	acc	1171
Gln	Gly	Pro	Met	Gly	Ala	Trp	Asn	Phe	Ala	Cys	Phe	Gly	Leu	Val	Thr	
			345					350					355			
ggc	gtg	ttc	ttg	ctc	tcc	att	aag	gaa	cga	aac	aag	gcc	atg	cgt	cag	1219
Gly	Val	Phe	Leu	Leu	Ser	Ile	Lys	Glu	Arg	Asn	Lys	Ala	Met	Arg	Gln	
		360					365					370				
gtt	tcc	ctg	ggg	ggc	atg	ttg	gct	ggg	ttg	ctc	ggc	ggc	att	tcc	gag	1267
Val	Ser	Leu	Gly	Gly	Met	Leu	Ala	Gly	Leu	Leu	Gly	Gly	Ile	Ser	Glu	
	375					380					385					
cct	tcc	ctc	tac	ggg	gtt	ctg	ctc	cga	ttc	aag	aag	acc	tac	ttc	cgc	1315
Pro	Ser	Leu	Tyr	Gly	Val	Leu	Leu	Arg	Phe	Lys	Lys	Thr	Tyr	Phe	Arg	
390					395					400					405	
ctc	ctg	ccg	ggg	tgt	ttg	gca	gca									1339
Leu	Leu	Pro	Gly	Cys	Leu	Ala	Ala									
				410												

&lt;210&gt; 32

&lt;211&gt; 413

&lt;212&gt; PRT

&lt;213&gt; Corynebacterium glutamicum

&lt;400&gt; 32

Met	Ala	Ser	Lys	Leu	Thr	Thr	Thr	Ser	Gln	His	Ile	Leu	Glu	Asn	Leu	
1				5					10					15		



Gly Gly Pro Asp Asn Ile Thr Ser Met Thr His Cys Ala Thr Arg Leu  
                     20                    25                    30  
 Arg Phe Gln Val Lys Asp Gln Ser Ile Val Asp Gln Gln Glu Ile Asp  
                     35                    40                    45  
 Ser Asp Pro Ser Val Leu Gly Val Val Pro Gln Gly Ser Thr Gly Met  
                     50                    55                    60  
 Gln Val Val Met Gly Gly Ser Val Ala Asn Tyr Tyr Gln Glu Ile Leu  
                     65                    70                    75                    80  
 Lys Leu Asp Gly Met Lys His Phe Ala Asp Gly Glu Ala Thr Glu Ser  
                     85                    90                    95  
 Ser Ser Lys Lys Glu Tyr Gly Gly Val Arg Gly Lys Tyr Ser Trp Ile  
                     100                    105                    110  
 Asp Tyr Ala Phe Glu Phe Leu Ser Asp Thr Phe Arg Pro Ile Leu Trp  
                     115                    120                    125  
 Ala Leu Leu Gly Ala Ser Leu Ile Ile Thr Leu Leu Val Leu Ala Asp  
                     130                    135                    140  
 Thr Phe Gly Leu Gln Asp Phe Arg Ala Pro Met Asp Glu Gln Pro Asp  
                     145                    150                    155                    160  
 Thr Tyr Val Phe Leu His Ser Met Trp Arg Ser Val Phe Tyr Phe Leu  
                     165                    170                    175  
 Pro Ile Met Val Gly Ala Thr Ala Ala Arg Lys Leu Gly Ala Asn Glu  
                     180                    185                    190  
 Trp Ile Gly Ala Ala Ile Pro Ala Ala Leu Leu Thr Pro Glu Phe Leu  
                     195                    200                    205  
 Ala Leu Gly Ser Ala Gly Asp Thr Val Thr Val Phe Gly Leu Pro Met  
                     210                    215                    220  
 Val Leu Asn Asp Tyr Ser Gly Gln Val Phe Pro Pro Leu Ile Ala Ala  
                     225                    230                    235                    240  
 Ile Gly Leu Tyr Trp Val Glu Lys Gly Leu Lys Lys Ile Ile Pro Glu  
                     245                    250                    255  
 Ala Val Gln Met Val Phe Val Pro Phe Phe Ser Leu Leu Ile Met Ile  
                     260                    265                    270  
 Pro Ala Thr Ala Phe Leu Leu Gly Pro Phe Gly Ile Gly Val Gly Asn  
                     275                    280                    285  
 Gly Ile Ser Asn Leu Leu Glu Ala Ile Asn Asn Phe Ser Pro Phe Ile  
                     290                    295                    300  
 Leu Ser Ile Val Ile Pro Leu Leu Tyr Pro Phe Leu Val Pro Leu Gly  
                     305                    310                    315                    320  
 Leu His Trp Pro Leu Asn Ala Ile Met Ile Gln Asn Ile Asn Thr Leu  
                     325                    330                    335  
 Gly Tyr Asp Phe Ile Gln Gly Pro Met Gly Ala Trp Asn Phe Ala Cys

340							345					350				
Phe	Gly	Leu	Val	Thr	Gly	Val	Phe	Leu	Leu	Ser	Ile	Lys	Glu	Arg	Asn	
		355					360					365				
Lys	Ala	Met	Arg	Gln	Val	Ser	Leu	Gly	Gly	Met	Leu	Ala	Gly	Leu	Leu	
	370					375					380					
Gly	Gly	Ile	Ser	Glu	Pro	Ser	Leu	Tyr	Gly	Val	Leu	Leu	Arg	Phe	Lys	
385					390					395					400	
Lys	Thr	Tyr	Phe	Arg	Leu	Leu	Pro	Gly	Cys	Leu	Ala	Ala				
				405					410							

```
<210> 33
<211> 428
<212> DNA
<213> Corynebacterium glutamicum
```

```
<220>  
<221> CDS  
<222> (1) .. (405)  
<223> FRXA01943
```

[illegible]

<210> 34  
 <211> 135  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 34  
 Pro Asp Pro Ile Phe Ala Ala Gly Lys Leu Gly Pro Gly Ile Ala Ile  
   1                  5                  10                  15  
 Gln Pro Thr Gly Asn Thr Val Val Ala Pro Ala Asp Ala Thr Val Ile  
           20                  25                  30  
 Leu Val Gln Lys Ser Gly His Ala Val Ala Leu Arg Leu Asp Ser Gly  
       35                  40                  45  
 Val Glu Ile Leu Val His Val Gly Leu Asp Thr Val Gln Leu Gly Gly  
       50                  55                  60  
 Glu Gly Phe Thr Val His Val Glu Arg Arg Gln Gln Val Lys Ala Gly  
       65                  70                  75                  80  
 Asp Pro Leu Ile Thr Phe Asp Ala Asp Phe Ile Arg Ser Lys Asp Leu  
                   85                  90                  95  
 Pro Leu Ile Thr Pro Val Val Val Ser Asn Ala Ala Lys Phe Gly Glu  
           100                  105                  110  
 Ile Glu Gly Ile Pro Ala Asp Gln Ala Asn Ser Ser Thr Thr Val Ile  
       115                  120                  125  
 Lys Val Asn Gly Lys Asn Glu  
       130                  135

**THIS PAGE BLANK (USPTO)**

## SEQUENCE LISTING

<110> BASF Aktiengesellschaft  
 <120> CORYNEBACTERIUM GLUTAMICUM GENES ENCODING  
 PHOSHOENOLPYRUVATE:SUGAR PHOSPHOTRANSFERASE  
 SYSTEM PROTEINS  
 <130> BGT-122CPPC  
 <140>  
 <141>  
 <160> 34  
  
 <210> 1  
 <211> 1527  
 <212> DNA  
 <213> Corynebacterium glutamicum  
  
 <220>  
 <221> CDS  
 <222> (101)..(1504)  
 <223> KXS00315  
  
 <400> 1  
 ctcacggcat ctgcccgtt ccgcttcttg ccagtgttgg ttggtttcac cgcacccaag 60  
  
 cgtttcgccg gcaatgagtt cctggggccc ggcatttggg atg gcc atg gtg ttc 115  
 Met Ala Met Val Phe  
 1 5  
  
 ccg agc lln qtg aac ggc tcc gac gtg gcc gcc acc atg gct gcc qgc 163  
 Pro Ser Leu Val Asn Gly Tyr Asp Val Ala Ala Thr Met Ala Ala Gly  
 10 15 20  
  
 gaa atg cca atg tgg tcc ctg ttt ggt tta gat gtt gcc caa gcc ggt 211  
 Glu Met Pro Met Trp Ser Leu Phe Gly Leu Asp Val Ala Gln Ala Gly  
 25 30 35  
  
 tac cag gcc acc gtg ctt cct gtg ctg gtg gtt tct tgg att ctg gca 259  
 Tyr Gln Gly Thr Val Leu Pro Val Leu Val Val Ser Trp Ile Leu Ala  
 40 45 50  
  
 aag atc gag aag ttc ctg cac aag cga ctc aag gcc act gca gac ttc 307  
 Thr Ile Glu Lys Phe Leu His Lys Arg Leu Lys Gly Thr Ala Asp Phe  
 55 60 65  
  
 ctg atc act cca gtg ctg aag ttg ctg ctc acc gga ttc ctc aca ttc 355  
 Leu Ile Thr Pro Val Leu Thr Leu Leu Leu Thr Gly Phe Leu Thr Phe  
 70 75 80 85  
  
 atc gcc att gcc cca gca atg cgc tgg gtg gcc gat gtg ctg gca cac 403  
 Ile Ala Ile Gly Pro Ala Met Arg Trp Val Gly Asp Val Leu Ala His  
 90 95 100  
  
 ggt cta cag gga ctt tat gat ttc ggt ggt cca gtc gcc ggt ctg ctc 451  
 Gly Leu Gln Gly Leu Tyr Asp Phe Gly Gly Pro Val Gly Gly Leu Leu  
 105 110 115  
  
 ttc ggt ctg gtc tac lca cca atc gtc atc act qgt ctg cag ccg tcc 499  
 Phe Gly Leu Val Tyr Ser Pro Ile Val Ile Thr Gly Leu His Gln Ser  
 120 125 130  
  
 ttc ccg cca att gag ctg gaa ctg ttt aac cag qgt gga tcc ttc ala 547

Phe	Pro	Pro	Ile	Glu	Leu	Glu	Leu	Phe	Asn	Gln	Gly	Gly	Ser	Phe	Ile	
135						140					145					
ttc	gca	acg	gca	tct	atg	ggt	aat	ctc	gcc	cag	ggt	gag	gca	tgt	ttg	595
Phe	Ala	Thr	Ala	Ser	Met	Ala	Asn	Ile	Ala	Gln	Gly	Ala	Ala	Cys	Leu	
150					155					160					165	
gca	gtg	ttc	ttc	ctg	gag	aag	agt	gaa	aag	ctc	aag	gac	ctt	gca	ggt	643
Ala	Val	Phe	Phe	Leu	Ala	Lys	Ser	Glu	Lys	Leu	Lys	Gly	Leu	Ala	Gly	
				170					175					180		
got	tca	ggt	gtc	tcc	gct	gtt	ctt	ggt	att	acg	gag	cct	gag	atc	ttc	691
Ala	Ser	Gly	Val	Ser	Ala	Val	Leu	Gly	Tle	Thr	Glu	Pro	Ala	Ile	Phe	
			185					190					195			
ggt	gtg	aac	ctt	cgc	ctg	cgc	tgg	cag	ttc	ttc	atc	ggt	atc	ggt	acc	739
Gly	Val	Asn	Leu	Arg	Leu	Arg	Trp	Pro	Phe	Phe	Tle	Gly	Ile	Gly	Thr	
		200					205					210				
gca	gct	atc	ggt	ggc	gct	ttg	att	gca	ctc	att	aat	atc	aag	gca	gtt	787
Ala	Ala	Ile	Gly	Gly	Ala	Leu	Tle	Ala	Leu	Phe	Asn	Tle	Lys	Ala	Val	
	215					220					225					
gag	ttg	ggc	gct	gca	ggt	ttc	ttg	ggt	gtt	gtt	tct	att	gat	gct	cca	835
Ala	Leu	Gly	Ala	Ala	Gly	Phe	Leu	Gly	Val	Val	Ser	Ile	Asp	Ala	Pro	
230					235					240					245	
gat	atg	gtc	atg	ttc	ttg	gtg	tgt	gca	gtt	gtt	acc	ttc	ttc	atc	gca	883
Asp	Met	Val	Met	Phe	Leu	Val	Cys	Ala	Val	Val	Thr	Phe	Phe	Ile	Ala	
				250					255					260		
ttc	ggc	gca	ggc	att	gct	tct	gac	att	tac	ttg	gtt	cgc	cgc	aac	ggc	931
Phe	Gly	Ala	Ala	Ile	Ala	Tyr	Gly	Leu	Tyr	Leu	Val	Arg	Arg	Asn	Gly	
			265					270					275			
agc	att	gat	cca	gat	gca	acc	gct	gct	cca	gtg	cct	gca	gga	acg	acc	979
Ser	Ile	Asp	Pro	Asp	Ala	Thr	Ala	Ala	Pro	Val	Pro	Ala	Gly	Thr	Thr	
		280					285					290				
aaa	gcc	gaa	gca	gaa	gca	ccc	gca	gaa	ttt	tca	aac	gat	too	acc	atc	1027
Lys	Ala	Glu	Ala	Glu	Ala	Pro	Ala	Glu	Phe	Ser	Asn	Asp	Ser	Thr	Ile	
	295					300					305					
atc	cag	gca	cct	ttg	acc	ggt	gaa	gct	att	gca	ctg	agc	agc	gtc	agc	1075
Ile	Gln	Ala	Pro	Leu	Thr	Gly	Glu	Ala	Ile	Ala	Leu	Ser	Ser	Val	Ser	
310					315					320					325	
gat	gcc	atg	ttt	gcc	agc	gga	aag	ctt	ggc	tgg	ggc	gtt	gca	atc	gtc	1123
Asp	Ala	Met	Phe	Ala	Ser	Gly	Lys	Leu	Gly	Ser	Gly	Val	Ala	Ile	Val	
				330					335					340		
caa	acc	aag	ggg	caq	tta	gtt	tct	cag	gtg	agt	gga	aag	att	gtg	gtg	1171
Pro	Thr	Lys	Gly	Gln	Leu	Val	Ser	Pro	Val	Ser	Gly	Lys	Ile	Val	Val	
			345					350					355			
gca	ttc	cca	tct	ggc	cat	gct	ttc	gca	gtt	cgc	ccc	aag	gct	gag	gct	1219
Ala	Phe	Pro	Ser	Gly	His	Ala	Phe	Ala	Val	Arg	Thr	Lys	Ala	Glu	Asp	
		360					365					370				
ggt	too	aat	gtg	gat	atc	ttg	atg	caa	att	ggt	ttc	gac	cca	gta	aac	1267
Gly	Ser	Asn	Val	Asp	Tle	Leu	Met	His	Tle	Gly	Phe	Asp	Thr	Val	Asn	

375	380	385	
ctc aac ggc acg cac ttt aac ccg ctg aag aag cag ggc gat gaa gtc			1315
Leu Asn Gly Thr His Phe Asn Pro Leu Lys Lys Gln Gly Asp Glu Val			
390	395	400	405
aaa gcc ggg gag ctg ctg tgt gaa ttc gat att gat gcc att aag gct			1363
Lys Ala Gly Glu Leu Leu Cys Glu Phe Asp Ile Asp Ala Ile Lys Ala			
	410	415	420
gca ggt tat gag gta aac acg ccg att gtt gtt tog ant tnc aag aaa			1411
Ala Gly Tyr Glu Val Thr Thr Pro Ile Val Val Ser Asn Tyr Lys Lys			
	425	430	435
acc gga cct gta aac act tac ggt ttg gcc gaa att gaa gcc gga gcc			1459
Thr Gly Pro Val Asn Thr Tyr Gly Leu Gly Glu Ile Glu Ala Gly Ala			
	440	445	450
aac ctg ctc aac gtc gca aag aaa gaa gcc gtg cca gca aca cca			1504
Asn Leu Leu Asn Val Ala Lys Lys Glu Ala Val Pro Ala Thr Pro			
	455	460	465
taagttgaan ccttgagtggt tog			1527

&lt;210&gt; 2

&lt;211&gt; 468

&lt;212&gt; ERT

&lt;213&gt; Corynebacterium glutamicum

&lt;400&gt; 2

Met Ala Met Val Phe Pro Ser Leu Val Asn Gly Tyr Asp Val Ala Ala  
1 5 10 15

Thr Met Ala Ala Gly Glu Met Pro Met Trp Ser Leu Phe Gly Leu Asp  
20 25 30

Val Ala Gln Ala Gly Tyr Gln Gly Thr Val Leu Pro Val Leu Val Val  
35 40 45

Ser Trp Ile Leu Ala Thr Ile Glu Lys Phe Leu His Lys Arg Leu Lys  
50 55 60

Gly Thr Ala Asp Phe Leu Ile Thr Pro Val Leu Thr Leu Leu Thr  
65 70 75 80

Gly Phe Leu Thr Phe Ile Ala Ile Gly Pro Ala Met Arg Trp Val Gly  
85 90 95

Asp Val Leu Ala His Gly Leu Gln Gly Leu Tyr Asp Phe Gly Gly Pro  
100 105 110

Val Gly Gly Leu Leu Phe Gly Leu Val Tyr Ser Pro Ile Val Ile Thr  
115 120 125

Gly Leu His Gln Ser Phe Pro Pro Ile Glu Leu Glu Leu Phe Asn Gln  
130 135 140

Gly Gly Ser Phe Ile Phe Ala Thr Ala S r Met Ala Asn Ile Ala Gln  
145 150 155 160

Gly Ala Ala Cys Leu Ala Val Phe Phe Leu Ala Lys Ser Glu Lys Leu  
 165 170 175  
 Lys Gly Leu Ala Gly Ala Ser Gly Val Ser Ala Val Leu Gly Ile Thr  
 180 185 190  
 Glu Pro Ala Ile Phe Gly Val Asn Leu Arg Leu Arg Trp Pro Phe Phe  
 195 200 205  
 Ile Gly Ile Gly Thr Ala Ala Ile Gly Gly Ala Leu Ile Ala Leu Phe  
 210 215 220  
 Asn Ile Lys Ala Val Ala Leu Gly Ala Ala Gly Phe Leu Gly Val Val  
 225 230 235 240  
 Ser Ile Asp Ala Pro Asp Met Val Met Phe Leu Val Cys Ala Val Val  
 245 250 255  
 Thr Phe Phe Ile Ala Phe Gly Ala Ala Ile Ala Tyr Gly Leu Tyr Leu  
 260 265 270  
 Val Arg Arg Asn Gly Ser Ile Asp Pro Asp Ala Thr Ala Ala Pro Val  
 275 280 285  
 Pro Ala Gly Thr Thr Lys Ala Glu Ala Glu Ala Pro Ala Glu Phe Ser  
 290 295 300  
 Asn Asp Ser Thr Ile Ile Gln Ala Pro Leu Thr Gly Glu Ala Ile Ala  
 305 310 315 320  
 Leu Ser Ser Val Ser Asp Ala Met Phe Ala Ser Gly Lys Leu Gly Ser  
 325 330 335  
 Gly Val Ala Ile Val Pro Thr Lys Gly Gln Leu Val Ser Pro Val Ser  
 340 345 350  
 Gly Lys Ile Val Val Ala Phe Pro Ser Gly His Ala Phe Ala Val Arg  
 355 360 365  
 Thr Lys Ala Glu Asp Gly Ser Asn Val Asp Ile Leu Met His Ile Gly  
 370 375 380  
 Phe Asp Thr Val Asn Leu Asn Gly Thr His Phe Asn Pro Leu Lys Lys  
 385 390 395 400  
 Gln Gly Asp Glu Val Lys Ala Gly Glu Leu Leu Cys Glu Phe Asp Ile  
 405 410 415  
 Asp Ala Ile Lys Ala Ala Gly Tyr Glu Val Thr Thr Pro Ile Val Val  
 420 425 430  
 Ser Asn Tyr Lys Lys Thr Gly Pro Val Asn Thr Tyr Gly Leu Gly Glu  
 435 440 445  
 Ile Glu Ala Gly Ala Asn Leu Leu Asn Val Ala Lys Lys Glu Ala Val  
 450 455 460  
 Pro Ala Thr Pro  
 465



<210> 3  
 <211> 1109  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (1)...(1086)  
 <223> FRXA00315

<400> 3  
 tat gat ttc ggc ggt cca gtc ggc ggt ctg ctg ttc ggt ctg gtc tac 48  
 Tyr Asp Phe Gly Gly Pro Val Gly Gly Leu Leu Phe Gly Leu Val Tyr  
 1 5 10 15  
 tca cca atc gtc atc act ggt ctg cac cag tcc ttc ccg cca att gag 96  
 Ser Pro Ile Val Ile Thr Gly Leu His Gln Ser Phe Pro Ile Glu  
 20 25 30  
 ctg gag ctg ttt aac cag ggt gga tcc ttc aac ttc gca acg gca tct 144  
 Leu Glu Leu Phe Asn Gln Gly Gly Ser Phe Ile Phe Ala Thr Ala Ser  
 35 40 45  
 atg gct aat atc gcc cag ggt gcg gca tgt ttg gca glg ttc ttc ctg 192  
 Met Ala Asn Ile Ala Gln Gly Ala Ala Cys Leu Ala Val Phe Phe Leu  
 50 55 60  
 gcg aag agt gaa aag ctg aag ggc ctt gca ggt gct tca ggt gtc tcc 240  
 Ala Lys Ser Glu Lys Leu Lys Gly Leu Ala Gly Ala Ser Gly Val Ser  
 65 70 75 80  
 gct gtt ctt ggt att acg gag cct gcg atc ttc ggt gtg aac ctt cgc 288  
 Ala Val Leu Gly Ile Thr Glu Pro Ala Ile Phe Gly Val Asn Leu Arg  
 85 90 95  
 ctg cgc tgg ccg ttc ttc atc ggt atc ggt acc gca got atc ggt ggc 336  
 Leu Arg Trp Pro Phe Phe Ile Gly Ile Gly Thr Ala Ala Ile Gly Gly  
 100 105 110  
 gcl llg all gca ctg ttc aat atc aag gca gtt gca ttg ggc got gca 384  
 Ala Leu Ile Ala Leu Phe Asn Ile Lys Ala Val Ala Leu Gly Ala Ala  
 115 120 125  
 ggt ttc tcc ggt gtt gtt tct att gat got cca gat atg gtc atg ttc 432  
 Gly Phe Leu Gly Val Val Ser Ile Asp Ala Pro Asp Met Val Met Phe  
 130 135 140  
 ttg gtg tgt gca gtt gtt acc ttc ttc aac gca ttc ggc gca gcg att 480  
 Leu Val Cys Ala Val Val Thr Phe Phe Ile Ala Phe Gly Ala Ala Ile  
 145 150 155 160  
 got tat ggc ctt tca ttg gtt cgc cgc aac ggc agc att gat cca gat 528  
 Ala Tyr Gly Leu Tyr Leu Val Arg Arg Asn Gly Ser Ile Asp Pro Asp  
 165 170 175  
 gca acc got got cca gtg cct gca gga acg acc aaa gcc gaa gca gaa 576  
 Ala Thr Ala Ala Pro Val Pro Ala Gly Thr Thr Lys Ala Glu Ala Glu  
 180 185 190  
 gca ccc gca gaa ttt tca aac gat tcc acc ctg ctg ccg gca cct ttg 624  
 Ala Pro Ala Glu Phe Ser Asn Asp Ser Thr Ile Ile Gln Ala Pro Leu

195					200					205					
acc ggt gaa gct att gca ctg agc agc gtc agc gat gcc atg ttt gcc	672														
Thr Gly Glu Ala Ile Ala Leu Ser Ser Val Ser Asp Ala Met Phe Ala															
210 215 220															
agc gga aag ctt ggc tgg ggc gtt gcc atc gtc cca acc aag ggg cag	720														
Ser Gly Lys Leu Gly Ser Gly Val Ala Ile Val Pro Thr Lys Gly Gln															
225 230 235 240															
tta gtt tct ccg gtg agt gga aag att gtg gtg gca ttc cca tct ggc	768														
Leu Val Ser Pro Val Ser Gly Lys Ile Val Val Ala Phe Pro Ser Gly															
245 250 255															
cat got ttc gca gtt cgc acc aag gct gag gat ggt tcc aat gtg gat	816														
Ris Ala Phe Ala Val Arg Thr Lys Ala Glu Asp Gly Ser Asn Val Asp															
260 265 270															
atc ttg atg cac att ggt ttc gac aca gta aac ctc aac ggc acg cac	864														
Ile Leu Met His Ile Gly Phe Asp Thr Val Asn Leu Asn Gly Thr His															
275 280 285															
ttt aac ccg ctg aag aag cag ggc gat gaa gtc aaa gca ggg gag ctg	912														
Phe Asn Pro Leu Lys Lys Gln Gly Asp Glu Val Lys Ala Gly Glu Leu															
290 295 300															
ctg tgt gaa ttc gat att gat gcc att aag gct gca ggt tat gag gta	960														
Leu Cys Glu Phe Asp Ile Asp Ala Ile Lys Ala Ala Gly Tyr Glu Val															
305 310 315 320															
acc acg ccg att gtt gtt tgg aat tac aaq aac acc gga cct gta aac	1008														
Thr Thr Pro Ile Val Val Ser Asn Tyr Lys Lys Thr Gly Pro Val Asn															
325 330 335															
act tac ggt ttg ggc gaa att gaa ggc ggc gcc aac ctg ctg aac gtc	1056														
Thr Tyr Gly Leu Gly Glu Ile Glu Ala Gly Ala Asn Leu Leu Asn Val															
340 345 350															
gca aag aaa gaa ggt gtc cca gca aca cca caagctgaa ccttgagctc	1106														
Ala Lys Lys Glu Ala Val Pro Ala Thr Pro															
355 360															
tgg	1109														

<210> 4  
 <211> 362  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 4  
 Tyr Asp Phe Gly Gly Pro Val Gly Gly Leu Leu Phe Gly Leu Val Tyr  
 1 5 10 15  
 Ser Pro Ile Val Ile Thr Gly Leu His Gln Ser Phe Pro Pro Ile Glu  
 20 25 30  
 Leu Glu Leu Phe Asn Gln Gly Gly Ser Phe Ile Phe Ala Thr Ala Ser  
 35 40 45  
 Met Ala Asn Ile Ala Gln Gly Ala Ala Cys Leu Ala Val Phe Phe Leu

50	55	60
Ala Lys Ser Glu Lys Leu Lys Gly Leu Ala Gly Ala Ser Gly Val Ser 65 70 75 80		
Ala Val Leu Gly Ile Thr Glu Pro Ala Ile Phe Gly Val Asn Leu Arg 85 90 95		
Leu Arg Trp Pro Phe Phe Ile Gly Ile Gly Thr Ala Ala Ile Gly Gly 100 105 110		
Ala Leu Ile Ala Leu Phe Asn Ile Lys Ala Val Ala Leu Gly Ala Ala 115 120 125		
Gly Phe Leu Gly Val Val Ser Ile Asp Ala Pro Asp Met Val Met Phe 130 135 140		
Leu Val Cys Ala Val Val Thr Phe Phe Ile Ala Phe Gly Ala Ala Ile 145 150 155 160		
Ala Tyr Gly Leu Tyr Leu Val Arg Arg Asn Gly Ser Ile Asp Pro Asp 165 170 175		
Ala Thr Ala Ala Pro Val Pro Ala Gly Thr Thr Lys Ala Glu Ala Glu 180 185 190		
Ala Pro Ala Glu Phe Ser Asn Asp Ser Thr Ile Ile Gln Ala Pro Leu 195 200 205		
Thr Gly Glu Ala Ile Ala Leu Ser Ser Val Ser Asp Ala Met Phe Ala 210 215 220		
Ser Gly Lys Leu Gly Ser Gly Val Ala Ile Val Pro Thr Lys Gly Gln 225 230 235 240		
Leu Val Ser Pro Val Ser Gly Lys Ile Val Val Ala Phe Pro Ser Gly 245 250 255		
His Ala Phe Ala Val Arg Thr Lys Ala Glu Asp Gly Ser Asn Val Asp 260 265 270		
Ile Leu Met His Ile Gly Phe Asp Thr Val Asn Leu Asn Gly Thr His 275 280 285		
Phe Asn Pro Leu Lys Lys Gln Gly Asp Glu Val Lys Ala Gly Glu Leu 290 295 300		
Leu Cys Glu Phe Asp Ile Asp Ala Ile Lys Ala Ala Gly Tyr Glu Val 305 310 315 320		
Thr Thr Pro Ile Val Val Ser Asn Tyr Lys Lys Thr Gly Pro Val Asn 325 330 335		
Thr Tyr Gly Leu Gly Glu Ile Glu Ala Gly Ala Asn Leu Leu Asn Val 340 345 350		
Ala Lys Lys Glu Ala Val Pro Ala Thr Pro 355 360		

&lt;210&gt; 5

<211> 372  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (101)..(349)  
 <223> RXA01503

<400> 5  
 gtatctctcga aggcctttcta gctgttgacag ctgcagcgca ctccggtggat acgacatccc 60  
 cgacctatca aattcttttat gctgcaggcg atgccttttc atg ttc ttg gca gtc 115  
 Met Phe Leu Ala Val  
 1 5  
 att ttg gcg att act gcg gct cgt aaa ttc ggt gcc aat gtc ttt aca 163  
 Ile Leu Ala Ile Thr Ala Ala Arg Lys Phe Gly Ala Asn Val Phe Thr  
 10 15 20  
 tca gtc gca ctc gct ggt gca ttg ctg cac aca cag ctc cag gca gta 211  
 Ser Val Ala Leu Ala Gly Ala Leu Leu His Thr Gln Leu Gln Ala Val  
 25 30 35  
 acc gtg ttg gtt gac ggt gaa ctc cag tgc atg act ctg gtg gct ttc 259  
 Thr Val Leu Val Asp Gly Glu Leu Gln Ser Met Thr Leu Val Ala Phe  
 40 45 50  
 cna aag gct ugt aat gac gtc acc ttc ctg gcc att cca gtg gtg ctg 307  
 Gln Lys Ala Gly Asn Asp Val Thr Phe Leu Gly Ile Pro Val Val Leu  
 55 60 65  
 cag ttg gcg ttg cat gla gcg agt ttg alu aaq ttg lcu cga 349  
 Gln Leu Ala Leu His Val Ala Ser Leu Met Lys Leu Ser Arg  
 70 75 80  
 taagaggagg ggcgtgtcgg tct 372

<210> 6  
 <211> 83  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 6  
 M t Phe Leu Ala Val Ile Leu Ala Ile Thr Ala Ala Arg Lys Phe Gly  
 1 5 10 15  
 Ala Asn Val Phe Thr Ser Val Ala Leu Ala Gly Ala Leu Leu His Thr  
 20 25 30  
 Gln Leu Gln Ala Val Thr Val Leu Val Asp Gly Glu Leu Gln Ser Met  
 35 40 45  
 Thr Leu Val Ala Phe Gln Lys Ala Gly Asn Asp Val Thr Phe Leu Gly  
 50 55 60  
 Ile Pro Val Val Leu Gln Leu Ala Leu His Val Ala Ser Leu Met Lys  
 65 70 75 80  
 Leu Ser Arg

<210> 7  
 <211> 2187  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (101)..(2164)  
 <223> RXN01299

<400> 7  
 cgaactggggc gtctcttctct ggcctctccca ttcctcgtctc tgaccaactc gccacagctg 60  
 gtgcaacgggt caccacaagtc aaaggattga aagaatcagc atg aat agc gta aat 115  
 Met Asn Ser Val Asn  
 7 5  
 aat tcc tog ott gtc cgg ctg gat gtc gat ttc ggc gac tcc acc acg 163  
 Asn Ser Ser Leu Val Arg Leu Asp Val Asp Phe Gly Asp Ser Thr Thr  
 10 15 20  
 gat gtc atc aac aac ott gcc act gtt att ttc gac gct ggc cga gct 211  
 Asp Val Ile Asn Asn Leu Ala Thr Val Ile Phe Asp Ala Gly Arg Ala  
 25 30 35  
 tcc tcc gcc gac gcc ott gcc aac gac ggc ctg gat cgt gaa gca aag 259  
 Ser Ser Ala Asp Ala Leu Ala Lys Asp Ala Leu Asp Arg Glu Ala Lys  
 40 45 50  
 tcc ggc acc ggc gtt cct ggt caa gtt gct atc ccc caa tgc cgt tcc 307  
 Ser Gly Thr Gly Val Pro Gly Gln Val Ala Ile Pro His Cys Arg Ser  
 55 60 65  
 gaa gcc gta tct gtc cct acc ttg ggc ttt gct cgc ctg agc aag ggt 355  
 Glu Ala Val Ser Val Pro Thr Leu Gly Phe Ala Arg Leu Ser Lys Gly  
 70 75 80 85  
 gtg gac ttc agc gga cct gat ggc gat gcc aac ttg gtg ttc ctc att 403  
 Val Asp Phe Ser Gly Pro Asp Gly Asp Ala Asn Leu Val Phe Leu Ile  
 90 95 100  
 gca gca cct gct ggc ggc ggc aac gag cac ctg aag atc ctg tcc aag 451  
 Ala Ala Pro Ala Gly Gly Gly Lys Glu His Leu Lys Ile Leu Ser Lys  
 105 110 115  
 ctt gct cgc tcc ttg gtg aag aag gat ttc atc aag gct ctg cag gaa 499  
 Leu Ala Arg Ser Leu Val Lys Lys Asp Phe Ile Lys Ala Leu Gln Glu  
 120 125 130  
 gcc acc acc ggc cag gaa atc gtc gac gtt gtc gat gcc gtg ctg acc 547  
 Ala Thr Thr Glu Glu Glu Ile Val Asp Val Val Asp Ala Val Leu Asn  
 135 140 145  
 cca gca cca aaa acc acc gag cca gct gca gct ccg gct ggc ggc ggc 595  
 Pro Ala Pro Lys Thr Thr Glu Pro Ala Ala Ala Pro Ala Ala Ala Ala  
 150 155 160 165  
 gtt gct gag agt ggg ggc ggc tgg aca agc gtt act cgt atc gtg gca 643

Val	Ala	Glu	Ser	Gly	Ala	Ala	Ser	Thr	Ser	Val	Thr	Arg	Ile	Val	Ala		
				170					175					180			
etc	acc	gca	tgc	caa	acc	ggt	atc	gca	cac	acc	tac	atg	gct	gcg	gat	691	
Ile	Thr	Ala	Cys	Pro	Thr	Gly	Ile	Ala	His	Thr	Tyr	Met	Ala	Ala	Asp		
			185					190					195				
tcg	ctg	acc	caa	aac	gcg	gaa	ggc	cgc	gat	gat	gtg	gaa	ctc	gtt	glg	739	
Ser	Leu	Thr	Gln	Asn	Ala	Glu	Gly	Arg	Asp	Asp	Val	Glu	Leu	Val	Val		
			200				205					210					
gag	act	cag	ggc	tct	tcc	gct	gtc	acc	cca	gtc	gct	ccg	aag	alc	atc	787	
Glu	Thr	Gln	Gly	Ser	Ser	Ala	Val	Thr	Pro	Val	Asp	Pro	Lys	Ile	Ile		
	215					220					225						
gga	gcl	gcc	gac	gcc	gtc	atc	llo	gcc	acc	gac	glg	gga	gtt	aac	gac	835	
Glu	Ala	Ala	Asp	Ala	Val	Ile	Phe	Ala	Thr	Asp	Val	Gly	Val	Lys	Asp		
230					235					240				245			
cgc	gag	cgt	ttc	gct	ggc	aag	cca	gtc	att	gaa	lnc	ggc	gtc	aag	cgc	883	
Arg	Glu	Arg	Phe	Ala	Gly	Lys	Pro	Val	Ile	Glu	Ser	Gly	Val	Lys	Arg		
			250					255						260			
gag	atc	aal	gag	cca	ggc	aag	atg	atc	gac	gag	gcc	alc	gca	gcc	lcc	931	
Ala	Ile	Asn	Glu	Pro	Ala	Lys	Met	Ile	Asp	Glu	Ala	Ile	Ala	Ala	Ser		
			265				270						275				
aag	aac	cca	aac	gcc	cgc	aag	gtt	tcc	ggt	tcc	ggt	gtc	gcg	gca	tct	979	
Lys	Asn	Pro	Asn	Ala	Arg	Lys	Val	Ser	Gly	Ser	Gly	Val	Ala	Ala	Ser		
		280					285					290					
gct	gaa	acc	acc	ggc	gag	aag	ctc	ggc	tgg	ggc	aag	cgc	atc	cag	cag	1027	
Ala	Glu	Thr	Thr	Gly	Glu	Lys	Leu	Gly	Trp	Gly	Lys	Arg	Ile	Gln	Gln		
	295					300					305						
gca	gtc	atg	acc	ggc	gtg	tcc	tac	atg	gtt	cca	ttc	gta	gct	gcc	ggc	1075	
Ala	Val	Met	Thr	Gly	Val	Ser	Tyr	Met	Val	Pro	Phe	Val	Ala	Ala	Gly		
310					315					320					325		
ggc	ctc	ctg	ttg	gct	ctc	ggc	ttc	gca	ttc	ggt	gga	tac	gac	atg	gcg	1123	
Gly	Leu	Leu	Leu	Ala	Leu	Gly	Phe	Ala	Phe	Gly	Gly	Tyr	Asp	Met	Ala		
				330				335						340			
aac	ggc	tgg	caa	gca	atc	gcc	acc	cag	ttc	tct	ctg	acc	aac	ctg	cca	1171	
Asn	Gly	Trp	Gln	Ala	Tic	Ala	Thr	Gln	Phe	Ser	Leu	Thr	Asn	Leu	Pro		
			345					350					355				
ggc	aac	acc	gtc	gat	gtt	gac	ggc	gtg	gcc	atg	acc	ttc	gag	cgt	tca	1219	
Gly	Asn	Thr	Val	Asp	Val	Asp	Gly	Val	Ala	Met	Thr	Phe	Glu	Arg	Ser		
		360				365						370					
ggc	ttc	ctg	ttg	tac	ttc	ggc	gca	gtc	ctg	ttc	gcc	acc	ggc	caa	gca	1267	
Gly	Phe	Leu	Leu	Tyr	Phe	Gly	Ala	Val	Leu	Phe	Ala	Thr	Gly	Gln	Ala		
	375					380					385						
gcc	atg	ggc	ttc	atc	gtg	gca	gcc	ctg	tct	ggc	tac	acc	gca	tac	gca	1315	
Ala	Met	Gly	Phe	Ile	Val	Ala	Ala	Leu	Ser	Gly	Tyr	Thr	Ala	Tyr	Ala		
	390				395					400					405		
ctc	gct	gga	cgc	cca	ggc	atc	gca	cag	ggc	ttc	gtc	ggt	ggc	gcc	alc	1363	
Leu	Ala	Gly	Arg	Pro	Gly	Il	Ala	Pro	Gly	Phe	Val	Gly	Gly	Ala	Ile		

410			415			420			
tcc gtc acc atc ggc gct ggc ttc att ggt ggt ctg gtt acc ggt atc	1411								
Ser Val Thr Ile Gly Ala Gly Phe Ile Gly Gly Leu Val Thr Gly Ile									
425	430			435					
ttg gct ggt ctc att gcc ctg tgg att ggc tcc tgg aag gta cca cgc	1459								
Leu Ala Gly Leu Ile Ala Leu Trp Ile Gly Ser Trp Lys Val Pro Arg									
440	445			450					
gtg gta cag tca ctg atg cct gta gta atc atc cca cta ctt acc tca	1507								
Val Val Gln Ser Leu Met Pro Val Val Ile Ile Pro Leu Leu Thr Ser									
455	460			465					
gtg gtt gtt ggt ctc gtc atg tac ctc ctg ctg ggt cgc cca ctc gca	1555								
Val Val Val Gly Leu Val Met Tyr Leu Leu Leu Gly Arg Pro Leu Ala									
470	475			480			485		
tcc atc atg act ggt ttg cag gac tgg cta tgg tcc atg tcc gga agc	1603								
Ser Ile Met Thr Gly Leu Gln Asp Trp Leu Ser Ser Met Ser Gly Ser									
490	495			500					
tcc gcc atc ttg ctg ggt atc atc ttg ggc ctc atg atg tgt ttc gac	1651								
Ser Ala Ile Leu Leu Gly Ile Ile Leu Gly Leu Met Met Cys Phe Asp									
505	510			515					
ctc ggc gga cca gta aac aag gca gcc tac ctc ttt ggt acc gca ggc	1699								
Leu Gly Gly Pro Val Asn Lys Ala Ala Tyr Leu Phe Gly Thr Ala Gly									
520	525			530					
ctg tct acc ggc gac caa gct tcc atg gaa atc atg gcc gcg atc atg	1747								
Leu Ser Thr Gly Asp Gln Ala Ser Met Glu Ile Met Ala Ala Ile Met									
535	540			545					
gca gct ggc atg gtc cca cca atc gcg ttg tcc att gct acc ctg ctg	1795								
Ala Ala Gly Met Val Pro Pro Ile Ala Leu Ser Ile Ala Thr Leu Leu									
550	555			560			565		
cgc aag aag ctg ttc acc cca gca gag caa gaa aac ggc aag tct tcc	1843								
Arg Lys Lys Leu Phe Thr Pro Ala Glu Gln Glu Asn Gly Lys Ser Ser									
570	575			580					
tgg ctg ctt ggc ctg gca ttc gtc tcc gaa ggt gcc atc cca ttc gcc	1891								
Trp Leu Leu Gly Leu Ala Phe Val Ser Glu Gly Ala Ile Pro Phe Ala									
585	590			595					
gca gct gac cca ttc cgt gtg atc cca gca atg atg gct ggc ggt gca	1939								
Ala Ala Asp Pro Phe Arg Val Ile Pro Ala Met Met Ala Gly Gly Ala									
600	605			610					
acc act ggt gca atc tcc atg gca ctg ggc gtc ggc tct cgg gct cca	1987								
Thr Thr Gly Ala Ile Ser Met Ala Leu Gly Val Gly Ser Arg Ala Pro									
615	620			625					
cac gcc ggt atc ttc gtg gta tgg gca atc gaa cca tgg tgg gcc tgg	2035								
Ile Gly Gly Ile Phe Val Val Trp Ala Ile Glu Pro Trp Trp Gly Trp									
630	635			640			645		
ctc atc gca ctt gcc gca gcc acc atc gtg tcc acc atc gtt gta atc	2083								
Leu Ile Ala Leu Ala Ala Gly Thr Ile Val Ser Thr Ile Val Val Ile									
650	655			660					

gca ctg aag cag ttc tgg cca aac aag gcc gtc gct gca gaa gtc ggc 2131  
 Ala Leu Lys Gln Phe Trp Pro Asn Lys Ala Val Ala Ala Glu Val Ala  
 665 670 675

aag caa gaa gca caa caa gca got gta aac gca taatgggccc ttgacccgat 2184  
 Lys Gln Glu Ala Gln Gln Ala Ala Val Asn Ala  
 680 685

gtc 2187

```
<210> 8
<211> 688
<212> PRT
<213> Corynebacterium glutamicum
```

```

<400> 8
Met Asn Ser Val Asn Asn Ser Ser Leu Val Arg Leu Asp Val Asp Phe
  1          5          10          15
Gly Asp Ser Thr Thr Asp Val Ile Asn Asn Leu Ala Thr Val Ile Phe
          20          25          30
Asp Ala Gly Arg Ala Ser Ser Ala Asp Ala Leu Ala Lys Asp Ala Leu
          35          40          45
Asp Arg Glu Ala Lys Ser Gly Thr Gly Val Pro Gly Gln Val Ala Ile
          50          55          60
Pro His Cys Arg Ser Glu Ala Val Ser Val Pro Thr Leu Gly Phe Ala
          65          70          75          80
Arg Leu Ser Lys Gly Val Asp Phe Ser Gly Pro Asp Gly Asp Ala Asn
          85          90          95
Leu Val Phe Leu Ile Ala Ala Pro Ala Gly Gly Gly Lys Gln His Leu
          100          105          110
Lys Ile Leu Ser Lys Leu Ala Arg Ser Leu Val Lys Lys Asp Phe Ile
          115          120          125
Lys Ala Leu Gln Glu Ala Thr Thr Glu Gln Glu Ile Val Asp Val Val
          130          135          140
Asp Ala Val Leu Asn Pro Ala Pro Lys Thr Thr Glu Pro Ala Ala Ala
          145          150          155          160
Pro Ala Ala Ala Ala Val Ala Glu Ser Gly Ala Ala Ser Thr Ser Val
          165          170          175
Thr Arg Ile Val Ala Ile Thr Ala Cys Pro Thr Gly Ile Ala His Thr
          180          185          190
Tyr Met Ala Ala Asp Ser Leu Thr Gln Asn Ala Glu Gly Arg Asp Asp
          195          200          205
Val Glu Leu Val Val Glu Thr Gln Gly Ser Ser Ala Val Thr Pro Val
          210          215          220
Asp Pro Lys Ile Il Glu Ala Ala Asp Ala Val Ile Phe Ala Thr Asp

```



225	230	235	240
Val Gly Val Lys Asp Arg Glu Arg Phe Ala Gly Lys Pro Val Ile Glu	245	250	255
Ser Gly Val Lys Arg Ala Ile Asn Glu Pro Ala Lys Met Ile Asp Glu	260	265	270
Ala Ile Ala Ala Ser Lys Asn Pro Asn Ala Arg Lys Val Ser Gly Ser	275	280	285
Gly Val Ala Ala Ser Ala Glu Thr Thr Gly Glu Lys Leu Gly Trp Gly	290	295	300
Lys Arg Ile Gln Gln Ala Val Met Thr Gly Val Ser Tyr Met Val Pro	305	310	320
Phe Val Ala Ala Gly Gly Leu Leu Leu Ala Leu Gly Phe Ala Phe Gly	325	330	335
Gly Tyr Asp Met Ala Asn Gly Trp Gln Ala Ile Ala Thr Gln Phe Ser	340	345	350
Leu Thr Asn Leu Pro Gly Asn Thr Val Asp Val Asp Gly Val Ala Met	355	360	365
Thr Phe Gln Arg Ser Gly Phe Leu Leu Tyr Phe Gly Ala Val Leu Phe	370	375	380
Ala Thr Gly Gln Ala Ala Met Gly Phe Ile Val Ala Ala Leu Ser Gly	385	390	395
Tyr Thr Ala Tyr Ala Leu Ala Gly Arg Pro Gly Ile Ala Pro Gly Phe	405	410	415
Val Gly Gly Ala Ile Ser Val Thr Ile Gly Ala Gly Phe Ile Gly Gly	420	425	430
Leu Val Thr Gly Ile Leu Ala Gly Leu Ile Ala Leu Trp Ile Gly Ser	435	440	445
Trp Lys Val Pro Arg Val Val Gln Ser Leu Met Pro Val Val Ile Ile	450	455	460
Pro Leu Leu Thr Ser Val Val Val Gly Leu Val Met Tyr Leu Leu Leu	465	470	475
Gly Arg Pro Leu Ala Ser Ile Met Thr Gly Leu Gln Asp Trp Leu Ser	485	490	495
Ser Met Ser Gly Ser Ser Ala Ile Leu Leu Gly Ile Ile Leu Gly Leu	500	505	510
Met Met Cys Phe Asp Leu Gly Gly Pro Val Asn Lys Ala Ala Tyr Leu	515	520	525
Phe Gly Thr Ala Gly Leu Ser Thr Gly Asp Gln Ala Ser Met Glu Ile	530	535	540
Met Ala Ala Ile Met Ala Ala Gly Met Val Pro Pro Ile Ala Leu Ser	545	550	555
			560

Ile Ala Thr Leu Leu Arg Lys Lys Leu Ph Thr Pro Ala Glu Gln Gly  
 565 570 575  
 Asn Gly Lys Ser Ser Trp Leu Leu Gly Leu Ala Phe Val Ser Glu Gly  
 580 585 590  
 Ala Ile Pro Phe Ala Ala Ala Asp Pro Phe Arg Val Ile Pro Ala Met  
 595 600 605  
 Met Ala Gly Gly Ala Thr Thr Gly Ala Ile Ser Met Ala Leu Gly Val  
 610 615 620  
 Gly Ser Arg Ala Pro His Gly Gly Ile Phe Val Val Trp Ala Ile Glu  
 625 630 635 640  
 Pro Trp Trp Gly Trp Leu Ile Ala Leu Ala Ala Gly Thr Ile Val Ser  
 645 650 655  
 Thr Ile Val Val Ile Ala Leu Lys Gln Phe Trp Pro Asn Lys Ala Val  
 660 665 670  
 Ala Ala Glu Val Ala Lys Gln Glu Ala Glu Gln Ala Ala Val Asn Ala  
 675 680 685

<210> 9  
 <211> 464  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (1)..(441)  
 <223> PRXA01299

<400> 9  
 atg gaa atc atg gcc gcc atc atc gca gct gcc atg gtc cca cca atc 48  
 Met Glu Ile Met Ala Ala Ile Met Ala Ala Gly Met Val Pro Pro Ile  
 1 5 10 15  
 gcc ttg tcc att gct acc ctg ctg cgc aag aag ctg ttc acc cca gca 96  
 Ala Leu Ser Ile Ala Thr Leu Leu Arg Lys Lys Leu Phe Thr Pro Ala  
 20 25 30  
 aag cca gaa aac ggc aag tat tcc tgg ctg ctt gcc ctg gca ttc gtc 144  
 Glu Gln Glu Asn Gly Lys Ser Ser Trp Leu Leu Gly Leu Ala Phe Val  
 35 40 45  
 tcc gaa ggt gcc atc cca ttc gcc gcc gct gac cca ttc cgt gtg atc 192  
 Ser Glu Gly Ala Ile Pro Phe Ala Ala Ala Asp Pro Phe Arg Val Ile  
 50 55 60  
 cca gca atg atg gct gcc ggt gca acc act ggt gca atc tcc atg gca 240  
 Pro Ala Met M t Ala Gly Gly Ala Thr Thr Gly Ala Ile Ser Met Ala  
 65 70 75 80  
 ctg gcc gtc gcc tat cgg gct cca cac gcc ggt atc ttc gtg gtc tgg 288

Leu Gly Val Gly Ser Arg Ala Pro His Gly Gly Ile Phe Val Val Trp  
                             85                            90                            95  
 gca atc gaa cca tgg tgg ggc tgg ctc atc gca ctt gca gca ggc acc 336  
 Ala Ile Glu Pro Trp Trp Gly Trp Leu Ile Ala Leu Ala Ala Gly Thr  
                             100                            105                            110  
 atc gtg tcc acc atc gtt gtc atc gca ctg aag caa ttc tgg cca aac 384  
 Ile Val Ser Thr Ile Val Val Ile Ala Leu Lys Gln Phe Trp Pro Asn  
                             115                            120                            125  
 aag gcc gtc gct gca gaa gtc gcg aag caa gaa gca caa cca gca gct 432  
 Lys Ala Val Ala Ala Glu Val Ala Lys Gln Glu Ala Gln Gln Ala Ala  
                             130                            135                            140  
 gta aac gca taatcggacc ttgaccgat gtc 464  
 Val Asn Ala  
 145

<210> 10  
 <211> 147  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 10  
 Met Glu Ile Met Ala Ala Ile Met Ala Ala Gly Met Val Pro Pro Ile  
   1                            5                            10                            15  
 Ala Leu Ser Ile Ala Thr Leu Leu Arg Lys Lys Leu Phe Thr Pro Ala  
                             20                            25                            30  
 Glu Gln Glu Asn Gly Lys Ser Ser Trp Leu Leu Gly Leu Ala Phe Val  
                             35                            40                            45  
 Ser Glu Gly Ala Ile Pro Phe Ala Ala Ala Asp Pro Phe Arg Val Ile  
                             50                            55                            60  
 Pro Ala Met Met Ala Gly Gly Ala Thr Thr Gly Ala Ile Ser Met Ala  
                             65                            70                            75                            80  
 Leu Gly Val Gly Ser Arg Ala Pro His Gly Gly Ile Phe Val Val Trp  
                             85                            90                            95  
 Ala Ile Glu Pro Trp Trp Gly Trp Leu Ile Ala Leu Ala Ala Gly Thr  
                             100                            105                            110  
 Ile Val Ser Thr Ile Val Val Ile Ala Leu Lys Gln Phe Trp Pro Asn  
                             115                            120                            125  
 Lys Ala Val Ala Ala Glu Val Ala Lys Gln Glu Ala Gln Gln Ala Ala  
                             130                            135                            140  
 Val Asn Ala  
 145

<210> 11  
 <211> 580  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (101)..(580)  
 <223> FRXA01883

<400> 11

```

cgactgcggc gtctcttctt ggcactacca ttctctgctcc tgaccaactc gccacagctg 60
gtgcacgggt cccccagtc aaaggattga aagaatcagc atg aat agc gta aat 115
                                         Met Asn Ser Val Asn
                                         1      5

aat tcc tgg ctt gtc cgg ctg gat gtc gat ttc ggc gac tcc acc acg 163
Asn Ser Ser Leu Val Arg Leu Asp Val Asp Phe Gly Asp Ser Thr Thr
                        10                        15                        20

gat gtc atc aac aac ctt gcc act gtt att ttc gac gct ggc cga gct 211
Asp Val Ile Asn Asn Leu Ala Thr Val Ile Phe Asp Ala Gly Arg Ala
                        25                        30                        35

tcc tcc gcc gac gcc ctt gcc aaa gac ggc ctg gat cgt gaa gca aag 259
Ser Ser Ala Asp Ala Leu Ala Lys Asp Ala Leu Asp Arg Glu Ala Lys
                        40                        45                        50

tcc ggc acc ggc gtt cct ggt caa gtt gct atc ccc ccc tgc cgt tcc 307
Ser Gly Thr Gly Val Pro Gly Gln Val Ala Ile Pro His Cys Arg Ser
                        55                        60                        65

gaa gcc gta tct gtc cct ccc ttg gcc ttt gct cgc ctg agc aag ggt 355
Glu Ala Val Ser Val Pro Thr Leu Gly Phe Ala Arg Leu Ser Lys Gly
                        70                        75                        80                        85

gtg gac ttc agc gga cct gat ggc gat gcc aac ttc gtc ttc ctc ctt 403
Val Asp Phe Ser Gly Pro Asp Gly Asp Ala Asn Leu Val Phe Leu Ile
                        90                        95                        100

gca gca cct gct ggc ggc ggc aaa gag cac ctg aag atc ctg tcc aag 451
Ala Ala Pro Ala Gly Gly Gly Lys Glu His Leu Lys Ile Leu Ser Lys
                        105                        110                        115

ctt gct cgc tcc ttg gtg aag aag gat ttc atc aag gct ctg cag gaa 499
Leu Ala Arg Ser Leu Val Lys Lys Asp Phe Ile Lys Ala Leu Gln Glu
                        120                        125                        130

gcc acc acc gag cag gaa atc gtc gac gtt gtc gat gcc gtg ctc aac 547
Ala Thr Thr Glu Gln Glu Ile Val Asp Val Val Asp Ala Val Leu Asn
                        135                        140                        145

cca gca cca aaa aac cac cga gcc agc tgc agc 580
Pro Ala Pro Lys Asn His Arg Ala Ser Cys Ser
150                        155                        160

```

<210> 12  
 <211> 160  
 <212> PRT  
 <213> Corynebact rium glutamicum

<400> 12

Met Asn Ser Val Asn Asn Ser Ser Leu Val Arg Leu Asp Val Asp Phe

1	5	10	15
Gly Asp Ser Thr Thr Asp Val Ile Asn Asn Leu Ala Thr Val Il Phe	20	25	30
Asp Ala Gly Arg Ala Ser Ser Ala Asp Ala Leu Ala Lys Asp Ala Leu	35	40	45
Asp Arg Glu Ala Lys Ser Gly Thr Gly Val Pro Gly Gln Val Ala Ile	50	55	60
Pro His Cys Arg Ser Glu Ala Val Ser Val Pro Thr Leu Gly Phe Ala	65	70	75
Arg Leu Ser Lys Gly Val Asp Phe Ser Gly Pro Asp Gly Asp Ala Asn	85	90	95
Leu Val Phe Leu Ile Ala Ala Pro Ala Gly Gly Gly Lys Gln His Leu	100	105	110
Lys Ile Leu Ser Lys Leu Ala Arg Ser Leu Val Lys Lys Asp Phe Ile	115	120	125
Lys Ala Leu Gln Glu Ala Thr Thr Glu Gln Glu Ile Val Asp Val Val	130	135	140
Asp Ala Val Leu Asn Pro Ala Pro Lys Asn His Arg Ala Ser Cys Ser	145	150	155
			160

<210> 13  
 <211> 631  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (77)..(631)  
 <223> FKXA01889

<400> 13  
 accgagccag ctgcagctcc ggetgcggcg gcgggttggt aagagtgggg cggcgctcgac 60  
 aagcggttaact cgtatcggtg gca atc acc gcc tgc cca acc ggt atc gcc cac 112  
 Val Ala Ile Thr Ala Cys Pro Thr Gly Ile Ala His  
 1 5 10  
 acc tac atg gct gcg gat tcc ctg acg caa aac gcg gaa ggc cgc gat 160  
 Thr Tyr Met Ala Ala Asp Ser Leu Thr Gln Asn Ala Gln Gly Arg Asp  
 15 20 25  
 gat gtg gaa ctg gtt gtg gag act cag ggc tct tcc gct gtc acc cca 208  
 Asp Val Glu Leu Val Val Glu Thr Gln Gly Ser Ser Ala Val Thr Pro  
 30 35 40  
 gtc gat ccg aag atc atc gaa gct gcc gac gcc gtc atc ttc gcc acc 256  
 Val Asp Pro Lys Ile Ile Glu Ala Ala Asp Ala Val Ile Phe Ala Thr  
 45 50 55 60

gag gtc gga gtt aca gac cgc gag cgt ttc gct ggc aag cca gtc att 304  
 Asp Val Gly Val Lys Asp Arg Glu Arg Phe Ala Gly Lys Pro Val Ile  
 65 70 75  
 gaa tcc ggc gtc aag cgc gcg atc aat gag cca gcc aag atg atc gac 352  
 Glu Ser Cys Val Lys Arg Ala Ile Asn Glu Pro Ala Lys Met Ile Asp  
 80 85 90  
 gag gcc atc gca gcc tcc aag aac cca aac gcc cgc aag gtt tcc ggt 400  
 Glu Ala Ile Ala Ala Ser Lys Asn Pro Asn Ala Arg Lys Val Ser Gly  
 95 100 105  
 tcc ggt gtc gcg gca tct gct gaa acc acc ggc gag aag ctc ggc tgg 448  
 Ser Gly Val Ala Ala Ser Ala Glu Thr Thr Gly Glu Lys Leu Gly Trp  
 110 115 120  
 ggc aag cgc atc cag cag gca gtc atg acc ggc gtg tcc tac atg gtt 496  
 Gly Lys Arg Ile Gln Gln Ala Val Met Thr Gly Val Ser Tyr Met Val  
 125 130 135 140  
 cca ttc gta gct gcc ggc ggc ctc ctg ttg gct ctc ggc ttc gca ttc 544  
 Pro Phe Val Ala Ala Gly Gly Leu Leu Ala Leu Gly Phe Ala Phe  
 145 150 155  
 ggt gga tac gac atg gcg aac ggc tgg caa gca atc gcc acc cag ttc 592  
 Gly Gly Tyr Asp Met Ala Asn Cys Trp Cln Ala Ile Ala Thr Gln Phe  
 160 165 170  
 tct ctg acc aac ctg cca ggc aac acc gtc gat att gac 631  
 Ser Leu Thr Asn Leu Pro Gly Asn Thr Val Asp Val Asp  
 175 180 185

&lt;210&gt; 14

&lt;211&gt; 185

&lt;212&gt; PRT

&lt;213&gt; Corynebacterium glutamicum

&lt;400&gt; 14

Val Ala Ile Thr Ala Cys Pro Thr Gly Ile Ala His Thr Tyr Met Ala  
 1 5 10 15  
 Ala Asp Ser Leu Thr Gln Asn Ala Glu Gly Arg Asp Asp Val Glu Leu  
 20 25 30  
 Val Val Glu Thr Gln Gly Ser Ser Ala Val Thr Pro Val Asp Pro Lys  
 35 40 45  
 Ile Ile Glu Ala Ala Asp Ala Val Ile Phe Ala Thr Asp Val Gly Val  
 50 55 60  
 Lys Asp Arg Glu Arg Phe Ala Gly Lys Pro Val Ile Glu Ser Gly Val  
 65 70 75 80  
 Lys Arg Ala Ile Asn Glu Pro Ala Lys Met Ile Asp Glu Ala Ile Ala  
 85 90 95  
 Ala S r Lys Asn Pro Asn Ala Arg Lys Val Ser Gly Ser Gly Val Ala  
 100 105 110

Ala Ser Ala Glu Thr Thr Gly Glu Lys Leu Gly Trp Gly Lys Arg Ile  
 115 120 125

Gln Gln Ala Val Met Thr Gly Val Ser Tyr Met Val Pro Phe Val Ala  
 130 135 140

Ala Gly Gly Leu Leu Leu Ala Leu Gly Phe Ala Phe Gly Gly Tyr Asp  
 145 150 155 160

Met Ala Asn Gly Trp Gln Ala Ile Ala Thr Gln Phe Ser Leu Thr Asn  
 165 170 175

Leu Pro Gly Asn Thr Val Asp Val Asp  
 180 185

<210> 15  
 <211> 416  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> {1}..(393)  
 <223> RXA00951

<400> 15  
 atc caa gca atc tta gag aag gca gca gcg ccg gcg aag cag aag got 48  
 Ile Gln Ala Ile Leu Glu Lys Ala Ala Ala Pro Ala Lys Gln Lys Ala  
 1 5 10 15

cct gct gtg gct cct gct gta aca ccc act gac gct cct gca gcc tca 96  
 Pro Ala Val Ala Pro Ala Val Thr Pro Thr Asp Ala Pro Ala Ala Ser  
 20 25 30

gtc caa tcc aac acc cac gac aag atc ctc acc gtc tgt gcc aac gcc 144  
 Val Gln Ser Lys Thr His Asp Lys Ile Leu Thr Val Cys Gly Asn Gly  
 35 40 45

ttg ggt acc tcc ctc ttc ctc aac aac acc ctt gag caa gtt ttc gac 192  
 Leu Gly Thr Ser Leu Phe Leu Lys Asn Thr Leu Glu Gln Val Phe Asp  
 50 55 60

acc tgg ggt tgg ggt cca tcc atg acc gtc gag gac acc gac act atc 240  
 Thr Trp Gly Trp Gly Pro Tyr Met Thr Val Glu Ala Thr Asp Thr Ile  
 65 70 75 80

tcc gcc aag gcc aac gcc aag gaa gct gat ctc atc atg acc tct ggt 288  
 Ser Ala Lys Gly Lys Ala Lys Glu Ala Asp Leu Ile Met Thr Ser Gly  
 85 90 95

gaa atc gcc cgc aag ttg ggt gat gtt gga atc ccg gtt cac gtg atc 336  
 Glu Ile Ala Arg Thr Leu Gly Asp Val Gly Ile Pro Val His Val Ile  
 100 105 110

aat gac ttc aag agc acc gat gaa atc gat gct gcg ctt cgt gaa cgc 384  
 Asn Asp Phe Thr Ser Thr Asp Glu Ile Asp Ala Ala Leu Arg Glu Arg  
 115 120 125

tac gac atc taactacttt aasaggacga aaa 416  
 Tyr Asp Ile

130

<210> 16  
 <211> 131  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 16  
 Ile Glu Ala Ile Leu Glu Lys Ala Ala Ala Pro Ala Lys Gln Lys Ala  
 1 5 10 15  
 Pro Ala Val Ala Pro Ala Val Thr Pro Thr Asp Ala Pro Ala Ala Ser  
 20 25 30  
 Val Gln Ser Lys Thr His Asp Lys Ile Leu Thr Val Cys Gly Asn Gly  
 35 40 45  
 Leu Gly Thr Ser Leu Phe Leu Lys Asn Thr Leu Glu Gln Val Phe Asp  
 50 55 60  
 Thr Trp Gly Trp Gly Pro Tyr Met Thr Val Glu Ala Thr Asp Thr Ile  
 65 70 75 80  
 Ser Ala Lys Gly Lys Ala Lys Glu Ala Asp Leu Ile Met Thr Ser Gly  
 85 90 95  
 Glu Ile Ala Arg Thr Leu Gly Asp Val Gly Ile Pro Val His Val Ile  
 100 105 110  
 Asn Asp Phe Thr Ser Thr Asp Glu Ile Asp Ala Ala Leu Arg Glu Arg  
 115 120 125  
 Tyr Asp Ile  
 130

<210> 17  
 <211> 1827  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> [101]..(1804)  
 <223> RXN01244

<400> 17  
 gatattgtgtt tgtttgtcaa tatcccaatg tttagaatgt tgcacaactg ttggttttgt 60  
 ggtgntcttg aggaatttaa ctcaatgatt gtgaggatgg gtg gct act gtg gct 115  
 Val Ala Thr Val Ala  
 1 5  
 gat gtg aat caa gac act gta ctg aag ggc acc ggc gtt gtc ggc gga 163  
 Asp Val Asn Gln Asp Thr Val Leu Lys Gly Thr Gly Val Val Gly Gly  
 10 15 20  
 gtc cgt tat gca agc ggc ggc tgg att acc cca cgc ccc gaa cta ccc 211  
 Val Arg Tyr Ala Ser Ala Val Trp Ile Thr Pro Arg Pro Glu Leu Pro  
 25 30 35



caa gca ggc gaa gtc gtc gcc gaa gaa aac cgt gaa gca gag cag gag	259
Gln Ala Gly Glu Val Val Ala Glu Glu Asn Arg Glu Ala Glu Gln Glu	
40 45 50	
cgt ttc gac gcc gct gcc gca gca gtc tct tct cgt ttg ctt gag cgc	307
Arg Phe Asp Ala Ala Ala Ala Thr Val Ser Ser Arg Leu Leu Glu Arg	
55 60 65	
tcc gaa gct gct gaa gga cca gcc gcc gag gtg ctt aaa gct act gct	355
Ser Glu Ala Ala Glu Gly Pro Ala Ala Glu Val Leu Lys Ala Thr Ala	
70 75 80 85	
ggc atg gtc aat gac cgt gcc tgg cgt aag gct gtc atc aag ggt gtc	403
Gly Met Val Asn Asp Arg Gly Trp Arg Lys Ala Val Ile Lys Gly Val	
90 95 100	
aag ggt ggt ccc cct gcc gaa tac gcc gtg gtt gca gca aca acc aag	451
Lys Gly Gly His Pro Ala Glu Tyr Ala Val Val Ala Ala Thr Thr Lys	
105 110 115	
ttc atc tcc atg ttc gaa gcc gca gcc gcc ctg atc gcc gag cgc acc	499
Phe Ile Ser Met Phe Glu Ala Ala Gly Gly Leu Ile Ala Glu Arg Thr	
120 125 130	
aca gac ttg cgc gac aac cgc gcc cgc gtc etc gca gaa ctt cgt gcc	547
Thr Asp Leu Arg Asp Ile Arg Asp Arg Val Ile Ala Glu Leu Arg Gly	
135 140 145	
gat gaa gag cca ggt ctg cca gcc gcc gcc cag gtc att etc ttt	595
Asp Glu Glu Pro Gly Leu Pro Ala Val Ser Gly Gln Val Ile Leu Phe	
150 155 160 165	
gca gat gac ctc tcc cca gca gac acc gcc gca cta gac aca gct ctc	643
Ala Asp Asp Leu Ser Pro Ala Asp Thr Ala Ala Leu Asp Thr Asp Leu	
170 175 180	
ttt gtg gga ctt gtc act gag ctg ggt gcc cca acg agc cac acc gcc	691
Phe Val Gly Leu Val Thr Glu Leu Gly Gly Pro Thr Ser His Thr Ala	
185 190 195	
atc atc gca cgc cag ctc aac gtg cct tgc atc gtc gca tcc gcc gcc	739
Ile Ile Ala Arg Glu Leu Asn Val Pro Cys Ile Val Ala Ser Gly Ala	
200 205 210	
ggc atc aag gnc atc aag tcc gcc gaa aag gtg ctt atc gac gcc agc	787
Gly Ile Lys Asp Ile Lys Ser Gly Glu Lys Val Leu Ile Asp Gly Ser	
215 220 225	
ctc gcc acc att gac cgc aac gcc gac gaa gct gaa gca acc aag ctc	835
Leu Gly Thr Ile Asp Arg Asn Ala Asp Glu Ala Glu Ala Thr Lys Leu	
230 235 240 245	
gtc tcc gag tcc ctc gag cgc gct gcc gcc atc gcc gag tgg aag ggt	883
Val Ser Glu Ser Leu Glu Arg Ala Ala Arg Ile Ala Glu Trp Lys Gly	
250 255 260	
cct gca caa acc aag gac gcc tac gcc gtt cag ctg ttg gcc aac gtc	931
Pro Ala Gln Thr Lys Asp Gly Tyr Arg Val Gln Leu Leu Ala Asn Val	
265 270 275	

caa gac ggc aac tct gcc cag cag gct gca cag acc gaa gca gaa ggc	979
Gln Asp Gly Asn Ser Ala Gln Gln Ala Ala Gln Thr Glu Ala Glu Gly	
280 285 290	
atc ggc ctg ttc cgc acc gaa ctg tgc ttc ctt tcc gcc acc gaa gag	1027
Ile Gly Leu Phe Arg Thr Glu Leu Cys Phe Leu Ser Ala Thr Glu Glu	
295 300 305	
cca agc gtt gat gag cag gct gcg gtc tac tca aag gtg ctt gaa gca	1075
Pro Ser Val Asp Glu Gln Ala Ala Val Tyr Ser Lys Val Leu Glu Ala	
310 315 320 325	
ttc cca gag tcc aag gtc gtt gtc cgc tcc ctc gac gca ggt tct gac	1123
Phe Pro Glu Ser Lys Val Val Val Arg Ser Leu Asp Ala Gly Ser Asp	
330 335 340	
aag cca gtt cca ttc gca tgc atg gct gat gag atg aac cca gca ctg	1171
Lys Pro Val Phe Ala Ser Met Ala Asp Glu Met Asn Pro Ala Leu	
345 350 355	
ggt gtt cgt ggc ctg cgt atc gca cgt gga cag gtc gat ctg ctg act	1219
Gly Val Arg Gly Leu Arg Ile Ala Arg Gly Gln Val Asp Leu Leu Thr	
360 365 370	
cgc cag ctc gac gca att gcg aag gcc agc gaa gaa ctc ggc cgt ggc	1267
Arg Gln Leu Asp Ala Ile Ala Lys Ala Ser Glu Glu Leu Gly Arg Gly	
375 380 385	
gac gac gcc cca acc tgg gtt atg gct cca atg gtg gct acc gct tat	1315
Asp Asp Ala Pro Thr Trp Val Met Ala Pro Met Val Ala Thr Ala Tyr	
390 395 400 405	
gaa gca aag tgg ttt gct gac atg tgc cgt gag cgt ggc cta atc gcc	1363
Glu Ala Lys Trp Phe Ala Asp Met Cys Arg Glu Arg Gly Leu Ile Ala	
410 415 420	
ggc gcc atg atc gaa gtt cca gca gca tcc ctg atg gca gac aag atc	1411
Gly Ala Met Ile Glu Val Pro Ala Ala Ser Leu Met Ala Asp Lys Ile	
425 430 435	
atg cct cac ctg gac ttt gtt tcc atc ggt acc aac gac ctg acc cag	1459
Met Pro His Leu Asp Phe Val Ser Ile Gly Thr Asn Asp Leu Thr Cln	
440 445 450	
tac acc atg gca gcg gac cgc atg tct cct gag ctt gcc tac ctg acc	1507
Tyr Thr Met Ala Ala Asp Arg Met Ser Pro Glu Leu Ala Tyr Leu Thr	
455 460 465	
gat cct tgg cag cca gca gtc ctg cgc ctg atc aag cac acc tgt gac	1555
Asp Pro Trp Gln Pro Ala Val Leu Arg Leu Ile Lys His Thr Cys Asp	
470 475 480 485	
gaa ggt gct cgc ttt aac aac cag gtc ggt gtt tgt ggt gaa gca gca	1603
Glu Gly Ala Arg Phe Asn Thr Pro Val Gly Val Cys Gly Glu Ala Ala	
490 495 500	
gca gac cca ctg ttg gca act gtc ctc acc ggt ctt gcc gtg aac tcc	1651
Ala Asp Pro Leu Leu Ala Thr Val Leu Thr Gly Leu Gly Val Asn Ser	
505 510 515	
ctg tcc gca gca tcc act gct ctc gca gca gtc ggt gcc aag cln tcc	1699

Leu Ser Ala Ala Ser Thr Ala Leu Ala Ala Val Gly Ala Lys Leu Ser  
 520 525 530  
 gag gtc acc ctg gaa acc tgl aag aag gca gca gaa gca gca ctt gac 1747  
 Glu Val Thr Leu Glu Thr Cys Lys Lys Ala Ala Glu Ala Ala Leu Asp  
 535 540 545  
 gct gaa ggt gca act gaa gca agc gat gct gta cgc gca glg atc gac 1795  
 Ala Glu Gly Ala Thr Glu Ala Arg Asp Ala Val Arg Ala Val Ile Asp  
 550 555 560 565  
 gca gca glc taaccacag ttgagctaa aag 1827  
 Ala Ala Val

&lt;210&gt; 18

&lt;211&gt; 568

&lt;212&gt; PRT

<213> *Corynebacterium glutamicum*

&lt;400&gt; 18

Val Ala Thr Val Ala Asp Val Asn Gln Asp Thr Val Leu Lys Gly Thr  
 1 5 10 15  
 Gly Val Val Gly Gly Val Arg Tyr Ala Ser Ala Val Trp Ile Thr Pro  
 20 25 30  
 Arg Pro Glu Leu Pro Gln Ala Gly Glu Val Val Ala Glu Glu Asn Arg  
 35 40 45  
 Glu Ala Glu Gln Glu Arg Phe Asp Ala Ala Ala Thr Val Ser Ser  
 50 55 60  
 Arg Leu Leu Glu Arg Ser Glu Ala Ala Glu Gly Pro Ala Ala Glu Val  
 65 70 75 80  
 Leu Lys Ala Thr Ala Gly Met Val Asn Asp Arg Gly Trp Arg Lys Ala  
 85 90 95  
 Val Ile Lys Gly Val Lys Gly Gly His Pro Ala Glu Tyr Ala Val Val  
 100 105 110  
 Ala Ala Thr Thr Lys Phe Ile Ser Met Phe Glu Ala Ala Gly Gly Leu  
 115 120 125  
 Ile Ala Glu Arg Thr Thr Asp Leu Arg Asp Ile Arg Asp Arg Val Ile  
 130 135 140  
 Ala Glu Leu Arg Gly Asp Glu Glu Pro Gly Leu Pro Ala Val Ser Gly  
 145 150 155 160  
 Gln Val Ile Leu Phe Ala Asp Asp Leu Ser Pro Ala Asp Thr Ala Ala  
 165 170 175  
 Leu Asp Thr Asp Leu Phe Val Gly Leu Val Thr Glu Leu Gly Gly Pro  
 180 185 190  
 Thr Ser His Thr Ala Ile Ile Ala Arg Gln L u Asn Val Pro Cys Ile  
 195 200 205

Val Ala Ser Gly Ala Gly Ile Lys Asp Ile Lys Ser Gly Glu Lys Val  
 210 215 220  
 Leu Ile Asp Gly Ser Leu Gly Thr Ile Asp Arg Asn Ala Asp Glu Ala  
 225 230 235 240  
 Glu Ala Thr Lys Leu Val Ser Glu Ser Leu Glu Arg Ala Ala Arg Ile  
 245 250 255  
 Ala Glu Trp Lys Gly Pro Ala Gln Thr Lys Asp Gly Tyr Arg Val Gln  
 260 265 270  
 Leu Leu Ala Asn Val Gln Asp Gly Asn Ser Ala Gln Gln Ala Ala Gln  
 275 280 285  
 Thr Glu Ala Glu Gly Ile Gly Leu Phe Arg Thr Glu Leu Cys Phe Leu  
 290 295 300  
 Ser Ala Thr Glu Glu Pro Ser Val Asp Glu Gln Ala Ala Val Tyr Ser  
 305 310 315 320  
 Lys Val Leu Glu Ala Phe Pro Glu Ser Lys Val Val Val Arg Ser Leu  
 325 330 335  
 Asp Ala Gly Ser Asp Lys Pro Val Pro Phe Ala Ser Met Ala Asp Glu  
 340 345 350  
 Met Asn Pro Ala Leu Gly Val Arg Gly Leu Arg Ile Ala Arg Gly Gln  
 355 360 365  
 Val Asp Leu Leu Thr Arg Gln Leu Asp Ala Ile Ala Lys Ala Ser Glu  
 370 375 380  
 Glu Leu Gly Arg Gly Asp Asp Ala Pro Thr Trp Val Met Ala Pro Met  
 385 390 395 400  
 Val Ala Thr Ala Tyr Glu Ala Lys Trp Phe Ala Asp Met Cys Arg Glu  
 405 410 415  
 Arg Gly Leu Ile Ala Gly Ala Met Ile Glu Val Pro Ala Ala Ser Leu  
 420 425 430  
 Met Ala Asp Lys Ile Met Pro His Leu Asp Phe Val Ser Ile Gly Thr  
 435 440 445  
 Asn Asp Leu Thr Gln Tyr Thr Met Ala Ala Asp Arg Met Ser Pro Glu  
 450 455 460  
 Leu Ala Tyr Leu Thr Asp Pro Trp Gln Pro Ala Val Leu Arg Leu Ile  
 465 470 475 480  
 Lys His Thr Cys Asp Glu Gly Ala Arg Phe Asn Thr Pro Val Gly Val  
 485 490 495  
 Cys Gly Glu Ala Ala Ala Asp Pro Leu Leu Ala Thr Val Leu Thr Gly  
 500 505 510  
 Leu Gly Val Asn Ser Leu Ser Ala Ala Ser Thr Ala Leu Ala Ala Val  
 515 520 525  
 Gly Ala Lys Leu Ser Glu Val Thr Leu Glu Thr Cys Lys Lys Ala Ala

530

535

540

Glu Ala Ala Leu Asp Ala Glu Gly Ala Thr Glu Ala Arg Asp Ala Val  
 545 550 555 560

Arg Ala Val Ile Asp Ala Ala Val  
 565

&lt;210&gt; 19

&lt;211&gt; 1629

&lt;212&gt; DNA

&lt;213&gt; Corynebacterium glutamicum

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; {98}..(1606)

&lt;223&gt; FRXA01244

&lt;400&gt; 19

agatgtcgat ttctcgagga agaagttaac gccgaagaaa accgtgaatc agagcaggag 60

cgttctgagc cgcctgagc cctctctctc tcttcttctc ctc cgc cgc tcc gaa 115  
 Leu Leu Glu Arg Ser Glu  
 1 5

gct gct gaa gga cca gca gct gaa gtc ctc asa gct act gct ggc atg 163  
 Ala Ala Glu Gly Pro Ala Ala Glu Val Leu Lys Ala Thr Ala Gly Met  
 10 15 20

gtc aat gac cgt ggc tgg cgt aag gct gtc atc aag ggt gtc aag ggt 211  
 Val Asn Asp Arg Gly Trp Arg Lys Ala Val Ile Lys Gly Val Lys Gly  
 25 30 35

ggt cac cct ggc gaa tac gcc gtg gtt gca gca aca acc aag ttc atc 259  
 Gly His Pro Ala Glu Tyr Ala Val Val Ala Ala Thr Thr Lys Phe Ile  
 40 45 50

tcc atg ttc gaa gcc gca ggc ggc ctg atc ggc gag cgc acc aca gac 307  
 Ser Met Phe Glu Ala Ala Gly Gly Leu Ile Ala Glu Arg Thr Thr Asp  
 55 60 65 70

ttg cgc gac atc cgc gac cgc gtc atc gca gaa ctt cgt ggc gat gaa 355  
 Leu Arg Asp Thr Arg Asp Arg Val Ile Ala Glu Leu Arg Gly Asp Glu  
 75 80 85

gag cca ggt ctg cca got gtt tcc gga cag gtc att ctc ttt gca gat 403  
 Glu Pro Gly Leu Pro Ala Val Ser Gly Gln Val Ile Leu Phe Ala Asp  
 90 95 100

gac ctc tcc cca gca gac acc ggc gca cta gac aca gat ctc ttt gtg 451  
 Asp Leu Ser Pro Ala Asp Thr Ala Ala Leu Asp Thr Asp Leu Phe Val  
 105 110 115

gga ctt gtc act gag ctg ggt ggc cca acg agc ccc acc ggc atc atc 499  
 Gly Leu Val Thr Glu Leu Gly Gly Pro Thr Ser His Thr Ala Ile Ile  
 120 125 130

gca cgc cag ttc aac gtg cct tgc atc gtc gca tcc ggc gcc ggc atc 547  
 Ala Arg Gln Leu Asn Val Pro Cys Ile Val Ala Ser Gly Ala Gly Ile  
 135 140 145 150

aag gac atc aag tcc ggc gaa aag gtg ctt atc gac ggc agc ctc ggc	595
Lys Asp Ile Lys Ser Gly Glu Lys Val Leu Ile Asp Gly Ser Leu Gly	
155 160 165	
acc att gac cgc aac gcg gac gaa gct gaa gca acc aag ctc gtc tcc	643
Thr Ile Asp Arg Asn Ala Asp Glu Ala Glu Ala Thr Lys Leu Val Ser	
170 175 180	
gag tcc ctc gag cgc gct gct cgc atc gcc gag tgg aag ggt cct gca	691
Glu Ser Leu Glu Arg Ala Ala Arg Ile Ala Glu Trp Lys Gly Pro Ala	
185 190 195	
caa acc aag gac ggc tac cgc gtt cag ctg ttg gcc aac gtc caa gac	739
Gln Thr Lys Asp Gly Tyr Arg Val Gln Leu Leu Ala Asn Val Gln Asp	
200 205 210	
ggc aac tct gca cag cag gct gca cag acc gaa gca gaa ggc atc ggc	787
Gly Asn Ser Ala Gln Gln Ala Ala Gln Thr Glu Ala Glu Gly Ile Gly	
215 220 225 230	
ctg ttc cgc acc gaa ctg tgc ttc ctt tcc gcc acc gaa gag cca agr	835
Leu Phe Arg Thr Glu Leu Cys Phe Leu Ser Ala Thr Glu Glu Pro Ser	
235 240 245	
gtt gat gag cag gct gcg gtc tac tca aag gtg ctt gaa gca ttc cca	883
Val Asp Glu Gln Ala Ala Val Tyr Ser Lys Val Leu Glu Ala Phe Pro	
250 255 260	
gag tcc aag gtc gtt gtc cgc tcc ctc gac gca ggt tct gac aag cca	931
Glu Ser Lys Val Val Val Arg Ser Leu Asp Ala Gly Ser Asp Lys Pro	
265 270 275	
gtt cca ttc gca tcc atg gct gat gag atg aac cca gca ctg ggt gtt	979
Val Pro Phe Ala Ser Met Ala Asp Glu Met Asn Pro Ala Leu Gly Val	
280 285 290	
cgt ggc ctg cgt atc gca cgt gga cag gtt gat ctg ctg act cgc cag	1027
Arg Gly Leu Arg Ile Ala Arg Gly Gln Val Asp Leu Leu Thr Arg Gln	
295 300 305 310	
ctc gac gca att gcg aag gcc agc gaa gaa ctc ggc cgt ggc gac gac	1075
Leu Asp Ala Ile Ala Lys Ala Ser Glu Glu Leu Gly Arg Gly Asp Asp	
315 320 325	
gcc cca acc tgg gtt atg gct cca atg gtg gct acc gct tat gaa gcc	1123
Ala Pro Thr Trp Val Met Ala Pro Met Val Ala Thr Ala Tyr Glu Ala	
330 335 340	
aag tgg ttt gct gac atg tgc cgt gag cgt ggc ctc atc gcc agc gcc	1171
Lys Trp Phe Ala Asp Met Cys Arg Glu Arg Gly Leu Ile Ala Gly Ala	
345 350 355	
atg atc gaa gtt cca gca gca tcc ctg atg gca gac aag atc atg cct	1219
Met Ile Glu Val Pro Ala Ala Ser Leu Met Ala Asp Lys Ile Met Pro	
360 365 370	
cac ctg gac ttt gtt tcc atc ggt acc aac gac ctg acc cag tac acc	1267
His Leu Asp Phe Val Ser Ile Gly Thr Asn Asp Leu Thr Gln Tyr Thr	
375 380 385 390	

atg gca gcg gac cgc atg tct cct gag ctt gcc tac ctg acc gat cct 1315  
 Met Ala Ala Asp Arg Met Ser Pro Glu Leu Ala Tyr Leu Thr Asp Pro  
 395 400 405  
 tgg cag cca gca gtc ctg cgc ctg atc aag cac acc tgt gac gaa ggt 1363  
 Trp Cln Pro Ala Val Leu Arg Leu Ile Lys His Thr Cys Asp Glu Gly  
 410 415 420  
 gct cgc ttt aac acc ccg gtc ggt gtt tgt ggt gaa gca gca gca gac 1411  
 Ala Arg Phe Asn Thr Pro Val Gly Val Cys Gly Cln Ala Ala Ala Asp  
 425 430 435  
 cca ctg ttg gca act gtc ctc acc ggt ctt ggc gtg aac tcc ctg tcc 1459  
 Pro Leu Leu Ala Thr Val Leu Thr Gly Leu Gly Val Asn Ser Leu Ser  
 440 445 450  
 gca gca tcc act gct ctc gca gca gtc ggt gca aag ctg tca gag gtc 1507  
 Ala Ala Ser Thr Ala Leu Ala Ala Val Gly Ala Lys Leu Ser Cln Val  
 455 460 465 470  
 acc ctg gaa acc tgt aag aag gca gca gaa gca gca ctt gac gct gaa 1555  
 Thr Leu Glu Thr Cys Lys Lys Ala Ala Glu Ala Ala Leu Asp Ala Cln  
 475 480 485  
 ggt gca act gaa gca cgc gat gct gta cgc gca gtg atc gac gca gca 1603  
 Gly Ala Thr Glu Ala Arg Asp Ala Val Arg Ala Val Ile Asp Ala Ala  
 490 495 500  
 gtc taaaccactg ttgagctaaa aag 1629  
 Val

<210> 20  
 <211> 503  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 20  
 Leu Leu Glu Arg Ser Cln Ala Ala Cln Gly Pro Ala Ala Glu Val Leu  
 1 5 10 15  
 Lys Ala Thr Ala Gly Met Val Asn Asp Arg Gly Trp Arg Lys Ala Val  
 20 25 30  
 Ile Lys Gly Val Lys Gly Gly His Pro Ala Glu Tyr Ala Val Val Ala  
 35 40 45  
 Ala Thr Thr Lys Phe Ile Ser Met Phe Glu Ala Ala Gly Gly Leu Ile  
 50 55 60  
 Ala Glu Arg Thr Thr Asp Leu Arg Asp Ile Arg Asp Arg Val Ile Ala  
 65 70 75 80  
 Glu Leu Arg Gly Asp Glu Glu Pro Gly Leu Pro Ala Val Ser Gly Gln  
 85 90 95  
 Val Ile L u Phe Ala Asp Asp Leu Ser Pro Ala Asp Thr Ala Ala Leu  
 100 105 110  
 Asp Thr Asp Leu Phe Val Gly Leu Val Thr Glu Leu Gly Gly Pro Thr

115					120					125					
Ser	His	Thr	Ala	Ile	Il	Ala	Arg	Gln	Leu	Asn	Val	Pro	Cys	Ile	Val
130					135					140					
Ala	Ser	Gly	Ala	Gly	Ile	Lys	Asp	Ile	Lys	Ser	Gly	Glu	Lys	Val	Leu
145					150					155					160
Ile	Asp	Gly	Ser	Leu	Gly	Thr	Ile	Asp	Arg	Asn	Ala	Asp	Glu	Ala	Glu
165					170					175					
Ala	Thr	Lys	Leu	Val	Ser	Glu	Ser	Leu	Glu	Arg	Ala	Ala	Arg	Ile	Ala
180					185					190					
Glu	Trp	Lys	Gly	Pro	Ala	Gln	Thr	Lys	Asp	Gly	Tyr	Arg	Val	Gln	Leu
195					200					205					
Leu	Ala	Asn	Val	Glu	Asp	Gly	Asn	Ser	Ala	Gln	Gln	Ala	Ala	Gln	Thr
210					215					220					
Glu	Ala	Glu	Gly	Ile	Gly	Leu	Phe	Arg	Thr	Glu	Leu	Cys	Phe	Leu	Ser
225					230					235					240
Ala	Thr	Glu	Glu	Pro	Ser	Val	Asp	Glu	Gln	Ala	Ala	Val	Tyr	Ser	Lys
245					250					255					
Val	Leu	Glu	Ala	Phe	Pro	Glu	Ser	Lys	Val	Val	Val	Arg	Ser	Leu	Asp
260					265					270					
Ala	Gly	Ser	Asp	Lys	Pro	Val	Pro	Phe	Ala	Ser	Met	Ala	Asp	Glu	Met
275					280					285					
Asn	Pro	Ala	Leu	Gly	Val	Arg	Gly	Leu	Arg	Ile	Ala	Arg	Gly	Gln	Val
290					295					300					
Asp	Leu	Leu	Thr	Arg	Gln	Leu	Asp	Ala	Ile	Ala	Lys	Ala	Ser	Glu	Glu
305					310					315					320
Leu	Gly	Arg	Gly	Asp	Asp	Ala	Pro	Thr	Trp	Val	Met	Ala	Pro	Met	Val
325					330					335					
Ala	Thr	Ala	Tyr	Glu	Ala	Lys	Trp	Phe	Ala	Asp	Met	Cys	Arg	Glu	Arg
340					345					350					
Gly	Leu	Ile	Ala	Gly	Ala	Met	Ile	Glu	Val	Pro	Ala	Ala	Ser	Leu	Met
355					360					365					
Ala	Asp	Lys	Ile	Met	Pro	His	Leu	Asp	Phe	Val	Ser	Ile	Gly	Thr	Asn
370					375					380					
Asp	Leu	Thr	Gln	Tyr	Thr	Met	Ala	Ala	Asp	Arg	Met	Ser	Pro	Glu	Leu
385					390					395					400
Ala	Tyr	Leu	Thr	Asp	Pro	Trp	Gln	Pro	Ala	Val	Leu	Arg	Leu	Ile	Lys
405					410					415					
His	Thr	Cys	Asp	Glu	Gly	Ala	Arg	Phe	Asn	Thr	Pro	Val	Gly	Val	Cys
420					425					430					
Gly	Glu	Ala	Ala	Ala	Asp	Pro	Leu	Leu	Ala	Thr	Val	Leu	Thr	Gly	Leu
435					440					445					



Gly Val Asn Ser Leu Ser Ala Ala Ser Thr Ala Leu Ala Ala Val Gly  
 450 455 460

Ala Lys Leu Ser Glu Val Thr Leu Glu Thr Cys Lys Lys Ala Ala Glu  
 465 470 475 480

Ala Ala Leu Asp Ala Glu Gly Ala Thr Glu Ala Arg Asp Ala Val Arg  
 485 490 495

Ala Val Ile Asp Ala Ala Val  
 500

<210> 21  
 <211> 390  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> [101]..(367)  
 <223> RXA01300

<400> 21  
 gatgacatt aaatccctc ccttgggggg tttactaac aaatcgtgc gccctaacc 60  
 gttaggatta acggcgtagc aacacqaaag gacactttcc atg gcl. tcc aag accl 115  
 Met Ala Ser Lys Thr  
 1 5

gta acc gtc ggt tcc tcc gtt ggc ctg cac gca cgt cca gca tcc atc 163  
 Val Thr Val Gly Ser Val Gly Leu His Ala Arg Pro Ala Ser Ile  
 10 15 20

atc gct gaa gcg gct gct gag tac gac gac gaa atc ttg ctg acc ctg 211  
 Ile Ala Glu Ala Ala Ala Glu Tyr Asp Asp Glu Ile Leu Leu Thr Leu  
 25 30 35

gtt ggc tcc gat gat gac gaa gag acc gac gcg tcc tct tcc ctg atg 259  
 Val Gly Ser Asp Asp Asp Glu Glu Thr Asp Ala Ser Ser Ser Leu Met  
 40 45 50

atc atg gcg ctg ggc gca gag cac ggc aac gaa gtt acc gtc acc tcc 307  
 Ile Met Ala Leu Gly Ala Glu His Gly Asn Glu Val Thr Val Thr Ser  
 55 60 65

gac aac gct gaa gct gtt gag aag atc gct gcg att atc gca cag gac 355  
 Asp Asn Ala Glu Ala Val Glu Lys Ile Ala Ala Leu Ile Ala Cln Asp  
 70 75 80 85

att gac gct gag taaacaacgc tctgcttgtt aaa 390  
 Leu Asp Ala Glu

<210> 22  
 <211> 89  
 <212> PRT  
 <213> Corynebacterium glutamicum

&lt;400&gt; 22

Met Ala Ser Lys Thr Val Thr Val Gly Ser Ser Val Gly Leu His Ala  
 1 5 10 15  
 Arg Pro Ala Ser Ile Il Ala Glu Ala Ala Glu Tyr Asp Asp Glu  
 20 25 30  
 Ile Leu Leu Thr Leu Val Gly Ser Asp Asp Asp Glu Glu Thr Asp Ala  
 35 40 45  
 Ser Ser Ser Leu Met Ile Met Ala Leu Gly Ala Glu His Gly Asn Glu  
 50 55 60  
 Val Thr Val Thr Ser Asp Asn Ala Glu Ala Val Glu Lys Ile Ala Ala  
 65 70 75 80  
 Leu Ile Ala Gln Asp Leu Asp Ala Glu  
 85

&lt;210&gt; 23

&lt;211&gt; 508

&lt;212&gt; DNA

&lt;213&gt; Corynebacterium glutamicum

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (101)..(508)

&lt;223&gt; RXN03002

&lt;400&gt; 23

ggaaacttcga ggtgtctctcg tgggggctac ggagatctat caagctgtacg lllslqlkky 60  
 acctatccg aatcaacatg cagtgaatta acatctactt atg ttt gta ctc aac 115  
 Met Phe Val Leu Lys  
 1 5  
 gat ctg cta aag gca gaa cgc ata gaa ctc gac cgc acg gtc acc gat 163  
 Asp Leu Leu Lys Ala Glu Arg Ile Glu Leu Asp Arg Thr Val Thr Asp  
 10 15 20  
 tgg cgt gaa ggc atc cgc gcc gca ggt gta ctc cta gaa acg acc acc 211  
 Trp Arg Glu Gly Ile Arg Ala Ala Gly Val Leu Leu Glu Lys Thr Asn  
 25 30 35  
 agc att gat tcc gcc tac acc gat gcc atg atc gcc agc gtc gaa gaa 259  
 Ser Ile Asp Ser Ala Tyr Thr Asp Ala Met Ile Ala Ser Val Glu Glu  
 40 45 50  
 aaa ggc ccc tac att gtg gtc gct cca ggt ttc gct ttc gag cac gcc 307  
 Lys Gly Pro Tyr Ile Val Val Ala Pro Gly Phe Ala Phe Ala His Ala  
 55 60 65  
 cgc ccc agc aga gca gtc cgc gag acc gct atg tcg tgg gtg cgc ctg 355  
 Arg Pro Ser Arg Ala Val Arg Glu Thr Ala Met Ser Trp Val Arg Leu  
 70 75 80 85  
 gcc tcc cct gtt tcc ttc ggt cac aat aag aal gct ccc ctc aat ctc 403  
 Ala Ser Pro Val Ser Phe Gly His Ser Lys Asn Asp Pro Leu Asn Leu  
 90 95 100

atc gtt gct ctc gct gcc aaa gat gcc acc gca cgt acc caa gcg atg 451  
 ile Val Ala Leu Ala Ala Lys Asp Ala Thr Ala His Thr Gln Ala Met  
                   105                  110                  115

gcg gca ttg gct aaa gct tta gga aaa tac cga aag gat ctc gac gag 499  
 Ala Ala Leu Ala Lys Ala Leu Gly Lys Tyr Arg Lys Asp Leu Asp Glu  
                   120                  125                  130

gca caa agt  
 Ala Gln Ser  
                   135

508

<210> 24  
 <211> 136  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 24  
 Met Phe Val Leu Lys Asp Leu Leu Lys Ala Glu Arg Ile Glu Leu Asp  
   1                  5                  10                  15

Arg Thr Val Thr Asp Trp Arg Glu Gly Ile Arg Ala Ala Gly Val Leu  
                   20                  25                  30

Leu Glu Lys Thr Asn Ser Ile Asp Ser Ala Tyr Thr Asp Ala Met Ile  
                   35                  40                  45

Ala Ser Val Glu Gln Lys Gly Pro Tyr Ile Val Val Ala Pro Gly Phe  
                   50                  55                  60

Ala Phe Ala His Ala Arg Pro Ser Arg Ala Val Arg Glu Thr Ala Met  
                   65                  70                  75                  80

Ser Trp Val Arg Leu Ala Ser Pro Val Ser Phe Gly His Ser Lys Asn  
                   85                  90                  95

Asp Pro Leu Asn Leu Ile Val Ala Leu Ala Ala Lys Asp Ala Thr Ala  
                   100                  105                  110

His Thr Gln Ala Met Ala Ala Leu Ala Lys Ala Leu Gly Lys Tyr Arg  
                   115                  120                  125

Lys Asp Leu Asp Glu Ala Gln Ser  
                   130                  135

<210> 25  
 <211> 789  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> (34)..(766)  
 <223> RXC00953

<400> 25  
 ctgcattcc ccaatg gcg cca cca acg gta ggc aac tac atc atg cag tcc 52  
                   Met Ala Pro Pro Thr Val Gly Asn Tyr Ile Met Gln Ser  
                   1                  5                  10

ttc act cca ggt ctg cag ttc ggc gtt gca gtt gcc gtg att ctc tct	100
Phe Thr Gln Gly Leu Gln Phe Gly Val Ala Val Ala Val Ile Leu Phe	
15 20 25	
ggg gtc cgc acc att ctt ggt gna ctg gtc ccc gca ttc caa ggt att	148
Gly Val Arg Thr Ile Leu Gly Glu Leu Val Pro Ala Phe Gln Gly Ile	
30 35 40 45	
gct gcg aag gtt gtt ccc gga got atc ccc gca ttg gat gca ccg atc	196
Ala Ala Lys Val Val Pro Gly Ala Ile Pro Ala Leu Asp Ala Pro Ile	
50 55 60	
gtg ttc ccc tac gcg cag aac gcc gtt ctc att ggt ttc ttg tct tcc	244
Val Phe Pro Tyr Ala Gln Asn Ala Val Leu Ile Gly Phe Leu Ser Ser	
65 70 75	
ttc gtc ggt ggc ttg gtt ggc ctg act gtt ctt gca tgc tgg ctg aac	292
Phe Val Gly Gly Leu Val Gly Leu Thr Val Leu Ala Ser Trp Leu Asn	
80 85 90	
cca got ttt ggt gtc gcg ttg att ctg cct ggt ttg gtc ccc cac ttc	340
Pro Ala Phe Gly Val Ala Leu Ile Leu Pro Gly Leu Val Pro His Phe	
95 100 105	
ttc act ggt ggc gcg gcg ggc gtt tac ggt aat gcc acg ggt ggt cgt	388
Phe Thr Gly Gly Ala Ala Gly Val Tyr Gly Asn Ala Thr Gly Gly Arg	
110 115 120 125	
aga gga gca gta ttt ggc gcc ttt gcc aac ggt ctt ctg att acc ttc	436
Arg Gly Ala Val Phe Gly Ala Phe Ala Asn Gly Leu Leu Ile Thr Phe	
130 135 140	
ctc cct gct ttc ctg ctt ggt gta ctt ggt tcc ttc ggt tca gag aac	484
Leu Pro Ala Phe Leu Leu Gly Val Leu Gly Ser Phe Gly Ser Glu Asn	
145 150 155	
acc act ttc ggt gat gcg gac ttt ggt tgg ttc gga atc gtt gtt ggt	532
Thr Thr Phe Gly Asp Ala Asp Phe Gly Trp Phe Gly Ile Val Val Gly	
160 165 170	
tct gca gcc aag gtg gaa ggt gct ggc ggg ctc atc ttg ttg ctc atc	580
Ser Ala Ala Lys Val Glu Gly Ala Gly Gly Leu Ile Leu Leu Leu Ile	
175 180 185	
atc gca gcg gtt ctt ctg ggt ggc gcg atg gtc ttc cag aag cgc gtc	628
Ile Ala Ala Val Leu Leu Gly Gly Ala Met Val Phe Gln Lys Arg Val	
190 195 200 205	
gtg aat ggg cac tgg gat cca gct ccc aac cgt gag cgc gtg gag aag	676
Val Asn Gly His Trp Asp Pro Ala Pro Asn Arg Glu Arg Val Glu Lys	
210 215 220	
gcg gaa gct gat gcc act cca acg gct ggg gct cgg acc tac cct aag	724
Ala Glu Ala Asp Ala Thr Pro Thr Ala Gly Ala Arg Thr Tyr Pro Lys	
225 230 235	
att gct cct ccg gcg ggc gct cct acc cca ccg gct cga agc	766
Ile Ala Pro Pro Ala Gly Ala Pro Thr Pro Pro Ala Arg Ser	
240 245 250	

taagatctcc aaaaacctga gat.

789

<210> 26  
 <211> 251  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 26  
 Met Ala Pro Pro Thr Val Gly Asn Tyr Ile Met Gln Ser Phe Thr Gln  
 1 5 10 15  
 Gly Leu Gln Phe Gly Val Ala Val Ala Val Ile Leu Phe Gly Val Arg  
 20 25 30  
 Thr Ile Leu Gly Glu Leu Val Pro Ala Phe Gln Gly Ile Ala Ala Lys  
 35 40 45  
 Val Val Pro Gly Ala Ile Pro Ala Leu Asp Ala Pro Ile Val Phe Pro  
 50 55 60  
 Tyr Ala Gln Asn Ala Val Leu Ile Gly Phe Leu Ser Ser Phe Val Gly  
 65 70 75 80  
 Gly Leu Val Gly Leu Thr Val Leu Ala Ser Trp Leu Asn Pro Ala Phe  
 85 90 95  
 Gly Val Ala Leu Ile Leu Pro Gly Leu Val Pro His Phe Phe Thr Gly  
 100 105 110  
 Gly Ala Ala Gly Val Tyr Gly Asn Ala Thr Gly Gly Arg Arg Gly Ala  
 115 120 125  
 Val Phe Gly Ala Phe Ala Asn Gly Leu Leu Ile Thr Phe Leu Pro Ala  
 130 135 140  
 Phe Leu Leu Gly Val Leu Gly Ser Phe Gly Ser Glu Asn Thr Thr Phe  
 145 150 155 160  
 Gly Asp Ala Asp Phe Gly Trp Phe Gly Ile Val Val Gly Ser Ala Ala  
 165 170 175  
 Lys Val Glu Gly Ala Gly Gly Leu Ile Leu Leu Ile Ile Ala Ala  
 180 185 190  
 Val Leu Leu Gly Gly Ala Met Val Phe Gln Lys Arg Val Val Asn Gly  
 195 200 205  
 His Trp Asp Pro Ala Pro Asn Arg Glu Arg Val Glu Lys Ala Glu Ala  
 210 215 220  
 Asp Ala Thr Pro Thr Ala Gly Ala Arg Thr Tyr Pro Lys Ile Ala Pro  
 225 230 235 240  
 Pro Ala Gly Ala Pro Thr Pro Pro Ala Arg Ser  
 245 250

<210> 27  
 <211> 553  
 <212> DNA

&lt;213&gt; Corynebacterium glutamicum

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (101)..(553)

&lt;223&gt; RXCU3001

&lt;400&gt; 27

cccggttcac gtgatcaatg acttcaacgag caacgatgaa atogatgctg cgettcgtga 60

acgetaacgac atctaactac tttaaaagga cgaatatatt atg gac tgg tta acc 115  
Met Asp Trp Leu Thr  
1 5

att cct ctt ttc ctc gtt aat gaa atc att gag gtt ccg ggt ttc ctc 163  
Ile Pro Leu Phe Leu Val Asn Glu Ile Leu Ala Val Pro Ala Phe Leu  
10 15 20

atc ggt atc atc acc ggc gtn gga ttg ggt gcc atg ggg cgt tcc gtc 211  
Ile Gly Ile Ile Thr Ala Val Gly Leu Gly Ala Met Gly Arg Ser Val  
25 30 35

ggt cag gtt atc ggt gga gca atc aaa gca acg ttg ggc ttt ttg ctc 259  
Gly Gln Val Ile Gly Gly Ala Ile Lys Ala Thr Leu Gly Phe Leu Leu  
40 45 50

att ggt gag ggt gcc acg ttg gtc act gcc tcc ctg gag cca ctg ggt 307  
Ile Gly Ala Gly Ala Thr Leu Val Thr Ala Ser Leu Glu Pro Leu Gly  
55 60 65

gag atg atc atg ggt gcc aca ggc atg cgt ggt gtt gtc cca acg aat 355  
Ala Met Ile Met Gly Ala Thr Gly Met Arg Gly Val Val Pro Thr Asn  
70 75 80 85

gaa gcc atc gcc gga atc gca cag gct gaa tac ggc gag cag gtg gcg 403  
Glu Ala Thr Ala Gly Ile Ala Gln Ala Glu Tyr Gly Ala Gln Val Ala  
90 95 100

tgg ctg atg att ctg ggc ttc gcc atc tot ttg gtg ttg got cgt ttc 451  
Trp Leu Met Ile Leu Gly Phe Ala Ile Ser Leu Val Leu Ala Arg Phe  
105 110 115

acc acc ctg cgt tat gtc ttg ctc aac gga cac cac gtg ctg ttg atg 499  
Thr Asn Leu Arg Tyr Val Leu Leu Asn Gly His His Val Leu Leu Met  
120 125 130

tgc acc atg ctc acc atg gtc ttg gcc acc gga aga gtt gat gag tgg 517  
Cys Thr Met Leu Thr Met Val Leu Ala Thr Gly Arg Val Asp Ala Trp  
135 140 145

atc ttc  
Ile Phe 553  
150

&lt;210&gt; 28

&lt;211&gt; 151

&lt;212&gt; PRT

&lt;213&gt; Corynebacterium glutamicum

&lt;400&gt; 28

Met Asp Trp Leu Thr Il Pro Leu Phe Leu Val Asn Glu Ile Leu Ala  
 1 5 10 15  
 Val Pro Ala Phe Leu Ile Gly Ile Ile Thr Ala Val Gly Leu Gly Ala  
 20 25 30  
 Met Gly Arg Ser Val Gly Gln Val Ile Gly Gly Ala Ile Lys Ala Thr  
 35 40 45  
 Leu Gly Phe Leu Leu Ile Gly Ala Gly Ala Thr Leu Val Thr Ala Ser  
 50 55 60  
 Leu Glu Pro Leu Gly Ala Met Ile Met Gly Ala Thr Gly Met Arg Gly  
 65 70 75 80  
 Val Val Pro Thr Asn Glu Ala Ile Ala Gly Ile Ala Gln Ala Glu Tyr  
 85 90 95  
 Gly Ala Gln Val Ala Trp Leu Met Ile Leu Gly Phe Ala Ile Ser Leu  
 100 105 110  
 Val Leu Ala Arg Phe Thr Asn Leu Arg Tyr Val Leu Leu Asn Gly His  
 115 120 125  
 His Val Leu Leu Met Cys Thr Met Leu Thr Met Val Leu Ala Thr Gly  
 130 135 140  
 Arg Val Asp Ala Trp Ile Phe  
 145 150

<210> 29  
 <211> 2172  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> [101]..(2149)  
 <223> RXN01943

<400> 29  
 cagattcttt ttccgcccac ttccgtacgg ccatctcttt aagtgacac gaaagtctct 60  
 tgcctcgagg agacagaccc tacgtttaga aaggtttgac atg gcg tcc aaa ctg 115  
 Met Ala Ser Lys Leu 5  
 acg acg aca tcg caa cat att ctg gaa aac att ggt gga cca gac aat 163  
 Thr Thr Thr Ser Gln His Ile Leu Glu Asn Leu Gly Gly Pro Asp Asn 20  
 10  
 att act tcg atg act cac tgt gcg act cgc att cgc ttc caa gtg aag 211  
 Ile Thr Ser Met Thr His Cys Ala Thr Arg Leu Arg Phe Gln Val Lys 35  
 25  
 gat caa tcc att gtt gal caa caa gaa att gaa tcc gac caa tca gtt 259  
 Asp Gln Ser Ile Val Asp Gln Gln Glu Ile Asp Ser Asp Pro Ser Val 50  
 40 45  
 ctt gcc gta gta ccc caa gga tcc acc ggt atg cag gta gtg atg ggt 307

Leu Gly Val Val Pro Gln Gly Ser Thr Gly Met Gln Val Val Met Gly  
55 60 65

gga tct gtt gca aac tat tac caa gaa atc ctc aac ctt gat gga atg 355  
Gly Ser Val Ala Asn Tyr Tyr Gln Glu Ile Leu Lys Leu Asp Gly Met  
70 75 80 85

aag cac ttc gcc gac ggt gaa gct aca gag agt tca tcc aag aag gaa 403  
Lys His Phe Ala Asp Gly Glu Ala Thr Glu Ser Ser Ser Lys Lys Glu  
90 95 100

tac ggc gga gtc cgt gcc aag tcc tcc tgg att gac tac gcc ttc gag 451  
Tyr Gly Gly Val Arg Gly Lys Tyr Ser Trp Ile Asp Tyr Ala Phe Glu  
105 110 115

ttc ttg tct gat act ttc cga cca atc ctg tgg gcc ctg ctt ggt gcc 499  
Phe Leu Ser Asp Thr Phe Arg Pro Ile Leu Trp Ala Leu Leu Gly Ala  
120 125 130

tca ctg att att acc ttg ttg gtt ctt gcg gat act ttc ggt ttg caa 547  
Ser Leu Ile Ile Thr Leu Leu Val Leu Ala Asp Thr Phe Gly Leu Gln  
135 140 145

gac ttc cgc gct cca atg gct gaa cag cct gat act tat gta ttc ctg 595  
Asp Phe Arg Ala Pro Met Asp Glu Gln Pro Asp Thr Tyr Val Phe Leu  
150 155 160 165

cac tcc atg tgg cgc tgg gtc ttc tac ttc ctg cca att ctg gtt ggt 643  
His Ser Met Trp Arg Ser Val Phe Tyr Phe Leu Pro Ile Met Val Gly  
170 175 180

gcc acc gca gct cga aag ctc gcc gca aac gag tgg att ggt gca gct 691  
Ala Thr Ala Ala Arg Lys Leu Gly Ala Asn Glu Trp Ile Gly Ala Ala  
185 190 195

att cca gcc gca ctt ctt act cca gaa ttc ttg gca ctg ggt tct gcc 739  
Ile Pro Ala Ala Leu Leu Thr Pro Glu Phe Leu Ala Leu Gly Ser Ala  
200 205 210

ggc gac acc gtc aca gtc ttt gcc ctg cca atg gtt ctg aat gac tac 787  
Gly Asp Thr Val Thr Val Phe Gly Leu Pro Met Val Leu Asn Asp Tyr  
215 220 225

tcc gcc cag gta ttc cca ccg ctg att gca gca att ggt ctg tac tgg 835  
Ser Gly Gln Val Phe Pro Pro Leu Ile Ala Ala Ile Gly Leu Tyr Trp  
230 235 240 245

gtg gaa aag gga ctg aag aag atc atc cct gaa gca gtc caa atg gtg 883  
Val Glu Lys Gly Leu Lys Lys Ile Ile Pro Glu Ala Val Gln Met Val  
250 255 260

ttc gtc cca ttc ttc tcc ctg ctg att atg atc cca gcg acc gca ttc 931  
Phe Val Pro Phe Phe Ser Leu Leu Ile Met Ile Pro Ala Thr Ala Phe  
265 270 275

ctg ctt gga cct ttc gcc atc ggt gll ggt aac gga att tcc aac ctg 979  
Leu Leu Gly Pro Phe Gly Ile Gly Val Gly Asn Gly Ile Ser Asn Leu  
280 285 290

ctt gaa gcg att aac aac ttc agc cca ttt att ctt tcc atc gtt atc 1027  
Leu Glu Ala Ile Asn Asn Phe Ser Pro Phe Ile Leu Ser Ile Val Ile



295	300	305	
cca ttg ctc tac cca ttc ttg gtt cca ctt gga ttg cac tgg cca ctc Pro Leu Leu Tyr Pro Phe Leu Val Pro Leu Gly Leu His Trp Pro Leu 310 315 320 325			1075
aac gcc atc atg atc cag aac atc aac acc ctg ggt tac gcc ttc att Asn Ala Ile Met Ile Gln Asn Ile Asn Thr Leu Gly Tyr Asp Phe Ile 330 335 340			1123
cag gga cca atg ggt gcc tgg aac ttc gcc tgc ttc ggc ctg gtc acc Gln Gly Pro Met Gly Ala Trp Asn Phe Ala Cys Phe Gly Leu Val Thr 345 350 355			1171
ggc qtg ttc ttg ctc tcc att aag gaa cga aac aag gcc atg cgt cag Gly Val Phe Leu Leu Ser Ile Lys Glu Arg Asn Lys Ala Met Arg Gln 360 365 370			1219
gtt tcc ctg ggt gcc atg ttg gct ggt ttg ctc gcc gcc att tcc gag Val Ser Leu Gly Gly Met Leu Ala Gly Leu Leu Gly Gly Ile Ser Glu 375 380 385			1267
cct tcc ctc tac ggt gtt ctg ctc cga ttc aag aag acc tac ttc cgc Pro Ser Leu Tyr Gly Val Leu Leu Arg Phe Lys Lys Thr Tyr Phe Arg 390 395 400 405			1315
ctc ctg acc ggt tgt ttg gca gcc ggt atc gtg atg gcc atc ttc gcc Leu Leu Pro Gly Cys Leu Ala Gly Gly Ile Val Met Gly Ile Phe Asp 410 415 420			1363
atc aag gcc tac gct ttc gtg ttc acc tcc ttg ctt acc atc cca gca Ile Lys Ala Tyr Ala Phe Val Phe Thr Ser Leu Leu Thr Ile Pro Ala 425 430 435			1411
atg gac cca tgg ttg gcc tac acc att ggt atc gca gtt gca ttc ttc Met Asp Pro Trp Leu Gly Tyr Thr Ile Gly Ile Ala Val Ala Phe Phe 440 445 450			1459
gtt tcc atg ttc ctt gtt ctc gca ctg gac tac cgt tcc aac gaa gag Val Ser Met Phe Leu Val Leu Ala Leu Asp Tyr Arg Ser Asn Glu Glu 455 460 465			1507
cgc gat gag gca cgt gca aag gtt gct gct gac aag cag gca gaa gaa Arg Asp Glu Ala Arg Ala Lys Val Ala Ala Asp Lys Gln Ala Glu Glu 470 475 480 485			1555
gat ctg aag gca gaa gct aat gca act cct gca gct cca gta gct gct Asp Leu Lys Ala Glu Ala Asn Ala Thr Pro Ala Ala Pro Val Ala Ala 490 495 500			1603
gca ggt gcc gga gcc ggt gca ggt gca gga gcc gct gct gcc gct gca Ala Gly Ala Gly Ala Gly Ala Gly Ala Gly Ala Ala Ala Gly Ala Ala 505 510 515			1651
acc gcc ctg gca gct aag ccg aag ctg gcc gct ggg gaa gta ctg gac Thr Ala Val Ala Ala Lys Pro Lys Leu Ala Ala Gly Glu Val Val Asp 520 525 530			1699
att gtt tcc cca ctc gaa gcc aag gca att cca ctt tot gaa gta cct Tle Val Ser Pro Leu Glu Gly Lys Ala Tle Pro Leu Ser Glu Val Pro 535 540 545			1747

gac cca atc ttt gca gca ggc aag ctt gga cca ggc att gca atc caa 1795  
 Asp Pro Ile Phe Ala Ala Gly Lys Leu Gly Pro Gly Ile Ala Ile Cln  
 550 555 560 565  
 cca act gga aac acc gtt gtt gct cca gca gac gct act gtc atc ctt 1843  
 Pro Thr Gly Asn Thr Val Val Ala Pro Ala Asp Ala Thr Val Ile Leu  
 570 575 580  
 gtc cag aaa tct gga cac gca gtg gca ttg cgc tta gat agc gga gtt 1891  
 Val Cln Lys Ser Gly His Ala Val Ala Leu Arg Leu Asp Ser Gly Val  
 585 590 595  
 gaa atc ctt gtc cac gtt gga ttg gac acc glg caa ttg ggc ggc gaa 1939  
 Glu Ile Leu Val His Val Gly Leu Asp Thr Val Gln Leu Gly Gly Glu  
 600 605 610  
 ggc ttc acc gtt cac gtt gag cgc agg cag caa gtc aag ggc ggc gat 1987  
 Gly Phe Thr Val His Val Glu Arg Arg Gln Gln Val Lys Ala Gly Asp  
 615 620 625  
 cca ctg atc act ttt gac gct gac ttc att cga tcc aag gat cta cct 2035  
 Pro Leu Ile Thr Phe Asp Ala Asp Phe Ile Arg Ser Lys Asp Leu Pro  
 630 635 640 645  
 ttg atc acc cca gtt gtg gtg tct aac gcc gcg aaa ttc ggt gaa att 2083  
 Leu Ile Thr Pro Val Val Val Ser Asn Ala Ala Lys Phe Gly Glu Ile  
 650 655 660  
 gaa ggt att cct gca gat cag gca aat tct tcc acg act gtc atc aag 2131  
 Glu Gly Ile Pro Ala Asp Gln Ala Asn Ser Ser Thr Thr Val Ile Lys  
 665 670 675  
 gtc aac ggc aag aac gag taacctggga tccatgttgc gca 2172  
 Val Asn Gly Lys Asn Glu  
 680

<210> 30  
 <211> 683  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 30  
 Met Ala Ser Lys Leu Thr Thr Thr Ser Gln His Ile Leu Glu Asn Leu  
 1 5 10 15  
 Gly Gly Pro Asp Asn Ile Thr Ser Met Thr His Cys Ala Thr Arg Leu  
 20 25 30  
 Arg Phe Gln Val Lys Asp Gln Ser Ile Val Asp Gln Gln Glu Ile Asp  
 35 40 45  
 Ser Asp Pro Ser Val Leu Gly Val Val Pro Gln Gly Ser Thr Gly Met  
 50 55 60  
 Gln Val Val Met Gly Gly Ser Val Ala Asn Tyr Tyr Gln Gln Ile Leu  
 65 70 75 80  
 Lys Leu Asp Gly Met Lys His Phe Ala Asp Gly Glu Ala Thr Glu Ser  
 85 90 95

S r Ser Lys Lys Glu Tyr Gly Gly Val Arg Gly Lys Tyr Ser Trp Ile  
 100 105 110  
 Asp Tyr Ala Phe Glu Phe Leu Ser Asp Thr Phe Arg Pro Ile Leu Trp  
 115 120 125  
 Ala Leu Leu Gly Ala Ser Leu Ile Ile Thr Leu Leu Val Leu Ala Asp  
 130 135 140  
 Thr Phe Gly Leu Gln Asp Phe Arg Ala Pro Met Asp Glu Gln Pro Asp  
 145 150 155 160  
 Thr Tyr Val Phe Leu His Ser Met Trp Arg Ser Val Phe Tyr Phe Leu  
 165 170 175  
 Pro Ile Met Val Gly Ala Thr Ala Ala Arg Lys Leu Gly Ala Asn Glu  
 180 185 190  
 Trp Ile Gly Ala Ala Ile Pro Ala Ala Leu Leu Thr Pro Glu Phe Leu  
 195 200 205  
 Ala Leu Gly Ser Ala Gly Asp Thr Val Thr Val Phe Gly Leu Pro Met  
 210 215 220  
 Val Leu Asn Asp Tyr Ser Gly Gln Val Phe Pro Pro Leu Ile Ala Ala  
 225 230 235 240  
 Ile Gly Leu Tyr Trp Val Glu Lys Gly Leu Lys Lys Ile Ile Pro Glu  
 245 250 255  
 Ala Val Gln Met Val Phe Val Pro Phe Phe Ser Leu Leu Ile Met Ile  
 260 265 270  
 Pro Ala Thr Ala Phe Leu Leu Gly Pro Phe Gly Ile Gly Val Gly Asn  
 275 280 285  
 Gly Ile Ser Asn Leu Leu Glu Ala Ile Asn Asn Phe Ser Pro Phe Ile  
 290 295 300  
 Leu Ser Ile Val Ile Pro Leu Leu Tyr Pro Phe Leu Val Pro Leu Gly  
 305 310 315 320  
 Leu His Trp Pro Leu Asn Ala Ile Met Ile Gln Asn Ile Asn Thr Leu  
 325 330 335  
 Gly Tyr Asp Phe Ile Gln Gly Pro Met Gly Ala Trp Asn Phe Ala Cys  
 340 345 350  
 Phe Gly Leu Val Thr Gly Val Phe Leu Leu Ser Ile Lys Glu Arg Asn  
 355 360 365  
 Lys Ala Met Arg Gln Val Ser Leu Gly Gly Met Leu Ala Gly Leu Leu  
 370 375 380  
 Gly Gly Ile Ser Glu Pro Ser Leu Tyr Gly Val Leu Leu Arg Phe Lys  
 385 390 395 400  
 Lys Thr Tyr Phe Arg Leu Leu Pro Gly Cys Leu Ala Gly Gly Ile Val  
 405 410 415

Met Gly Ile Phe Asp Ile Lys Ala Tyr Ala Phe Val Phe Thr Ser Leu  
 420 425 430

Leu Thr Ile Pro Ala Met Asp Pro Trp Leu Gly Tyr Thr Ile Gly Ile  
 435 440 445

Ala Val Ala Phe Phe Val Ser Met Phe Leu Val Leu Ala Leu Asp Tyr  
 450 455 460

Arg Ser Asn Glu Glu Arg Asp Glu Ala Arg Ala Lys Val Ala Ala Asp  
 465 470 475 480

Lys Gln Ala Glu Glu Asp Leu Lys Ala Glu Ala Asn Ala Thr Pro Ala  
 485 490 495

Ala Pro Val Ala Ala Ala Gly Ala Gly Ala Gly Ala Gly Ala  
 500 505 510

Ala Ala Gly Ala Ala Thr Ala Val Ala Ala Lys Pro Lys Leu Ala Ala  
 515 520 525

Gly Glu Val Val Asp Ile Val Ser Pro Leu Glu Gly Lys Ala Ile Pro  
 530 535 540

Leu Ser Glu Val Pro Asp Pro Ile Phe Ala Ala Gly Lys Leu Gly Pro  
 545 550 555 560

Gly Ile Ala Ile Gln Pro Thr Gly Asn Thr Val Val Ala Pro Ala Asp  
 565 570 575

Ala Thr Val Ile Leu Val Gln Lys Ser Gly His Ala Val Ala Leu Arg  
 580 585 590

Leu Asp Ser Gly Val Glu Ile Leu Val His Val Gly Leu Asp Thr Val  
 595 600 605

Gln Leu Gly Gly Glu Gly Phe Thr Val His Val Glu Arg Arg Gln Gln  
 610 615 620

Val Lys Ala Gly Asp Pro Leu Ile Thr Phe Asp Ala Asp Phe Ile Arg  
 625 630 635 640

S r Lys Asp Leu Pro Leu Ile Thr Pro Val Val Val Ser Asn Ala Ala  
 645 650 655

Lys Phe Gly Glu Ile Glu Gly Ile Pro Ala Asp Gln Ala Asn Ser Ser  
 660 665 670

Thr Thr Val Ile Lys Val Asn Gly Lys Asn Glu  
 675 680

<210> 31  
 <211> 1339  
 <212> DNA  
 <213> Corynebacterium glutamicum

<220>  
 <221> CDS  
 <222> {1011..(1339)}  
 <223> FRXA02191

&lt;100&gt; 31

ccgattcttt ttcgcccaaa ttctaacgg cgtctctctt aagtggacaa gaaagtctct 60

tgcacgcggg agacagacc tacgtttaga aaggtttgac atg ggc tcc aaa ctg 115  
Met Ala Ser Lys L u  
1 5

acg acg acc tgg caa cat att ctg gaa aac att ggt gga cca gac aat 163  
Thr Thr Thr Ser Gln His Ile Leu Glu Asn Leu Gly Gly Pro Asp Asn  
10 15 20

att act tgg atg act cac tgt ggg act cgc att cgc ttc caa gtg aag 211  
Ile Thr Ser Met Thr His Cys Ala Thr Arg Leu Arg Phe Gln Val Lys  
25 30 35

gat caa tcc att gtt gat caa caa gaa att gac tcc gac cca tca gtt 259  
Asp Gln Ser Ile Val Asp Gln Gln Glu Ile Asp Ser Asp Pro Ser Val  
40 45 50

ctt ggc gta gta ccc caa gga tcc acc ggt atg cag gtg gtg atg ggt 307  
 Leu Gly Val Val Pro Gln Gly Ser Thr Gly Met Gln Val Val Met Gly  
     55                    60                    65

gga tct gtt gca aac tat tac caa gaa atc ctc aac ctt ggt gga atg 355  
Gly Ser Val Ala Asn Tyr Tyr Gln Glu Ile Leu Lys Leu Asp Gly Met  
70 75 80 85

aag cac ttc gcc gac ggt gan got aca gag ayt tcc tcc aag aag gaa 403  
 Lys His Phe Ala Asp Gly Glu Ala Thr Glu Ser Ser Ser Lys Lys Glu  
 90 95 100

Lac ggc gga gln cgt ggc aag Lac Lcg Lgg att gac tac gcc ttc gag 451  
Tyr Gly Gly Val Arg Gly Lys Tyr Ser Trp Ile Asp Tyr Ala Phe Glu  
105 110 115

ttc ttg tct gat act ttc cga cca atc ctg tgg gcc ctg ctt ggt gcc 499  
 Phe Leu Ser Asp Thr Phe Arg Pro Ile Leu Trp Ala Leu Leu Gly Ala  
 120 125 130

tca ctg att att acc ttg ttg gtt ctt gcg gat act ttc ggt ttg caa 547  
Ser Leu Ile Ile Thr Leu Leu Val Leu Ala Asp Thr Phe Gly Leu Cln  
135 140 145

gac ttc cgc gct cca atg gat gag cag cct gat act tat gta ttc ctg 595  
Asp Phe Arg Ala Pro Met Asp Glu Gln Pro Asp Thr Tyr Val Phe Leu  
150 155 160 165

cac hcc atg tgg cgc tcg gtc ttc tac ttc ctg cca att atg gtt ggt      613  
His Ser Met Trp Arg Ser Val Phe Tyr Phe Leu Pro Ile Met Val Gly

170                  175                  180

gac gcc gca gct cga agc ctc ggc gca aac gaa tgg att ggt gca gct 691  
Ala Thr Ala Ala Arg Lys Leu Gly Ala Asn Glu Trp Ile Gly Ala Ala  
185 190 195

att coa gcc gca att att act coa gaa ttc ttg gca ctg ggt tct gcc 739  
Tle Pro Ala Leu Leu Thr Pro Glu Phe Leu Ala Leu Gly Ser Ala  
200 205 210

ggc gat acc gtc acc gtc ttt ggc ctg cca atg gtt ctg aat gac taa 787

Gly	Asp	Thr	Val	Thr	Val	Phe	Gly	Leu	Pro	Met	Val	L	u	Asn	Asp	Tyr	
215						220					225						
tcg	gga	cag	gta	ttc	cca	ccg	ctg	att	gca	gca	att	ggt	ctg	tac	tgg		835
S r	Gly	Gln	Val	Phe	Pro	Pro	Leu	Ile	Ala	Ala	Ile	Gly	Leu	Tyr	Trp		
230					235				240						245		
gtg	gaa	aag	gga	ctg	aag	aag	atc	atc	cct	gaa	gca	gtc	caa	atg	gtg		883
Val	Glu	Lys	Gly	Leu	Lys	Lys	Ile	Ile	Pro	Glu	Ala	Val	Gln	Met	Val		
				250					255					260			
ttc	gtc	cca	ttc	ttc	tcg	ctg	att	atg	atc	cca	gcg	acc	gca	ttc			931
Phe	Val	Pro	Phe	Phe	Ser	Leu	Leu	Ile	Met	Ile	Pro	Ala	Thr	Ala	Phe		
			265					270					275				
ctg	ctt	gga	cct	ttc	ggc	atc	ggc	gtt	ggc	acc	gga	att	tac	aac	ctg		979
Leu	Leu	Gly	Pro	Phe	Gly	Ile	Gly	Val	Gly	Asn	Gly	Ile	Ser	Asn	Leu		
		280					285					290					
ctt	gaa	gcg	att	aac	aac	ttc	ggc	cca	ttt	att	ctt	tcc	atc	gtt	atc		1027
Leu	Glu	Ala	Ile	Asn	Asn	Phe	Ser	Pro	Phe	Ile	Leu	Ser	Ile	Val	Ile		
	295					300					305						
cca	ttg	ctc	tac	cca	ttc	ttg	gtt	cca	ctt	gga	ttg	cac	tgg	cca	ctc		1075
Pro	Leu	Leu	Tyr	Pro	Phe	Leu	Val	Pro	Leu	Gly	Leu	His	Trp	Pro	Leu		
310					315					320					325		
aac	gcc	atc	atg	atc	cag	aac	atc	aac	acc	ctg	ggt	tac	gac	ttc	att		1123
Asn	Ala	Ile	Met	Ile	Gln	Asn	Ile	Asn	Thr	Leu	Gly	Tyr	Asp	Phe	Ile		
				330				335						340			
cag	gga	cca	atg	ggt	gcc	tgg	aac	ttc	gcc	tgc	ttc	ggc	ctg	gtc	acc		1171
Gln	Gly	Pro	Met	Gly	Ala	Trp	Asn	Phe	Ala	Cys	Phe	Gly	Leu	Val	Thr		
		345					350					355					
ggc	gtg	ttc	ttg	ctc	tcg	att	aag	gaa	cca	aac	ang	goc	atg	cgt	cag		1219
Gly	Val	Phe	Leu	Leu	Ser	Thr	Lys	Glu	Arg	Asn	Lys	Ala	Met	Arg	Gln		
	360						365					370					
gtt	tcc	ctg	ggt	ggc	atg	ttg	ggt	ggt	ttg	ctc	ggc	ggc	att	tcc	gag		1267
Val	Ser	Leu	Gly	Gly	Met	Leu	Ala	Gly	Leu	Leu	Gly	Gly	Ile	Ser	Glu		
	375					380					385						
cct	tcc	ctc	tac	ggt	gtt	ctg	ctc	cca	ttc	aag	aag	acc	tac	ttc	cgc		1315
Pro	Ser	Leu	Tyr	Gly	Val	Leu	Leu	Arg	Phe	Lys	Lys	Thr	Tyr	Phe	Arg		
390					395				400						405		
ctc	ctg	ccg	ggt	tgt	ttg	gca	gca										1339
Leu	Leu	Pro	Gly	Cys	Leu	Ala	Ala										
				410													

&lt;210&gt; 32

&lt;211&gt; 413

&lt;212&gt; PRT

&lt;213&gt; Corynebacterium glutamicum

&lt;400&gt; 32

Met	Ala	Ser	Lys	Leu	Thr	Thr	Thr	Ser	Gln	His	Ile	Leu	Glu	Asn	Leu	
1				5					10					15		

Gly Gly Pro Asp Asn Ile Thr Ser Met Thr His Cys Ala Thr Arg Leu  
 20 25 30  
 Arg Ph Gln Val Lys Asp Gln Ser Ile Val Asp Gln Gln Glu Ile Asp  
 35 40 45  
 Ser Asp Pro Ser Val Leu Gly Val Val Pro Gln Gly Ser Thr Gly Met  
 50 55 60  
 Gln Val Val Met Gly Gly Ser Val Ala Asn Tyr Tyr Gln Glu Ile Leu  
 65 70 75 80  
 Lys Leu Asp Gly Met Lys His Phe Ala Asp Gly Glu Ala Thr Glu Ser  
 85 90 95  
 Ser Ser Lys Lys Glu Tyr Gly Gly Val Arg Gly Lys Tyr Ser Trp Ile  
 100 105 110  
 Asp Tyr Ala Phe Glu Phe Leu Ser Asp Thr Phe Arg Pro Ile Leu Trp  
 115 120 125  
 Ala Leu Leu Gly Ala Ser Leu Ile Ile Thr Leu Leu Val Leu Ala Asp  
 130 135 140  
 Thr Phe Gly Leu Gln Asp Phe Arg Ala Pro Met Asp Glu Gln Pro Asp  
 145 150 155 160  
 Thr Tyr Val Phe Leu His Ser Met Trp Arg Ser Val Phe Tyr Phe Leu  
 165 170 175  
 Pro Ile Met Val Gly Ala Thr Ala Ala Arg Lys Leu Gly Ala Asn Glu  
 180 185 190  
 Trp Ile Gly Ala Ala Ile Pro Ala Ala Leu Leu Thr Pro Glu Phe Leu  
 195 200 205  
 Ala Leu Gly Ser Ala Gly Asp Thr Val Thr Val Phe Gly Leu Pro Met  
 210 215 220  
 Val Leu Asn Asp Tyr Ser Gly Gln Val Phe Pro Pro Leu Ile Ala Ala  
 225 230 235 240  
 Ile Gly Leu Tyr Trp Val Glu Lys Gly Leu Lys Lys Ile Ile Pro Glu  
 245 250 255  
 Ala Val Gln Met Val Phe Val Pro Phe Phe Ser Leu Leu Ile Met Ile  
 260 265 270  
 Pro Ala Thr Ala Phe Leu Leu Gly Pro Phe Gly Ile Gly Val Gly Asn  
 275 280 285  
 Gly Ile Ser Asn Leu Leu Glu Ala Ile Asn Asn Phe Ser Pro Phe Ile  
 290 295 300  
 Leu Ser Ile Val Ile Pro Leu Leu Tyr Pro Phe Leu Val Pro Leu Gly  
 305 310 315 320  
 Leu His Trp Pro Leu Asn Ala Ile Met Ile Gln Asn Ile Asn Thr L u  
 325 330 335  
 Gly Tyr Asp Phe Ile Gln Gly Pro Met Gly Ala Trp Asn Ph Ala Cys

340

345

350

Phe Gly Leu Val Thr Gly Val Phe Leu Leu Ser Ile Lys Glu Arg Asn  
 355 360 365

Lys Ala Met Arg Gln Val Ser Leu Gly Gly Met L u Ala Gly Leu L u  
 370 375 380

Gly Gly Ile Ser Glu Pro Ser Leu Tyr Gly Val Leu Leu Arg Phe Lys  
 385 390 395 400

Lys Thr Tyr Phe Arg Leu Leu Pro Gly Cys Leu Ala Ala  
 405 410

&lt;210&gt; 33

&lt;211&gt; 428

&lt;212&gt; DNA

&lt;213&gt; Corynebacterium glutamicum

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(405)

&lt;223&gt; FRXA01943

&lt;400&gt; 33

cct gac cca atc ttt gca gca ggc aag ctt gga cca ggc att gca atc 48  
 Pro Asp Pro Ile Phe Ala Ala Gly Lys Leu Gly Pro Gly Ile Ala Ile  
 1 5 10 15

caa cca not gga aac aac gtt gtt gct cca gca gac gct act gtc atc 96  
 Gln Pro Thr Gly Asn Thr Val Val Ala Pro Ala Asp Ala Thr Val Ile  
 20 25 30

ctt gtc cag aac tct gga cca gca gtc gca tta cga gat agc gga 144  
 Leu Val Gln Lys Ser Gly His Ala Val Ala Leu Arg Leu Asp Ser Gly  
 35 40 45

gtt gaa atc ctt gtc cac ggt gga ttc gac aac gtc cca ttc ggc ggc 192  
 Val Glu Ile Leu Val His Val Gly Leu Asp Thr Val Gln Leu Gly Gly  
 50 55 60

gaa ggc ttc aac gtt cac ggt gag cgc agg cag caa gtc aag ggc ggc 240  
 Glu Gly Phe Thr Val His Val Glu Arg Arg Gln Gln Val Lys Ala Gly  
 65 70 75 80

gac cca ctc atc act ttt gac gct gac ttc att cga tcc aag gat cta 288  
 Asp Pro Leu Ile Thr Phe Asp Ala Asp Phe Ile Arg Ser Lys Asp Leu  
 85 90 95

cct ttg atc acc cca gtt gtc gtc tct aac gcc ggc aac ttc ggt gaa 336  
 Pro Leu Ile Thr Pro Val Val Val Ser Asn Ala Ala Lys Phe Gly Glu  
 100 105 110

att gaa ggt att cct gca gat cag gca aat tct tcc aag act gtc atc 384  
 Ile Glu Gly Ile Pro Ala Asp Gln Ala Asn Ser Ser Thr Thr Val Ile  
 125 120 125

aag gtc aac ggc aag aac gag taacctggga tccatgttgc gca 428  
 Lys Val Asn Gly Lys Asn Glu  
 130 135



<210> 34  
 <211> 135  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 34  
 Pro Asp Pro Ile Phe Ala Ala Gly Lys Leu Gly Pro Gly Ile Ala Ile  
   1                  5                  10                  15  
 Gln Pro Thr Gly Asn Thr Val Val Ala Pro Ala Asp Ala Thr Val Ile  
           20                  25                  30  
 Leu Val Gln Lys Ser Gly His Ala Val Ala Leu Arg Leu Asp Ser Gly  
           35                  40                  45  
 Val Glu Ile Leu Val His Val Gly Leu Asp Thr Val Gln Leu Gly Gly  
           50                  55                  60  
 Glu Gly Phe Thr Val His Val Glu Arg Arg Gln Gln Val Lys Ala Gly  
           65                  70                  75                  80  
 Asp Pro Leu Ile Thr Phe Asp Ala Asp Phe Ile Arg Ser Lys Asp Leu  
                   85                  90                  95  
 Pro Leu Ile Thr Pro Val Val Val Ser Asn Ala Ala Lys Phe Gly Glu  
           100                  105                  110  
 Ile Gln Gly Ile Pro Ala Asp Gln Ala Asn Ser Ser Thr Thr Val Ile  
           115                  120                  125  
 Lys Val Asn Gly Lys Asn Glu  
           130                  135

THIS PAGE BLANK (USPTO)